

**PUTRIZEN ARNICA BRUISE CREAM- glycerin cream**  
**Shantou South Moon Biotechnology Co., Ltd.**

-----  
MENTHA SUAVEOLENS LEAF □ PANTHENOL □ ARNICA MONTANA FLOWER □ COCONUT OIL □ MENAQUINONE 7 □ ASCORBIC ACID □ TOCOPHEROL

For the temporary relief of pain, swelling, and discoloration due to minor bruising · muscle overexertion · or skin discomfort.

- 1.Clean and dry the bruise area.
- 2.Apply a small amount of cream directly onto the bruise or affected area.
- 3.Gently massage in upward circular motions until fully absorbed.
- 4.Pay special attention to fragile or delicate skin areas.
- 5.Use 2-3 times daily or as needed.

For external use only.

Do not use if you are allergic to Arnica Montana or any of the ingredients.

Avoid contact with eyes, mucous membranes, or broken skin.

Stop use and ask a doctor if condition worsens or symptoms persist for more than 3 days.

Keep out of reach of children. In case of ingestion, seek medical help or contact a Poison Control Center immediately.

Do not use if you are allergic to Arnica Montana or any of the ingredients.

Stop use and ask a doctor if condition worsens or symptoms persist for more than 3 days.

Keep out of reach of children.

Store in a cool and dry place.

WATER □ GLYCERIN □ MINERAL OIL □ GLYCERYL MONOSTEARATE □ 1,2-HEXANEDIOL □ CARBOMER HOMOPOLYMER TYPE A □ TROLAMINE □ ALLANTOIN

- 1.Clean and dry the bruise area.
- 2.Apply a small amount of cream directly onto the bruise or affected area.
- 3.Gently massage in upward circular motions until fully absorbed.
- 4.Pay special attention to fragile or delicate skin areas.
- 5.Use 2-3 times daily or as needed.

- 1.Clean and dry the bruise area.
- 2.Apply a small amount of cream directly onto the bruise or affected area.
- 3.Gently massage in upward circular motions until fully absorbed.
- 4.Pay special attention to fragile or delicate skin areas.
- 5.Use 2-3 times daily or as needed.

# 纸盒 (哑膜)

规格尺寸: 长7.2x宽7.2x高5.1cm



PUTRIZEN ARNICA BRUISE CREAM			
glycerin cream			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84983-026
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	

<b>TOCOPHEROL</b> (UNII: R0ZB2556P8) (TOCOPHEROL - UNII:R0ZB2556P8)	TOCOPHEROL	0.12 mg in 120 mg
<b>ASCORBIC ACID</b> (UNII: PQ6CK8PD0R) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	0.12 mg in 120 mg
<b>PANTHENOL</b> (UNII: VW9CM0067Z) (PANTHENOL - UNII:VW9CM0067Z)	PANTHENOL	0.12 mg in 120 mg
<b>MENAQUINONE 7</b> (UNII: 8427BML8NY) (MENAQUINONE 7 - UNII:8427BML8NY)	MENAQUINONE 7	0.12 mg in 120 mg
<b>MENTHA SUAVEOLENS LEAF</b> (UNII: 1341ZC68MK) (MENTHA SUAVEOLENS LEAF - UNII:1341ZC68MK)	MENTHA SUAVEOLENS LEAF	1.08 mg in 120 mg
<b>ARNICA MONTANA FLOWER</b> (UNII: OZ0E5Y15PZ) (ARNICA MONTANA FLOWER - UNII:OZ0E5Y15PZ)	ARNICA MONTANA FLOWER	0.12 mg in 120 mg
<b>COCONUT OIL</b> (UNII: Q9L0073W7L) (COCONUT OIL - UNII:Q9L0073W7L)	COCONUT OIL	0.12 mg in 120 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>ALLANTOIN</b> (UNII: 344S277G0Z)	0.12 mg in 120 mg
<b>CARBOMER HOMOPOLYMER TYPE A (ALLYL PENTAERYTHRITOL CROSSLINKED)</b> (UNII: F68VH75CJC)	0.24 mg in 120 mg
<b>1,2-HEXANEDIOL</b> (UNII: TR046Y3K1G)	0.6 mg in 120 mg
<b>WATER</b> (UNII: 059QF0KO0R)	101.4 mg in 120 mg
<b>TROLAMINE</b> (UNII: 9O3K93S3TK)	0.24 mg in 120 mg
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	9.6 mg in 120 mg
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	3.6 mg in 120 mg
<b>GLYCERYL MONOSTEARATE</b> (UNII: 230OU9XXE4)	2.4 mg in 120 mg

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84983-026-01	120 mg in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/09/2026	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M016	04/09/2026	

**Labeler** - Shantou South Moon Biotechnology Co., Ltd. (457126192)

## Establishment

Name	Address	ID/FEI	Business Operations
Shantou South Moon Biotechnology Co., Ltd.		457126192	manufacture(84983-026)

