

**SCOT-TUSSIN SENIOR SF DMEXP- dextromethorphan hydrobromide, guaifenesin liquid**  
**SCOT-TUSSIN Pharmacal Co., Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Scot-Tussin Senior SF DMEXP**

**Active Ingredients:**

*(in each 5 mL. tsp. teaspoon)*

Dextromethorphan HBr. USP 15mg

Guaifenesin USP 200mg

**Purpose**

Cough Suppressant

Expectorant

**Uses**

- Temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold. Quiets and calms a dry cough.
- Helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive thereby relieving chest congestion.

**Warnings**

\*Phenylketonurics: Contains Phenylalanine.

**Do not use**

- more than the recommended dosage.
- If drowsiness occurs, do not drive or operate machinery.
- if you are taking a prescription monoamine oxidase inhibitor (MAOI); (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- A chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema.
- Difficulty in urination due to enlargement of the prostate gland.
- A Cough accompanied by excessive phlegm (mucus).
- Glaucoma.
- Thyroid disease.

**Stop use and ask a doctor if**

- Cough lasts more than seven days, returns or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

Do Not Use.

**Keep out of reach of children.**

In case of accidental overdose get medical help or contact a Poison Control Center right away.

**Directions**

- Follow Dosage Chart.
- Do not take more than 4 doses in any 24 hour period.
- This product is not intended for use in children under 12 years of age.

| age                                   | dose   |
|---------------------------------------|--|
| Adults and children over 12 years old | 2 teaspoons<br>(2 tsp. 10 ml)<br>every 6 hours |
| Children Under 12 years old           | DO NOT USE                                     |

**Other Information**

- Each teaspoon contains Phenylalanine 7.5 mg.
- Store at 20° – 25° C (68° – 77° F).
- Do not refrigerate.
- Dosage cup provided.

**Inactive Ingredients**

\*Aspartame (see warning), Benzoic Acid, Citric Acid, Glycerin, Hydroxypropylmethylcellulose, Menthol, Methyl Paraben, Peppermint Stick Flavor, Propyl Paraben, Propylene Glycol, Purified Water.

**PRINCIPAL DISPLAY PANEL**

NDC 0372-0050-04

**SCOT-TUSSIN**

**SENIOR**

**Maximum Strength COUGH SUPPRESSANT & EXPECTORANT**

4 FL OZ (118 ml)

**SUGAR-FREE**

# SCOT-TUSSIN®

**Drug Facts** (Continued)

**Stop use and ask a doctor if**

- cough lasts more than seven days, returns or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding, Do Not Use. Keep out of reach of children.** In case of overdose get medical help or contact a Poison Control Center right away.

**Directions**

- follow Dosage Chart.
- do not take more than 4 doses in any 24-hour period.
- this product is not intended for use in children under 12 years of age.

| age                                   | dose                                     |
|---------------------------------------|--|
| Adults and children over 12 years old | 2 teaspoons (2 tsp, 10 mL) every 6 hours |
| Children under 12 years old           | DO NOT USE                               |

**Other information**

- each teaspoon contains Phenylalanine 7.5mg.
- store at 20°-25° C (68°-77° F).
- do not refrigerate.
- dosage cup provided.

**Inactive ingredients**

aspartame\* (see warning), benzoic acid, citric acid, glycerin, hydroxypropylmethylcellulose, menthol, methyl-paraben, peppermint stick flavor, propyl-paraben, propylene glycol, purified water.

**SUGAR-FREE**

# SCOT-TUSSIN®

**SCOT-TUSSIN QUALITY ASSURANCE**

Decades of experience since 1956 in providing you with quality medicines goes into every product. Scot-Tussin has been an innovator in the formulation of sugar-free and alcohol-free medicines for the diabetic consumer and has been fortunate to have received recommendations from generations of doctors. Scot-Tussin has been and still is committed to the often over-looked special needs patient. Our medicines are safe for the typical consumer as well as those with diabetes, celiac sprue, high blood pressure, and heart disease.

**All Scot-Tussin Brand Medicines are Satisfaction Guaranteed or Your Money Back**

Distributed by:  
Scot-Tussin Pharmacal Co., Inc.  
Cranston, RI 02920 U.S.A.  
Toll Free # 1-800-638-7268 (Scot)  
1-401-942-8555  
© Copyright 1983  
© TM Reg. US Patent Office #657-674

**SUGAR-FREE**

**ADULT** NOC 0372-0050-04

**For Diabetics since 1956**

# SCOT-TUSSIN®

**SENIOR  
MAXIMUM STRENGTH**

**COUGH SUPPRESSANT  
& EXPECTORANT**

DEXTROMETHORPHAN HBr (Cough Suppressant)  
GUAIFENESIN (Expectorant)

**FAST RELIEF OF:  
COUGHING &  
CHEST CONGESTION**

- FIRST SUGAR & ALCOHOL-FREE
- HIGH BLOOD PRESSURE SAFE
- LACTOSE & GLUTEN-FREE
- SODIUM-FREE
- DYE-FREE

LIQUID  
4 FL OZ (118 ml)

**SUGAR-FREE**

# SCOT-TUSSIN®

**IF TAMPER-EVIDENT SEAL AROUND CAP IS BROKEN OR MISSING, DO NOT USE.**

**Drug Facts**

**Active ingredients Purposes**  
(in each 5 mL, tsp, teaspoon)

Dextromethorphan HBr, USP  
15mg.....Cough Suppressant  
Guaifenesin USP 200mg.....Expectorant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold. Quiets and calms a dry cough.
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive thereby relieving chest congestion.

**Warnings**

- \*Phenylketoneurics: Contains Phenylalanine.

**Do not use**

- more than the recommended dosage.
- if drowsiness occurs do not drive or operate machinery.
- if you are taking a prescription monoamine oxidase inhibitor (MAOI); (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema.
- difficulty in urination due to enlargement of the prostate gland.
- a cough accompanied by excessive phlegm (mucus).
- glaucoma.
- thyroid disease.

## SCOT-TUSSIN SENIOR SF DMEXP

dextromethorphan hydrobromide, guaifenesin liquid

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:0372-0050 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength                | Strength          |
|--|----------------------------------|-------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)<br>(DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN<br>HYDROBROMIDE | 15 mg<br>in 5 mL  |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)                           | GUAIFENESIN                      | 200 mg<br>in 5 mL |

### Inactive Ingredients

| Ingredient Name                            | Strength |
|--|----------|
| ASPARTAME (UNII: Z0H242BBR1)               |          |
| BENZOIC ACID (UNII: 8SKN0B0MIM)            |          |
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) |          |
| GLYCERIN (UNII: PDC6A3C0OX)                |          |

|  |  |
|--|--|
| <b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)    |  |
| <b>MENTHOL</b> (UNII: L7T10EP3A)           |  |
| <b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)    |  |
| <b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)    |  |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3) |  |
| <b>WATER</b> (UNII: 059QF0KO0R)            |  |

### Product Characteristics

|                 |                               |                     |  |
|-----------------|-------------------------------|---------------------|--|
| <b>Color</b>    |                               | <b>Score</b>        |  |
| <b>Shape</b>    |                               | <b>Size</b>         |  |
| <b>Flavor</b>   | PEPPERMINT (Peppermint Stick) | <b>Imprint Code</b> |  |
| <b>Contains</b> |                               |                     |  |

### Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:0372-0050-04 | 118 mL in 1 BOTTLE  |                      |                    |

### Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part341                                  | 12/07/1999           |                    |

**Labeler** - SCOT-TUSSIN Pharmacial Co., Inc. (001203918)

Revised: 2/2013

SCOT-TUSSIN Pharmacial Co., Inc.