

MOTION SICKNESS RELIEF- dimenhydrinate tablet
Rite Aid Corporation

Rite Aid 44-198

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- alcohol, sedatives, and tranquilizers may increase drowsiness
- marked drowsiness may occur
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Other information

- **each tablet contains:** calcium 35 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- protect from moisture
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 11822-0198-7

Compare to the active ingredient in
Dramamine® Original Formula*

MOTION SICKNESS

RELIEF

DIMENHYDRINATE 50 mg

ANTIEMETIC**PREVENTS MOTION SICKNESS**

Prevents nausea, vomiting & dizziness

For children & adults

ACTUAL SIZE

36 TABLETS

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

* This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine®
Original Formula. 50844 REV0518G19807

DISTRIBUTED BY: RITE AID,
200 NEWBERRY COMMONS
ETTERS, PA 17319
www.riteaid.com

**SATISFACTION
GUARANTEE**

If you're not satisfied, we'll
happily refund your money.

MOTION SICKNESS RELIEF

DIMENHYDRINATE 50 mg

ANTIEMETIC

NDC 11822-0198-7

Compare to the active ingredient in
Dramamine® Original Formula*

MOTION SICKNESS RELIEF

DIMENHYDRINATE 50 mg

ANTIEMETIC
PREVENTS MOTION SICKNESS

Prevents nausea, vomiting & dizziness
For children & adults



ACTUAL SIZE

36
TABLETS

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

DISTRIBUTED BY: RITE AID,
200 NEWBERRY COMMONS
ETTERS, PA 17319
www.riteaid.com

SATISFACTION GUARANTEE

If you're not satisfied, we'll happily refund your money.

No print/No varnish area
Lot no/Exp date



B-1702-198-07-R3
REV0518619807

*This product is not manufactured or distributed by Mediatech Products Inc., owner of the registered trademark Dramamine®
Original Formula, 50844 REV0518619807

Drug Facts (continued)	Other information ■ each tablet contains: calcium 35 mg ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture
Drug Facts (continued)	Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid
Drug Facts (continued)	Questions or comments? 1-800-426-9391

Drug Facts (continued)	Warnings Do not use for children under 2 years of age unless directed by a doctor. ■ Ask a doctor before use if you have a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. ■ When using this product alcohol, sedatives, and tranquilizers may increase drowsiness ■ marked drowsiness may occur
Drug Facts (continued)	Uses for prevention and treatment of these symptoms associated with motion sickness: ■ nausea ■ vomiting ■ dizziness
Drug Facts (continued)	Active ingredient (in each tablet) Purpose Dimenhydrinate 50 mg..... Antiemetic
Drug Facts (continued)	Directions ■ to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity adults and children 1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor children 6 to under 12 years ½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor children 2 to under 6 years ½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor
Drug Facts (continued)	Keep out of reach of children. In case of overdose, get before use. If pregnant or breast-feeding, ask a health professional for use. ■ use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages

Rite Aid 44-198

MOTION SICKNESS RELIEF

dimenhydrinate tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:11822-0198

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;198
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0198-2	2 in 1 CARTON	12/01/1992	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-0198-7	6 in 1 CARTON	12/01/1992	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	12/01/1992	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11822-0198)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-0198) , pack(11822-0198)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-0198)

Revised: 1/2025

Rite Aid Corporation