

**MOTION SICKNESS RELIEF- dimenhydrinate tablet**  
**Rite Aid Corporation**

-----  
**Rite Aid 44-198**

***Active ingredient (in each tablet)***

Dimenhydrinate 50 mg

***Purpose***

Antiemetic

***Uses***

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

***Warnings***

**Do not use**

for children under 2 years of age unless directed by a doctor.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- alcohol, sedatives, and tranquilizers may increase drowsiness
- marked drowsiness may occur
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

**Other information**

- **each tablet contains:** calcium 35 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- protect from moisture
- see end flap for expiration date and lot number

**Inactive ingredients**

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

**Questions or comments?**

**1-800-426-9391**

**Principal Display Panel**

NDC 11822-0198-7

Compare to the active ingredient in  
**Dramamine® Original Formula\***

**MOTION SICKNESS**

RELIEF

DIMENHYDRINATE 50 mg

**ANTIEMETIC****PREVENTS MOTION SICKNESS**

Prevents nausea, vomiting & dizziness

For children & adults

ACTUAL SIZE

**36** TABLETS

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

\* This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine®  
Original Formula. 50844 REV0518G19807

**DISTRIBUTED BY:** RITE AID,  
200 NEWBERRY COMMONS  
ETTERS, PA 17319  
[www.riteaid.com](http://www.riteaid.com)

**SATISFACTION  
GUARANTEE**

If you're not satisfied, we'll  
happily refund your money.

# MOTION SICKNESS RELIEF

**DIMENHYDRINATE 50 mg**

ANTIEMETIC

NDC 11822-0198-7

Compare to the active ingredient in  
Dramamine® Original Formula\*

# MOTION SICKNESS RELIEF

**DIMENHYDRINATE 50 mg**

ANTIEMETIC  
PREVENTS MOTION SICKNESS

Prevents nausea, vomiting & dizziness  
For children & adults



ACTUAL SIZE

**36**  
TABLETS

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

DISTRIBUTED BY: RITE AID,  
200 NEWBERRY COMMONS  
ETTERS, PA 17319  
www.riteaid.com

**SATISFACTION GUARANTEE**

If you're not satisfied, we'll happily refund your money.

No print/No varnish area  
Lot no/Exp date



B-1702-198-07-R3  
REV0518619807

\*This product is not manufactured or distributed by Mediatech Products Inc., owner of the registered trademark Dramamine®  
Original Formula, 50844 REV0518619807

<p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>each tablet contains: calcium 35 mg</li> <li>TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN</li> <li>store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p>■ see end flap for expiration date and lot number</p> <p><b>Inactive ingredients</b> croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid</p> <p><b>Questions or comments?</b> 1-800-426-9391</p>
--	--

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Purpose Dimenhydrinate 50 mg, ..... Antiemetic</p> <p><b>Uses</b> for prevention and treatment of these symptoms associated with motion sickness: nausea ■ vomiting ■ dizziness</p> <p><b>Warnings</b> Do not use for children under 2 years of age unless directed by a doctor. Ask a doctor before use if you have a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. When using this product alcohol, sedatives, and tranquilizers may increase drowsiness ■ marked drowsiness may occur</p>	<p><b>Directions</b> ■ to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity adults and children 1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor children 6 to under 12 years ½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor children 2 to under 6 years ½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor</p> <p><b>Drug Facts (continued)</b> ■ use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>
--	--

Rite Aid 44-198

## MOTION SICKNESS RELIEF

dimenhydrinate tablet

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:11822-0198

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIMENHYDRINATE</b> (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	44;198
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0198-2	2 in 1 CARTON	12/01/1992	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-0198-7	6 in 1 CARTON	12/01/1992	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	12/01/1992	

**Labeler** - Rite Aid Corporation (014578892)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11822-0198)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-0198) , pack(11822-0198)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-0198)

Revised: 1/2026

Rite Aid Corporation