ZATEAN-PN PLUS- levomefolate calcium, folic acid, ascorbic acid, cholecalciferol, .alpha.tocopherol, pyridoxine, cyanocobalamin, biotin, calcium carbonate, ferrous fumarate, potassium iodide, magnesium oxide, doconexent, and icosapent capsule Trigen Laboratories, LLC

-----

Zatean<sup>™</sup>-Pn Plus Softgels

SUPPLEMENT FACTS

# Supplement Facts

Serving Size: 1 Softgel Capsule Servings Per Container: 30

10 0.5 g 85 mg 5 mcg (200 IU) 7 mg* (10 IU) 25 mg 1667 mcg DFE	<1%** 71% 33% 37% 1250% 278%
85 mg 5 mcg (200 IU) 7 mg* (10 IU) 25 mg 1667 mcg	71% 33% 37% 1250%
5 mcg (200 IU) 7 mg* (10 IU) 25 mg 1667 mcg	33% 37% 1250%
(200 IŬ) 7 mg* (10 IU) 25 mg 1667 mcg	37% 1250%
(10 IU) 25 mg 1667 mcg	1250%
1667 mcg	
	278%
12 mcg	429%
250 mcg	714%
140 mg	11%
28 mg	104%
150 mcg	52%
45 mg	11%
300 mg	†
40 mg	†
	250 mcg 140 mg 28 mg 150 mcg 45 mg 300 mg

**Other Ingredients:** Orange Cream Flavor O.S., Citric Acid, Biomin Potassium Citrate, Soy Lecithin, Yellow Beeswax

**Shell Ingredients**: Bovine Gelatin, Glycerin, Purified Water, Sorbitol, Titanium Dioxide, Ethyl Vanillin, FD&C Yellow #5, FD&C Blue #1, and Caramel.

# THIS PRODUCT CONTAINS SOY AND FISH OIL (ANCHOVY).

Zatean<sup>™</sup>-Pn Plus is a prescription prenatal/postnatal multivitamin/ mineral softgel with an essential fatty acid and is indicated for the supplemental requirements of patients with nutritional deficiencies or are in

need of nutritional supplementation.

# CONTRAINDICATIONS

This product is contraindicated in patients with a known hypersensitivity to any of the ingredients including soy or fish oil.

# WARNING

Administration of omega-3 fatty acids – including DHA, should be avoided in patients with inherited or acquired bleeding diatheses, including those taking anticoagulants.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. KEEP THIS PRODUCT OUT OF REACH OF CHILDREN. In case of accidental overdose, call a doctor or poison control center immediately.

# PRECAUTIONS

**General:** Folic acid alone is an improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where vitamin B12 is deficient. Folic acid in doses above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurologic manifestations remain progressive. Reduced folates may be less likely than folic acid to mask vitamin B12 deficiency. Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

The patient's medical conditions and consumption of other drugs, herbs, and/or supplements should be considered.

Biotin levels higher than the recommended daily allowance may cause interference with some laboratory tests, including cardiovascular diagnostic tests (e.g. troponin) and hormone tests, and may lead to incorrect test results. Tell your healthcare provider about all prescription and over-the-counter medicines, vitamins, and dietary supplements that you take, including biotin.

This product contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible patients. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

## **DRUG INTERACTIONS**

Pyridoxine hydrochloride should not be given to patients receiving the drug levodopa, because the action of levodopa is antagonized by pyridoxine hydrochloride. However, pyridoxine hydrochloride may be used concurrently in patients receiving a preparation containing both carbidopa and levodopa.

There is a possibility of increased bleeding due to pyridoxine interaction with anticoagulants (e.g., Aspirin, Heparin, Clopidogrel).

# **ADVERSE REACTIONS**

Allergic sensitization has been reported following both oral and parenteral administration of folic acid, as well as possibly the use of other forms of folates, including reduced folates. Paresthesia, somnolence, nausea and headaches have been reported with pyridoxine hydrochloride. Mild transient diarrhea, polycythemia vera, itching, transitory exanthema and the feeling of swelling of the entire body have been associated with cyanocobalamin.

## DESCRIPTION

Zatean<sup>TM</sup> - Pn Plus is an oblong shaped, dark green soft gelatin capsule containing brown paste with white "T582" laser etched.

## **DIRECTIONS FOR USE**

Take one softgel capsule by mouth daily or as prescribed by a physician.

## HOW SUPPLIED

Zatean<sup>™</sup> - Pn Plus is available in bottles of 30 softgel capsules.

## **PRODUCT CODE 13811-582-30**

#### STORAGE

Store at 20°-25°C (68°-77°F), excursions permitted to 15°-30°C (59°-86°F) [See USP Controlled Room Temperature].

## KEEP OUT OF REACH OF CHILDREN.

For use on the order of a healthcare practitioner.

Call your doctor about side effects. To report side effects, call Trigen Laboratories, LLC at 1-877-482-3788 or FDA at 1-800-FDA-1088.

Rev. 09/2019

Customer Service: 1-877-482-3788

Manufactured for: Trigen Laboratories, LLC Bridgewater, NJ 08807 www.trigenlab.com



PRINCIPAL DISPLAY PANEL - 30 Softgel Bottle Label

13811-582-30

Zatean<sup>™</sup>-Pn Plus Softgels

**30 SOFTGELS** 

DIRECTIONS FOR USE Supplement Facts Take one softgel capsule by mouth daily or as presaribed by a 13811-582-30 Sewing Stee: 1 Softgel Capsule Sewings Ver Container: 30 physician. STORAGE Store at 20°-25°C (68°-77°F), excursions permitted to 15°-30°C Amount Ner %EV for Pregnant Sewing and Lactating Women (59° 86°F) [See USP Controlled Room Temperature]. Zatean<sup>™</sup>-Pn Calories Total Fat 10 0.5 g WA RNING: Accidental overdose of iron-containing <1%\*\* products is a leading cause of fatal poisoning in children under 6. KEEP THIS PRODUCT OUT OF Peel label for prescribing and product information. Vitamin C (Ascorbic Acid) Vitamin D (as Cholecalaferol) 85 mg 5 mg (200 IJ) 71% Plus 33% **REACH OF CHILDREN. In case of accidental overd** Vitamin E (as dAlpha Toopheryl Acetate) 7 mg\* (10 IU) 37% Vitamin B<sub>6</sub> (as Pyridaxine HCl) Folate (as L-Wethylfolate (6(S)-5-MTHF) 600 mcg and Folic Add 400 mcg) 25 mg 1667 mg DFE call a doctor or paison control center immediately. 1250% Softgels 278% Call your doctor about side effects. To report side effects, call Trigen Laboratories, LLC at 429% Vitamin B<sub>12</sub> (as Granocobalamin) 12 m cg 1-877-482-3788 or FDA at 1-800-FDA -1088. Rintin 250 mg 7145 PRENATAL DIETARY KEEP OUT OF THE REACH OF CHILDREN. Calcium (as Calcium Carbonate) 140 mg 11% SUPPLEMENT Manufactured for: Trigen Laboratories, LLC, Bridgewater, NJ 08807 Iron (as Ferrons Furnarate) Iodine (as Patassium Iodide) 28 mg 104% 52% 150 mg www.trigenlab.com Magnesium (as Magnesium Oxide 11% 45 mg **30 SOFTGELS** Rev. 10/2019 DHA (Domynhermenoic Ario 300 mg EPA (Ecceapentaenoic Acid) 40 mg † Daily Value (DV) not established "Equal to mg Alpha - Tocopherol \*"Percent Daily Values are based on a 2,000 calorie diet Exp: GE 13811 58230 III 7 3

## ZATEAN-PN PLUS

levomefolate calcium, folic acid, ascorbic acid, cholecalciferol, .alpha.-tocopherol, pyridoxine, cyanocobalamin, biotin, calcium carbonate, ferrous fumarate, potassium iodide, magnesium oxide, doconexent, and icosapent capsule

Product Information				
Product Type	DIETARY SUPPLEMENT	Item Code (Source)	NHRIC:13811-582	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
LEVOMEFOLIC ACID (UNII: 8S95DH25XC) (LEVOMEFOLIC ACID - UNII:8S95DH25XC)	LEVOMEFOLIC ACID	600 ug	
folic acid (UNII: 935E97BOY8) (FOLIC ACID - UNII:935E97BOY8)	folic acid	400 ug	
ASCORBIC ACID (UNII: PQ6CK8PD0R) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	85 mg	
CHOLECALCIFEROL (UNII: 1C6V77QF41) (CHOLECALCIFEROL - UNII:1C6V77QF41)	CHOLECALCIFEROL	200 [iU]	
.ALPHATOCOPHEROL (UNII: H4N855PNZ1) (.ALPHATOCOPHEROL - UNII:H4N855PNZ1)	.ALPHATOCOPHEROL	10 [iU]	
PYRIDO XINE (UNII: KV2JZ1BI6Z) (PYRIDO XINE - UNII:KV2JZ1BI6Z)	PYRIDO XINE	25 mg	
CYANOCOBALAMIN (UNII: P6 YC3EG204) (CYANOCOBALAMIN - UNII:P6 YC3EG204)	CYANOCOBALAMIN	12 ug	
Biotin (UNII: 6SO6U10H04) (Biotin - UNII:6SO6U10H04)	Biotin	250 ug	
calcium carbonate (UNII: H0G9379FGK) (CALCIUM CATION - UNII:2M83C4R6ZB)	calcium carbonate	140 mg	
ferrous fumarate (UNII: R5L488RY0Q) (FERROUS CATION - UNII:GW895810WR)	FERROUS CATION	28 mg	
potassium iodide (UNII: 1C4QK22F9J) (IODIDE ION - UNII:09G4I6V86Q)	potassium iodide	150 ug	
magnesium oxide (UNII: 3A3U0GI71G) (MAGNESIUM CATION - UNII:T6V3LHY838)	magnesium oxide	45 mg	
Doconexent (UNII: ZAD9OKH9JC) (DOCONEXENT - UNII:ZAD9OKH9JC)	Doconexent	300 mg	
Icosapent (UNII: AAN7QOV9EA) (ICOSAPENT - UNII:AAN7QOV9EA)	Icosapent	40 mg	

**Inactive Ingredients** 

**Ingredient Name** 

Strength

caramel (UNII: T9D99G2B	1R)				
ethyl vanillin (UNII: YC9ST449YJ)					
FD&C Blue no. 1 (UNII: H3	R47K3TBD)				
FD&C Yellow no. 5 (UNII:	I753WB2F1M)				
gelatin (UNII: 2G86QN327	L)				
glycerin (UNII: PDC6A3C0OX)					
lecithin, soybean (UNII: 1DI56QDM62)					
water (UNII: 059QF0KO0R)					
sorbitol (UNII: 506T60A25R)					
titanium dioxide (UNII: 15FIX9V2JP)					
potassium citrate anhydrous (UNII: 86R1NVR0HW)					
yellow wax (UNII: 2ZA36H	yellow wax (UNII: 2ZA36H0S2V)				
ANHYDROUS CITRIC AC	ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)				
<b>ORANGE OIL</b> (UNII: AKN3	3KSD11B)				
Packaging					
# Item Code	Package Description	Marketin	ig Start Date	Ma	rketing End Date
1 NHRIC:13811-582-30	30 in 1 BOTTLE				
Maultating Inform					
Marketing Inform					
Marketing Category	Application Number or Monogra	ph Citation	Marketing Start D	Date	Marketing End Date
DIETARY SUPPLEMENT			0 5/0 1/20 10		

Supplement Facts				
Serving Size :		Serving per Container :		
	Amount Per Serving	% Daily Value		
color				
imprint				
scoring	1			
shape				
size (solid drugs)	26 mm			

Labeler - Trigen Laboratories, LLC (830479668)

Revised: 9/2019

Trigen Laboratories, LLC