

**ROMPE PECHO MAX- acetaminophen, dextromethorphan hbr,
guaifenesin liquid
Efficient Laboratories Inc**

Rompe Pecho Max (PE Free)

Active Ingredients:(in each 30ml doses cup) **Purpose**

Acetaminophen 650mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough Suppressant
Guaifenesin 400 mg	Expectorant

Purpose

Pain reliever/fever reducer
Cough Suppressant
Expectorant

Uses

- temporarily relieves these common cold/flu symptoms:
 - cough due to minor throat and bronchial irritation
- minor aches and pains • headache • sore throat
- temporarily reduces fever.
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks daily while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease • liver disease • high blood pressure • thyroid disease • diabetes • difficulty in urination due to an enlarged prostate gland • a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema • a cough that is followed by excessive phlegm (mucus).

Ask a doctor or pharmacist before use if you are:

- taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if:

- symptoms do not improve within 7 days, tend to recur, or are accompanied by a fever, rash, or persistent headache
- Nervousness, dizziness or sleeplessness occurs • fever gets worse or last more than 3 days • redness or swelling is present or new symptoms occur. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions do not take more than directed (see overdose warning)

- Do not exceed more than 6 doses in any 24-hour period
- Shake well before use
- measure only with dosing cup provided

- keep dosing cup with product
- mL=miliiter TSP=teaspoonful
- this adult product is not intended for use in children under 12 years of age

Age	• Dose
• adults and children 12 years of age and over	• 20 mL every 4 hours
• children under 12 years of age	• do not use

Inactive ingredients: blue cohosh root extract, citric acid, echinacea root extract, eucalyptus oil, ginkgo biloba leaf extract, glycerin, goldenseal root extract, honey, horehound (flower, leaf, stem) extract, licorice root extract, menthol, methylparaben, mullein leaf extract, myrrh gum, potassium citrate, potassium sorbate, propylene glycol, Propylparaben, slippery elm bark extract, sodium chloride, sucralose, water, wild cherry bark extract and zinc sulfate.

Questions or Comments?

305-805-3456 Monday - Friday 9AM to 5PM EST

www.efficientlabs.com

Distributed by

Efficient Laboratories, Inc., Miami, Fl 33166

Made in U.S.A.

7 Inch.

<p>Active Ingredients (in each 20 mL)</p> <p>Acetaminophen, 325 mg.....Pain reliever/fever reducer</p> <p>Dextromethorphan, 10 mg.....Cough suppressant</p> <p>Guafenesin, 400 mg.....Expectorant</p> <p>Uses Temporarily relieves these common cold flu symptoms:</p> <ul style="list-style-type: none"> cough due to throat and bronchial irritation minor aches and pains - sore throat - headache temporarily reduces hoarseness, nasal congestion, nasal and the bronchial secretions to rid the bronchial passages of bothersome mucus and make coughing more productive. <p>Warnings</p> <p>Read the entire product information carefully.</p> <p>Do not take more than 8 doses in any 24 hours period - unless you have been told to do so by your doctor. Do not take more than 2 doses every 4 hours. Do not take more than 1 dose every 4 hours. Do not take more than 1 dose every 4 hours. Do not take more than 1 dose every 4 hours.</p> <p>Directions</p> <p>Adults and Children 12 years & over: 20 mL every 4 hours.</p> <p>Children under 12 years of age: 10 mL every 4 hours.</p> <p>Do not take more than 8 doses in any 24 hours period - unless you have been told to do so by your doctor. Do not take more than 2 doses every 4 hours. Do not take more than 1 dose every 4 hours. Do not take more than 1 dose every 4 hours.</p> <p>REV 10/2022 Rev. 1 Rompe Pecho for complete instructions in Spanish.</p>	 <p>Rompe Pecho.</p> <p>MAX Multi-Symptom relief</p> <p>PAIN RELIEVER</p> <p>FEVER REDUCER</p> <p>COUGH SUPPRESSANT</p> <p>EXPECTORANT</p> <p>ALONG WITH</p> <p>Menthol</p> <p>Adults & Children 12 years & over</p> <p>NET WEIGHT: 6 FL OZ (182 mL)</p>	<p>Warnings</p> <p>Do not take more than 8 doses in any 24 hours period - unless you have been told to do so by your doctor. Do not take more than 2 doses every 4 hours. Do not take more than 1 dose every 4 hours. Do not take more than 1 dose every 4 hours.</p> <p>Directions</p> <p>Adults and Children 12 years & over: 20 mL every 4 hours.</p> <p>Children under 12 years of age: 10 mL every 4 hours.</p> <p>Do not take more than 8 doses in any 24 hours period - unless you have been told to do so by your doctor. Do not take more than 2 doses every 4 hours. Do not take more than 1 dose every 4 hours. Do not take more than 1 dose every 4 hours.</p> <p>Uses</p> <p>Temporarily relieves these common cold flu symptoms:</p> <ul style="list-style-type: none"> cough due to throat and bronchial irritation minor aches and pains - sore throat - headache temporarily reduces hoarseness, nasal congestion, nasal and the bronchial secretions to rid the bronchial passages of bothersome mucus and make coughing more productive.
--	--	---

3 Inch.



Rompe Pecho.

MAX Multi-Symptom relief

EFFICIENT

NEW PROSOLVED

Drug Facts (continued)

Directions

Do not take more than 8 doses in any 24 hours - unless you have been told to do so by your doctor. Do not take more than 2 doses every 4 hours. Do not take more than 1 dose every 4 hours. Do not take more than 1 dose every 4 hours.

Warnings

Do not take more than 8 doses in any 24 hours period - unless you have been told to do so by your doctor. Do not take more than 2 doses every 4 hours. Do not take more than 1 dose every 4 hours. Do not take more than 1 dose every 4 hours.

Uses

Temporarily relieves these common cold flu symptoms:

- cough due to throat and bronchial irritation
- minor aches and pains - sore throat - headache
- temporarily reduces hoarseness, nasal congestion, nasal and the bronchial secretions to rid the bronchial passages of bothersome mucus and make coughing more productive.

Drug Facts

Active Ingredients (in each 20 mL)	Purpose
Acetaminophen, 325 mg	Pain reliever/fever reducer
Dextromethorphan, 10 mg	Cough suppressant
Guafenesin, 400 mg	Expectorant

Uses

Temporarily relieves these common cold flu symptoms:

- cough due to throat and bronchial irritation
- minor aches and pains - sore throat - headache
- temporarily reduces hoarseness, nasal congestion, nasal and the bronchial secretions to rid the bronchial passages of bothersome mucus and make coughing more productive.

Warnings

Do not take more than 8 doses in any 24 hours period - unless you have been told to do so by your doctor. Do not take more than 2 doses every 4 hours. Do not take more than 1 dose every 4 hours. Do not take more than 1 dose every 4 hours.

Directions

Adults and Children 12 years & over: 20 mL every 4 hours.

Children under 12 years of age: 10 mL every 4 hours.

Do not take more than 8 doses in any 24 hours period - unless you have been told to do so by your doctor. Do not take more than 2 doses every 4 hours. Do not take more than 1 dose every 4 hours. Do not take more than 1 dose every 4 hours.

Uses

Temporarily relieves these common cold flu symptoms:

- cough due to throat and bronchial irritation
- minor aches and pains - sore throat - headache
- temporarily reduces hoarseness, nasal congestion, nasal and the bronchial secretions to rid the bronchial passages of bothersome mucus and make coughing more productive.

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
POTASSIUM CITRATE ANHYDROUS (UNII: 86R1NVR0HW)	
CAULOPHYLLUM THALICTROIDES ROOT (UNII: JTJ6HH6YEH)	
GLYCERIN (UNII: PDC6A3C0OX)	
ECHINACEA ANGUSTIFOLIA ROOT (UNII: D982V7VT3P)	
EUCALYPTUS OIL (UNII: 2R04ONI662)	
GINKGO BILOBA LEAF OIL (UNII: Y5967KO1JH)	
GOLDENSEAL (UNII: ZW3Z11D0JV)	
HONEY (UNII: Y9H1V576FH)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
HOREHOUND (UNII: K08036XEJV)	
MENTHOL (UNII: L7T10EIP3A)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
LICORICE (UNII: 61ZBX54883)	
MULLEIN LEAF (UNII: 9936O846LI)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
ULMUS RUBRA BARK (UNII: 91QY4PXU8Q)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
PRUNUS SEROTINA BARK (UNII: 5D48E975HA)	
ZINC SULFATE (UNII: 89DS0H96TB)	
MYRRH (UNII: JC71GJ1F3L)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58593-005-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/01/2026	

Labeler - Efficient Laboratories Inc (969044932)

Registrant - Efficient Laboratories Inc (969044932)

Revised: 4/2026

Efficient Laboratories Inc