

**COLEMAN BITE AND STING RELIEF- external anesthetic and first aid
antiseptic liquid
Wisconsin Pharmacal Company**

Warnings For external use only. **When using this product** ■ Do not get in eyes ■ Do not apply over large areas of the body ■ In case of deep or puncture wounds, animal bites or serious burns consult a doctor. ■ Do not bandage. **Stop use and ask doctor if** ■ condition worsens, or symptoms persist for more than 7 days or clear up and occur again within a few days. Do not use longer than one week unless directed by doctor. **Keep out of reach of children.** If swallowed, get medical help or contact a poison control center right away. If in eyes, flush with water for 15 minutes and call a physician. **Drug Facts Continued** ►

Warnings For external use only. **When using this product** ■ Do not get in eyes ■ Do not apply over large areas of the body ■ In case of deep or puncture wounds, animal bites or serious burns consult a doctor. ■ Do not bandage. **Stop use and ask doctor if** ■ condition worsens, or symptoms persist for more than 7 days or clear up and occur again within a few days. Do not use longer than one week unless directed by doctor. **Keep out of reach of children.** If swallowed, get medical help or contact a poison control center right away. If in eyes, flush with water for 15 minutes and call a physician. **Drug Facts Continued** ►

Warnings For external use only. **When using this product** ■ Do not get in eyes ■ Do not apply over large areas of the body ■ In case of deep or puncture wounds, animal bites or serious burns consult a doctor. ■ Do not bandage. **Stop use and ask doctor if** ■ condition worsens, or symptoms persist for more than 7 days or clear up and occur again within a few days. Do not use longer than one week unless directed by doctor. **Keep out of reach of children.** If swallowed, get medical help or contact a poison control center right away. If in eyes, flush with water for 15 minutes and call a physician. **Drug Facts Continued** ►

Warnings For external use only. **When using this product** ■ Do not get in eyes ■ Do not apply over large areas of the body ■ In case of deep or puncture wounds, animal bites or serious burns consult a doctor. ■ Do not bandage. **Stop use and ask doctor if** ■ condition worsens, or symptoms persist for more than 7 days or clear up and occur again within a few days. Do not use longer than one week unless directed by doctor. **Keep out of reach of children.** If swallowed, get medical help or contact a poison control center right away. If in eyes, flush with water for 15 minutes and call a physician. **Drug Facts Continued** ►

Directions Apply soon after an insect bite, to counteract the effects of bites and stings from bees, wasps, mosquitoes, ticks, hornets, ants, fire ants, deerflies, sand fleas, horseflies and chiggers ■ Clean area ■ Apply a small amount to the bite area while massaging with sponge tip applicator. If itching persists, apply again in 10-15 minutes ■ For ticks and bees, remove tick or stinger before treatment ■ Adults and children 2 years and over: Apply to affected area 1 to 3 times daily ■ Children under 2 years: Consult a doctor

Directions Apply soon after an insect bite, to counteract the effects of bites and stings from bees, wasps, mosquitoes, ticks, hornets, ants, fire ants, deerflies, sand fleas, horseflies and chiggers ■ Clean area ■ Apply a small amount to the bite area while massaging with sponge tip applicator. If itching persists, apply again in 10-15 minutes ■ For ticks and bees, remove tick or stinger before treatment ■ Adults and children 2 years and over: Apply to affected area 1 to 3 times daily ■ Children under 2 years: Consult a doctor

Inactive ingredients Propylene Glycol

Questions or comments? 1-800-558-6614

Uses First aid to help prevent infection in minor cuts, scrapes, and burns.

Drug Facts

Active ingredients

Benzocaine 10%	Analgesic, Anesthetic, Antipruritic	Purpose
Phenol 1.35%	First Aid Antiseptic	
Camphor 3%	External Analgesic	

Drug Facts

Active ingredients

Benzocaine 10%	Analgesic, Anesthetic, Antipruritic	Purpose
Phenol 1.35%	First Aid Antiseptic	
Camphor 3%	External Analgesic	

Directions Apply soon after an insect bite, to counteract the effects of bites and stings from bees, wasps, mosquitoes, ticks, hornets, ants, fire ants, deerflies, sand fleas, horseflies and chiggers ■ Clean area ■ Apply a small amount to the bite area while massaging with sponge tip applicator. If itching persists, apply again in 10-15 minutes ■ For ticks and bees, remove tick or stinger before treatment ■ Adults and children 2 years and over: Apply to affected area 1 to 3 times daily ■ Children under 2 years: Consult a doctor

Inactive ingredients Propylene Glycol

Questions or comments? 1-800-558-6614

Drug Facts

Active ingredients

Benzocaine 10%	Analgesic, Anesthetic, Antipruritic	Purpose
Phenol 1.35%	First Aid Antiseptic	
Camphor 3%	External Analgesic	

Uses First aid to help prevent infection in minor cuts, scrapes, and burns.

Warnings For external use only. **When using this product** ■ Do not get in eyes ■ Do not apply over large areas of the body ■ In case of deep or puncture wounds, animal bites or serious burns consult a doctor. ■ Do not bandage. **Stop use and ask doctor if** ■ condition worsens, or symptoms persist for more than 7 days or clear up and occur again within a few days. Do not use longer than one week unless directed by doctor. **Keep out of reach of children. If swallowed, get medical help or contact a poison control center right away. If in eyes, flush with water for 15 minutes and call a physician.** **Drug Facts Continued** ▶

COLEMAN BITE AND STING RELIEF

external anesthetic and first aid antiseptic liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68093-7428
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CAMPHOR (SYNTHETIC) (UNII: 5TJD82A1ET) (CAMPHOR (SYNTHETIC) -	CAMPHOR	0.44 g

UNII:5TJD82A1ET)	(SYNTHETIC)	in 15 mL
PHENOL (UNII: 339NCG44TV) (PHENOL - UNII:339NCG44TV)	PHENOL	0.2 g in 15 mL
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	1.48 g in 15 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white (clear, colorless to pale amber)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68093-7428-1	15 mL in 1 VIAL, PLASTIC; Type 0: Not a Combination Product	08/29/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	08/29/2023	

Labeler - Wisconsin Pharmacal Company (800873986)

Registrant - Wisconsin Pharmacal Company (800873986)

Establishment

Name	Address	ID/FEI	Business Operations
Wisconsin Pharmacal Company		800873986	manufacture(68093-7428) , pack(68093-7428) , label(68093-7428)

Revised: 4/2026

Wisconsin Pharmacal Company