# SINUS PRESSURE AND PAIN MAXIMUM STRENGTH- acetaminophen, phenylephrine hcl tablet DOLGENCORP, LLC

-----

**Dollar General 44-502 RR** 

### Active ingredients (in each caplet)

Acetaminophen 325 mg Phenylephrine HCl 5 mg

#### **Purpose**

Pain reliever/fever reducer Nasal decongestant

#### Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
  - minor aches and pains
  - nasal congestion
  - headache
  - sinus congestion and pressure
- promotes sinus drainage
- temporarily reduces fever

# Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

## Ask a doctor before use if you have

- diabetes
- liver disease
- heart disease
- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland

### Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

### When using this product

do not exceed recommended dosage.

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- do not use more than directed
- adults and children 12 years and over
  - take 2 caplets every 4 hours
  - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

#### Other information

# ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

#### Inactive ingredients

corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, povidone, silicon dioxide, stearic acid

Questions or comments?

1-888-309-9030

Principal display panel

DG™ | health

Compare to active ingredients of Sudafed PE® Sinus Pressure + Pain\*

**Maximum Strength** 

Sinus Pressure & Pain

**Acetaminophen** • Pain Reliever/Fever Reducer Phenylephrine HCl • Nasal Decongestant

**Relieves:** • Sinus pressure & congestion

• Sinus headache

Pseudoephedrine Free

Non Drowsy

24 Caplets

Actual Caplet Size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark SUDAFED PE® SINUS PRESSURE + PAIN.

50844 REV0820L50208

DISTRIBUTED BY OLD EAST MAIN CO. 100 MISSION RIDGE GOODLETTSVILLE, TN 37072

# **Maximum Strength**

# Sinus Pressure & Pain

**DG**™ health

Compare to active ingredients of SUDAFED PE<sup>®</sup> SINUS PRESSURE + PAIN

S 80

ZM

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

PRINT/NO VARNISH Lot & exp date

**Maximum Strength** 

# Sinus Pressure & Pain

Acetaminophen • Pain Reliever/Fever Reducer Phenylephrine HCl • Nasal Decongestant

Relieves: • Sinus pressure & congestion

Sinus headache

Pseudoephedrine Free

24 Caplets

Questions or comments? 1-888-309-9030

microcrystalline cellulose, povidone, silicon dioxide, stearic acid crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, Inactive ingredients corn starch, croscarmellose sodium,

nund racts (continued)

■ see end flap for expiration date and lot number ■ store at Z5 U(1/1); excursions permitted between 15-30 U(20-86) BLISTER IS TORN OR BROKEN

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR Other information

nund Lacts (continued)

■ children under 12 years: aska doctor

■ do not take more than 10 caplets in 24 hours ■ adults and children 12 years and over ■ take 2 caplets every 4 hours

Directions a do not use more than directed

help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any Keep out of reach of children. In case of accidental overdose, get medical If pregnant or breast-feeding, ask a health professional before use.

These could be signs of a serious condition. ■ redness or swelling is present ■ new symptoms occur

■ Tever gets worse or lasts more than 3 days ■ pain or nasal congestion gets worse or lasts more than 7 days

■ nervousness, dizziness, or sleeplessness occur 2 top use and ask a doctor if

When using this product do not exceed recommended dosage.

Ask a doctor or pharmacist before use if you are taking the blood thinning

 difficulty in urination due to enlargement of the prostate gland ■ heart disease ■ thyroid disease ■ high blood pressure aceasib tavil ■ safadsib ■ Ask a doctor before use if you have

■ it you have ever had an allergic reaction to this product or any of its or pharmacist before taking this product.

Urug Facts (continued)

do not know if your prescription drug contains an MAOI, ask a doctor Parkinson's disease), or for 2 weeks after stopping the MAO I drug. If yo (certain drugs for depression, psychiatric or emotional conditions, or ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)

acetaminophen, ask a doctor or pharmacist. nonprescription). If you are not sure whether a drug contains

■ with any other drug containing acetaminophen (prescription or

If a skin reaction occurs, stop use and seek medical help right away. may include: mskin reddening mblisters mrash Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms ■3 or more alcoholic drinks every day while using this product

may occur if you take

more than 4,000 mg of acetaminophen in 24 hours

Liver warning: This product contains acetaminophen. Severe liver damage Marnings

■ promotes sinus drainage ■ temporarily reduces fever

 sinus congestion and pressure ■ minor aches and pains ■ nasal congestion ■ headache

or other upper respiratory allergies, and the common cold: USES I temporanly relieves these symptoms associated with hay tever

Nasal decongestant Phenylephrine HCI 5 mg. Active ingredients (in each caplet) Purpose

COMPLETE PRODUCT INFORMATION KEEP OUTER PACKAGE FOR

Drug Facts

B-0315-502-08RR REV0820L50208

\*This product is not manufactured or Corporation, owner of the registered or SINUS PRESSURE + PAIN. 50844 REV0820L50208

or distributed by Johnson & Johnson of trademark SUDAFED PE®

DISTRIBUTED BY OLD EAST MAIN CO. 100 MISSION RIDGE GOODLETTSVILLE, TN 37072

100%
Satisfaction
Guaranteed!
(888) 309-9030



# SINUS PRESSURE AND PAIN MAXIMUM STRENGTH

acetaminophen, phenylephrine hcl tablet

**Route of Administration** ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
1 11=111==1 1111111= 111=1111== (0:1111 0 1), (1:11=111==1:11111=	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3SJ)			
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STEARIC ACID (UNII: 4ELV7Z65AP)			

Product Characteristics				
Color	orange	Score	no score	
Shape	OVAL	Size	18mm	
Flavor		Imprint Code	44;502	
Contains				

l	Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1	NDC:55910- 925-08	2 in 1 CARTON	06/23/2005		
	1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing	Application Number or Monograph	Marketing Start	Marketing End	
Category	Citation	Date	Date	

OTC Monograph Drug	M012	06/23/2005	

# Labeler - DOLGENCORP, LLC (068331990)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-925) , pack(55910-925)

Establishment					
Name	Address	ID/FEI	<b>Business Operations</b>		
LNK International, Inc.		117025878	manufacture(55910-925)		

Revised: 6/2024 DOLGENCORP, LLC