

**BISACODYL- bisacodyl suppository**  
**AARNA USA INC.**

-----  
**BISACODYL 10 mg Suppositories**  
**Fast Acting Stimulant Laxative**

**Active Ingredients (per suppository)**

Bisacodyl USP, 10 mg

**Purpose**

Laxative

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Uses**

- for relief of occasional constipation
- this product usually produces bowel movement in 1/4 to 1 hour

**Warnings**

For rectal use only.

**Do not use laxative products**

- when abdominal pain, nausea or vomiting are present
- for a period longer than one week

**Ask a doctor before use if you have**

noticed a sudden change in bowel habits that persist over a period of 2 weeks

**When using this product**

it may cause abdominal discomfort, faintness, rectal burning and mild cramps

**Stop use and ask a doctor if**

rectal bleeding occurs, or you fail to have a bowel movement after using a laxative. This may indicate a serious condition.

**If pregnant or breast-feeding**

ask a health professional before use.

**Directions**

- adults and children 12 years of age and older
- detach one suppository from the strip
- remove from wrapper before inserting into the rectum
- the rectal suppository dose is one suppository per day or as directed by a doctor
- children under 12 years of age consult a doctor

**Other information**

store below 30° C (86° F)

**Inactive ingredient**

hydrogenated vegetable oil

**Manufactured for:**

Aarna USA Inc.

Leland, NC 28451

[www.aarnaus.com](http://www.aarnaus.com)

**Bisacodyl 10mg  
Suppositories**

**Drug Facts (continued)**

**Warnings** For rectal use only  
Do not use when abdominal pain, nausea, or vomiting are present.

Ask a doctor before use if you have a sudden change in bowel habits that lasts over a period of 2 weeks.

When using this product you may experience rectal burning, rectal cramps, mild cramps, abdominal discomfort, faintness, Stop use and ask a doctor if you have rectal bleeding or you feel to have a bowel movement after using a laxative. These may be signs of a serious condition. You need to use a laxative for more than one week.


If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
Tear off one suppository from the strip.  
Tear and peel the wrapper in a downward motion.  
Continue tearing downward to almost the full length of the suppository (see illustration at the right).  
Gently remove the suppository from the wrapper.  
Lie on your side and push suppository, with pointed end first, high into the rectum so it will not slip out.  
Remain for 15 to 20 minutes.  
Do not exceed recommended dose. If suppository seems soft, place in refrigerator for a short time before use.  
Adults and children 12 years and over: 1 suppository once daily.  
Ask a doctor.  
Children under 12 years: Ask a doctor.

**Other information** ■ store at temperatures below 30°C (86°F)

**Inactive ingredient** hydrogenated vegetable oil



**Drug Facts**

**Active ingredient (in each suppository)** Bisacodyl USP 10 mg.....Stimulant laxative

**Purpose** Stimulant laxative

**Uses** ■ relieves occasional constipation ■ this product generally produces a bowel movement in 15 minutes to 1 hour

**Bisacodyl 10mg  
Suppositories**

NDC 82568-0028-2

**Bisacodyl 10mg  
Suppositories**

FAST ACTING STIMULANT LAXATIVE

Compare to the active ingredient  
in Dulcolax Suppositories\*

**12  
SUPPOSITORIES**

TAMPER EVIDENT: SUPPOSITORIES ARE WRAPPED IN TAMPER EVIDENT SEALED WRAPPERS. DO NOT USE IF WRAPPER IS TORN OR OPEN WHEN PURCHASED

Rev: 04/26

**Aarna USA**  
Manufactured for: AARNA USA Inc. Leland, NC-28451  
[www.aarnaus.com](http://www.aarnaus.com)  
Questions or comments? toll-free (877)225-6999

Made in USA

3 82568 00282 6

\*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, owner of the registered trademark Dulcolax®



## BISACODYL

bisacodyl suppository

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:82568-0028
<b>Route of Administration</b>	RECTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	10 mg

### Inactive Ingredients

Ingredient Name	Strength
HYDROGENATED PALM KERNEL OIL (UNII: FM8D1RE2VP)	

## Product Characteristics

<b>Color</b>	white (WHITE TO OFF WHITE)	<b>Score</b>	
<b>Shape</b>	OVAL (TORPEDO SHAPE)	<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82568-0028-2	12 in 1 BOX	04/21/2026	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:82568-0028-5	50 in 1 BOX	04/21/2026	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	04/21/2026	

**Labeler** - AARNA USA INC. (118515992)

Revised: 4/2026

AARNA USA INC.