ARTHRITIS PAIN RELIEVER- acetaminophen tablet, film coated, extended release Walgreen Company

waigi een Com

Walgreen Co. Arthritis Pain Reliever Drug Facts

Active ingredient (in each caplet)

Acetaminophen 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults	 take 2 caplets every 8 hours with water swallow whole; do not crush, chew, split or dissolve do not take more than 6 caplets in 24 hours do not use for more than 10 days unless directed by a doctor
under 18 years of age	ask a doctor

Other information

- store at 20-25°C (68-77°F).
- do not use if printed foil under cap is broken or missing

• meets the requirements of USP Dissolution Test 2

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Walgreens

WALGREENS PHARMACIST RECOMMENDED

Compare to the active ingredient in Tylenol® 8HR Arthritis Pain

Arthritis Pain Reliever

ACETAMINOPHEN EXTENDED-RELEASE TABLETS, 650 mg

PAIN RELIEVER / FEVER REDUCER

Arthritis

8 Hour

• For the temporary relief of minor arthritis pain

24 CAPLETS* 650 mg EACH (*capsule-shaped tablets)

ACTUAL SIZE



Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

'This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Tylenol® BHR Arthritis Pain. DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
WAIGreens.com ©2021 Walgreen Co. Patent 7,897,172

W3ORG1221-F

Drug Facts

Active ingredient (in each caplet)

Purpose

Acetaminophen 650 mg.......Pain reliever/fever reducer

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- hackache
- premenstrual and menstrual cramps
- the common cold headache toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains aceta minophen. Se vere liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening blisters rash
- If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do notuse

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Aska doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- rednessor swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for

adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

■ do not take more than directed (see over dose warning)

adults

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
 - do not take more than 6 cap lets in 24 hours
 - do not use for more than 10 days unless directed by a doctor

under 18 years ■ ask a doctor

- Other information ■store at 20-25°C (68-77°F) ■ do not use if printed foil under cap is broken or
- meets the requirements of USP Dissolution Test 2

Inactive ingredients camauta wax, colloidal silicon dioxide, croscarm ellose so dium, hypromellose, magnesium stearate, mattodextrin, microcrystalline ce llulose, polye thylene gtycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments? 1-800-719-9260



DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

ITEM 358600 W10106-0722-L



CODE AREA

54462 94 C7

ARTHRITIS PAIN RELIEVER

acetaminophen tablet, film coated, extended release

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0363-0544

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) ACETAMINOPHEN 650 mg

Inactive Ingredients

l	Ingredient Name	Strength
ı	CARNAUBA WAX (UNII: R12CBM0EIZ)	
ı	SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

CROSCARMELLOSE SODIUM (UNII: M280L1HH48)
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

MAGNESIUM STEARATE (UNII: 70097M6I30)

MALTODEXTRIN (UNII: 7CVR7L4A2D)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

POLYSORBATE 80 (UNII: 60ZP39ZG8H)

POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)

STEARIC ACID (UNII: 4ELV7Z65AP)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	L544
Contains			

Packaging

	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0544- 71	1 in 1 CARTON	02/11/2011	01/31/2022
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0544- 47	1 in 1 CARTON	02/11/2011	04/23/2013
		150 in 1 BOTTLE; Type 0: Not a Combination		

4		Product		
3	NDC:0363-0544- 85	1 in 1 CARTON	02/04/2011	03/23/2017
3		250 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0363-0544- 78	1 in 1 CARTON	07/31/2014	
4		100 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:0363-0544- 62	1 in 1 CARTON	08/15/2014	
5		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075077	05/26/2000	

Labeler - Walgreen Company (008965063)

Revised: 11/2022 Walgreen Company