MOTION SICKNESS RELIEF- meclizine hcl tablet Amerisource Bergen

Good Neighbor Pharmacy 44-403

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

Warnings

Do not use

for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- take first dose 1/2 to 1 hour before starting activity
- adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from heat and humidity
- see end flap for expiration date and lot number

Inactive ingredients

colloidal silicon dioxide, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, pregelatinized starch

Questions or comments?

1-800-426-9391

Principal display panel

GOOD NEIGHBOR PHARMACY®

Compare to Dramamine® All Day Less Drowsy active ingredient*

NDC 46122-535-51

Motion Sickness Relief

meclizine HCl 25 mg / antiemetic

Less Drowsy Formula

Helps prevent Nausea, Vomiting and Dizziness due to Motion Sickness for up to 24 hours

Ages 12 Years and Over

8 Tablets | 25 mg Each

Actual Size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

^{*}This product is not manufactured or distributed by Medtech

Products Inc., owner of the registered trademark Dramamine® Less Drowsy. 50844 ORG042340319

Distributed By AmerisourceBergen 1 West First Avenue Conshohocken, PA 19428 Questions or Concerns? www.mygnp.com



Motion Sickness Relief

Motion Sickness Relief

Less Drowsy Formula

Compare to Dramamine® Less Drowsy active ingredient®



NDC 46122-535-51

Motion Sickness Relief

meclizine HCl 25 mg / antiemetic

Less Drowsy Formula

Helps prevent Nausea, Vomiting and Dizziness due to Motion Sickness for up to 24 hours

Ages 12 Years and Over

8 Tablets 25 mg Each

Actual Size



8-0M5-403A-1 0R60423403



www.mygnp.com

Distributed By AmericourceBergen 1 West First Avenue Conshohocken, PA 19428 Questions or Concerns?





Motion Relief *This products inc, owner of the registered trademark
Products Inc., owner of the registered trademark
Dramamine® Less Drowsy. 50844 0PR3042340319

Questions or comments? 1-800-426-9391

Inactive ingredients coloidal silicon dioxide, D&C yellow #10 aluminum lake, lactose mono hydrale, magnesium stearate, pregelatini zed starch

- see end flap for expiration date and lot number
 - 15°-30°C (59°-86°F)
 protect from heat and humidity
- store at 25°C (77°F); excursions permitted between
- OTHER INTORMATION TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

daily, or as directed by a doctor

If pregnant or breast-feeding, ask a health professional

- drowsiness

 use caution when driving a motor vehicle or operating
 - a slcohol, sedatives, and tranquilizers may increase
 - sovid alcoholic beverages
 - quowajuesa jusik occort

Sedatives or Tanquilizers.
When using this product

Ask a doctor or pharmacist before use if you are taking

camonora

- gland Boucoms arch as emphysema or chronic
- Ask a doctor before use if you have difficulty in unination due to enlargement of the prostate

directed by a doctor.

Warnings Do not use for children under 12 years of age unless TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

> arnish area o date

MOTION SICKNESS RELIEF

meclizine hcl tablet

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:46122-535

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name

Basis of Strength

MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - MECLIZINE - MECLIZINE

UNII:3L5TQ84570)

HYDROCHLORIDE

25 mg

Inactive Ingredients

Ingredient Name Strength

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)
MAGNESIUM STEARATE (UNII: 70097M6I30)

STARCH, CORN (UNII: O8232NY3SJ)

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;403
Contains			

_		-
שבע	ran	ına
Pac	Ray	шу

" .. _ . Marketing Start Marketing End

+	item code	Раскаде резсприон	Date	Date
1	NDC:46122-535- 51	1 in 1 CARTON	06/24/2002	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M009	06/24/2002			

Labeler - Amerisource Bergen (007914906)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(46122-535)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(46122-535) , pack(46122-535)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(46122-535)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(46122-535)

Revised: 11/2023 Amerisource Bergen