

**TRIPLE ANTIBIOTIC PLUS PAIN RELIEF- bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment
GENOMMA LAB USA, INC.**

TRIPLE ANTIBIOTIC PLUS PAIN RELIEF

Drug Facts

<i>Active ingredients (in each gram)</i>	<i>Purpose</i>
Bacitracin zinc 500 units	First aid antibiotic
Neomycin sulfate 3.5 mg	First aid antibiotic
Polymyxin B sulfate 10,000 units	First aid antibiotic
Pramoxine HCl 10 mg	External analgesic

Uses

first aid to help prevent infection and for the temporary relief of pain in minor:

- cuts
- scrapes
- burns

Warnings

For external use only.

Do not use

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 2 years of age and older:
 - clean the affected area
 - apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
 - may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information

- store at 20° to 25°C (68° to 77°F)
- to open: unscrew cap, pull tab to remove foil seal
- see carton or tube crimp for lot number and expiration date

Inactive ingredient

white petrolatum

Questions?

Call toll free **1-877-994-3666** Monday to Friday 8am to 6pm, Central time

Distributed by:
Genomma Lab USA Inc.
Houston, TX 77098

PRINCIPAL DISPLAY PANEL - 28.4 g Tube Carton

cicatricure[®]

OPEN WOUND

FOR MINOR CUTS, SCRAPES AND BURNS

PAIN, ITCH & SCAR

- Triple Antibiotic Ointment
- Up to 24-Hour Infection Protection
- Maximum Strength: Infection Protection, Pain & Itch Relief
- Helps Healing Wounds, Minimizing the Appearance of Scars**

FIRST AID ANTIBIOTIC

PAIN RELIEVING OINTMENT

Bacitracin Zinc – Neomycin Sulfate – Polyximin B Sulfate – Pramoxine HCl

*Based on published studies. For best results use sterile bandage or gauze.

**Preventing infection helps minimize appearance of scars.

NET WT.

1 oz (28.4 g)

5247116
1223
56



2000013449

cicatricure®

OPEN WOUND

FOR MINOR CUTS, SCRAPES AND BURNS

PAIN, ITCH & SCAR

- Triple Antibiotic Ointment
- Up to 24-Hour Infection Protection
- Maximum Strength: Infection Protection, Pain & Itch Relief
- Helps Healing Wounds, Minimizing the Appearance of Scars**



FIRST AID ANTIBIOTIC PAIN RELIEVING OINTMENT

Bacitracin Zinc - Neomycin Sulfate - Polyximin B Sulfate - Pramoxine HCl

NET WT.
1 oz (28.4 g)

*Based on published studies. For best results use sterile bandage or gauze.
**Preventing infection helps minimize appearance of scars.

Made in Canada
Distributed by:
Genomma Lab USA Inc.
Houston, TX 77098
www.genommalab.us

NO VARNISH/NO AQ
NO COPY / NO COLOR
THIS FLAP FOR LOT #
AND EXP DATE PRINT



104
B272.0
ENG 19.13

OPEN WOUND
FOR MINOR CUTS, SCRAPES
AND BURNS



TRIPLE ANTIBIOTIC PLUS PAIN RELIEF

bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:50066-202

Drug Facts

Active ingredients (in each gram)

Bacitracin zinc 500 units.....First aid antibiotic
 Neomycin sulfate 3.5 mg.....First aid antibiotic
 Polymyxin B sulfate 10,000 units.....First aid antibiotic
 Pramoxine HCl 10 mg.....External analgesic

Uses
 first aid to help prevent infection and for the temporary relief of pain in minor: • cuts • scrapes • burns

Warnings
 For external use only.
Do not use

- if you are allergic to any of the ingredients
- in the eyes • over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds • animal bites • serious burns

Drug Facts (continued)

Stop use and ask a doctor if

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 2 years of age and older:
- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Drug Facts (continued)

Inactive ingredient white petrolatum

Questions?
 Call toll free 1-877-994-3666 Monday to Friday 8am to 6pm, Central time

1201

Pharmaceutical

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN ZINC	500 [USP'U] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ 07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50066-202-01	1 in 1 CARTON	12/21/2023	
1		28.4 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M004	12/21/2023	

Labeler - GENOMMA LAB USA, INC. (832323534)

Revised: 12/2023

GENOMMA LAB USA, INC.