



## **INDICATIONS AND USAGE**

Nystatin Oral Suspension is indicated for the treatment of candidiasis in the oral cavity.

## **CONTRAINDICATIONS**

The preparation is contraindicated in patients with a history of hypersensitivity to any of its components.

## **PRECAUTIONS**

### **General**

This medication is not to be used for the treatment of systemic mycoses. Discontinue treatment if sensitization or irritation is reported during use.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

### **Pregnancy**

### **Teratogenic Effects Category C**

Animal reproduction studies have not been conducted with nystatin oral suspension. It is also not known whether nystatin oral suspension can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin oral suspension should be given to a pregnant woman only if clearly needed.

### **Nursing Mothers**

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

### **Pediatric Use**

See **DOSAGE AND ADMINISTRATION**.

## **ADVERSE REACTIONS**

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported. (See **PRECAUTIONS: General**).

*Gastrointestinal:* Diarrhea (including one case of bloody diarrhea), nausea, vomiting, gastrointestinal upset/disturbances.

*Dermatologic:* Rash, including urticaria has been reported rarely. Stevens-Johnson syndrome has been reported very rarely.

*Other:* Tachycardia, bronchospasm, facial swelling, and non-specific myalgia have also been rarely reported.

**To report SUSPECTED ADVERSE REACTIONS, contact Kesin Pharma at 1-833-537-4679 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **OVERDOSAGE**

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects of superinfections (see **CLINICAL PHARMACOLOGY, Pharmacokinetics**).

## **DOSAGE AND ADMINISTRATION**

**INFANTS:** 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

**NOTE:** Limited clinical studies in premature and low birth weight infants indicate that 1 mL four times daily is effective.

**CHILDREN AND ADULTS:** 4 to 6 mL (400,000 to 600,000 units) four times daily (one-half of dose in each side of mouth). The preparation should be retained in the mouth as long as possible before swallowing.

Continue treatment for at least 48 hours after perioral symptoms have disappeared and cultures demonstrate eradication of *Candida albicans*.

## **HOW SUPPLIED**

Nystatin Oral Suspension, USP 100,000 USP Nystatin Units per mL, is available as a cherry-mint flavored, yellow, ready-to-use suspension, supplied in the following oral dosage forms:

NDC 81033-015-60: 60 mL bottle with a 1 mL calibrated dropper

NDC 81033-015-16: 16 oz (473 mL) bottle

## **Rx Only**

## **SHAKE WELL BEFORE USE**

## **SHAKE WELL BEFORE USE**

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15° C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

AVOID FREEZING

**Pharmacists:** Dispense in a tight, light-resistant container as defined in the USP.

Distributed by:

**Kesin Pharma**

Oldsmar, FL 34677

Revised: 12/2025

PI-015B-V01



## Nystatin 16 oz (473 mL)

**Rx Only**

NDC 81033- **015**-16

**Nystatin Oral Suspension, USP**

**SHAKE WELL BEFORE USING**

The image shows a detailed view of the product label for Nystatin Oral Suspension, USP. The label is white with yellow and blue accents. It features the Kasim logo (a stylized 'K' with a vertical bar) in the top left and bottom left corners. The text on the label includes: 'Rx Only' and 'NDC 81033-015-16' in the top left; 'Nystatin Oral Suspension, USP' in large blue letters in the center; '(100,000 units per mL)' in a yellow box below the product name; 'SHAKE WELL BEFORE USING' and 'Cherry-Mint Flavored For Institutional Use Only' in the middle; 'Net: 473 mL' in the bottom right; and detailed usage instructions, warnings, and distribution information on the right side. A barcode is located on the right side of the label, with the number 'N 81033 01516 3' printed vertically next to it. A red dashed line indicates a 'NO VARNISH ZONE (20 x 25 mm)' on the right edge of the label.

**Rx Only**      NDC 81033-**015**-16

**Nystatin Oral Suspension, USP**

**(100,000 units per mL)**

**SHAKE WELL BEFORE USING**  
Cherry-Mint Flavored  
For Institutional Use Only

**Net: 473 mL**

**Each mL contains:** 100,000 units Nystatin, USP in a vehicle containing cherry flavor, dibasic sodium phosphate heptahydrate, disodium edetate, glycerin, methylparaben, monobasic sodium phosphate monohydrate, peppermint oil, propyl paraben, sodium benzoate, sodium hexametaphosphate, and sucrose.

**Usual Dosage: For Infants:** 2 mL (200,000 units) four times daily (in infants and young children, place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

**Usual Dosage: For Children and Adults:** See package insert for full prescribing information.

**Warnings:** Keep this and all drugs out of reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

**AVOID FREEZING.**

Dispense in a tight light-resistant container as defined in the USP. Do not use if inner foil seal printed "SEALED FOR YOUR PROTECTION" is broken or missing.

**Distributed by:**  
Kasim Pharma  
Oldsmar, FL 34677      Rev. 01/2026

N 81033 01516 3

NO VARNISH ZONE  
(20 x 25 mm)

## Nystatin 60 mL Bottle and Carton Labels

**Rx Only**

NDC 81033- **015**-60

**Nystatin Oral Suspension, USP**

**SHAKE WELL BEFORE USING**

Rx Only

NDC 81033-015-60

# Nystatin Oral Suspension, USP

(100,000 units per mL)

SHAKE WELL BEFORE USING  
Cherry-Mint Flavored  
For Institutional Use Only

Net: 60 mL

**Each mL contains:** 100,000 units Nystatin, USP in a vehicle containing cherry flavor, dibasic sodium phosphate heptahydrate, disodium edetate, glycerin, methylparaben, monobasic sodium phosphate monohydrate, peppermint oil, propyl paraben, sodium benzoate, sodium hexametaphosphate, and sucrose.

**Usual Dosage: For Infants:** 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

**Usual Dosage: For Children and Adults:** See package insert for full prescribing information.

**Warnings: Keep this and all drugs out of reach of children.** In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

**Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].**

**AVOID FREEZING.**

Dispense in a tight light-resistant container as defined in the USP.

**Before dispensing, replace cap with safety dropper.**

**Distributed by:**

Kesin Pharma  
Oldsmar, FL 34677

Rev. 12/2025



NO VARNISH ZONE  
(20 x 25 mm)

Rx Only

NDC 81033- 015-60

Nystatin Oral Suspension, USP

SHAKE WELL BEFORE USING

(Includes safe cap dropper)



## NYSTATIN

nystatin suspension

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:81033-015(NDC:85742-013)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>NYSTATIN</b> (UNII: BDF1O1C72E) (NYSTATIN - UNII:BDF1O1C72E)	NYSTATIN	100000 U in 1 mL		
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>		<b>Strength</b>		
EDETATE DISODIUM (UNII: 7FLD91C86K)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
HEXASODIUM HEXAMETAPHOSPHATE (UNII: N40N91DW96)				
SODIUM PHOSPHATE, DIBASIC, HEPTAHYDRATE (UNII: 70WT22SF4B)				
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)				
GLYCERIN (UNII: PDC6A3C0OX)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
SUCROSE (UNII: C151H8M554)				
PEPPERMINT OIL (UNII: AV092KU4JH)				
<b>Product Characteristics</b>				
<b>Color</b>	yellow	<b>Score</b>		
<b>Shape</b>		<b>Size</b>		
<b>Flavor</b>	CHERRY (MINT)	<b>Imprint Code</b>		
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:81033-015-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/30/2026	
2	NDC:81033-015-60	1 in 1 CARTON	08/31/2026	
2		60 mL in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANDA	ANDA062832	08/14/2020		

**Labeler** - Kesin Pharma Corporation (117447816)