

# DOXAZOSIN- doxazosin tablet

## Rising Pharma Holdings, Inc.

-----

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DOXAZOSIN TABLETS safely and effectively. See full prescribing information for DOXAZOSIN TABLETS.

DOXAZOSIN tablets, for oral use  
Initial U.S. Approval: 1990

### INDICATIONS AND USAGE

Doxazosin tablets are an alpha<sub>1</sub> adrenergic antagonist indicated for: (1)

- Signs and symptoms of Benign Prostatic Hyperplasia (BPH)
- Treatment of Hypertension

### DOSAGE AND ADMINISTRATION

- For the treatment of BPH: Initiate therapy at 1 mg once daily. Dose maybe titrated at 1 to 2-week intervals, up to 8 mg once daily. (2.2)
- For the treatment hypertension: Initiate therapy at 1 mg once daily. Dose may be titrated as needed, up to 16 mg once daily. (2.3)

### DOSAGE FORMS AND STRENGTHS

- Tablets: 1 mg, 2 mg, 4 mg, 8 mg.

### CONTRAINDICATIONS

- Hypersensitivity to doxazosin, other quinazolines, or any other ingredient in doxazosin tablets. (4)

### WARNINGS AND PRECAUTIONS

- Postural hypotension with or without syncope may occur. (5.1)
- Risk of Intraoperative Floppy Iris Syndrome during cataract surgery. (5.2)
- Screen for the presence of prostate cancer prior to treatment for BPH and at regular intervals afterwards. (5.3)

### ADVERSE REACTIONS

The most commonly reported adverse reactions from clinical trials are Fatigue, malaise, hypotension, and dizziness. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Rising Pharma Holdings, Inc. at 1-844-874-7464 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### DRUG INTERACTIONS

- Strong cytochrome P450 (CYP) 3A inhibitors may increase exposure to doxazosin and increased risk of hypotension (7.1)
- Concomitant administration of doxazosin tablets with a phosphodiesterase-5 (PDE-5) inhibitor can result in additive blood pressure lowering effects and symptomatic hypotension. (7.2)

### USE IN SPECIFIC POPULATIONS

- Hepatic Impairment: Monitor for hypotension. (8.6, 12.3)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 5/2024

---

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

- 1.1 Benign Prostatic Hyperplasia (BPH)
- 1.2 Hypertension

## **2 DOSAGE AND ADMINISTRATION**

- 2.1 Dosing Information
- 2.2 Benign Prostatic Hyperplasia
- 2.3 Hypertension

## **3 DOSAGE FORMS AND STRENGTHS**

## **4 CONTRAINDICATIONS**

## **5 WARNINGS AND PRECAUTIONS**

- 5.1 Postural Hypotension
- 5.2 Cataract Surgery
- 5.3 Prostate Cancer
- 5.4 Priapism

## **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

## **7 DRUG INTERACTIONS**

- 7.1 CYP 3A Inhibitors
- 7.2 Phosphodiesterase-5 (PDE-5) inhibitors

## **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and Pharmacology

## **14 CLINICAL STUDIES**

- 14.1 Benign Prostatic Hyperplasia (BPH)
- 14.2 Hypertension

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

#### **1.1 Benign Prostatic Hyperplasia (BPH)**

Doxazosin tablets are indicated for the treatment of the signs and symptoms of BPH.

## **1.2 Hypertension**

Doxazosin tablets are indicated for the treatment of hypertension, to lower blood pressure. Lowering blood pressure reduces the risk of fatal and nonfatal cardiovascular events, primarily strokes and myocardial infarctions. These benefits have been seen in controlled trials of antihypertensive drugs from a wide variety of pharmacologic classes, including this drug.

Control of high blood pressure should be part of comprehensive cardiovascular risk management, including, as appropriate, lipid control, diabetes management, antithrombotic therapy, smoking cessation, exercise, and limited sodium intake. Many patients will require more than one drug to achieve blood pressure goals. For specific advice on goals and management, see published guidelines, such as those of the National High Blood Pressure Education Program's Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC).

Numerous antihypertensive drugs, from a variety of pharmacologic classes and with different mechanisms of action, have been shown in randomized controlled trials to reduce cardiovascular morbidity and mortality, and it can be concluded that it is blood pressure reduction, and not some other pharmacologic property of the drugs, that is largely responsible for those benefits. The largest and most consistent cardiovascular outcome benefit has been a reduction in the risk of stroke, but reductions in myocardial infarction and cardiovascular mortality also have been seen regularly.

Elevated systolic or diastolic pressure causes increased cardiovascular risk, and the absolute risk increase per mmHg is greater at higher blood pressures, so that even modest reductions of severe hypertension can provide substantial benefit. Relative risk reduction from blood pressure reduction is similar across populations with varying absolute risk, so the absolute benefit is greater in patients who are at higher risk independent of their hypertension (for example, patients with diabetes or hyperlipidemia), and such patients would be expected to benefit from more aggressive treatment to a lower blood pressure goal.

Some antihypertensive drugs have smaller blood pressure effects (as monotherapy) in black patients, and many antihypertensive drugs have additional approved indications and effects (e.g., on angina, heart failure, or diabetic kidney disease). These considerations may guide selection of therapy.

Doxazosin tablets may be used alone or in combination with other antihypertensives.

## **2 DOSAGE AND ADMINISTRATION**

### **2.1 Dosing Information**

Following the initial dose and with each dose increase of doxazosin tablets, monitor blood pressure for at least 6 hours following administration. If doxazosin tablets administration is discontinued for several days, therapy should be restarted using the initial dosing regimen.

### **2.2 Benign Prostatic Hyperplasia**

The recommended initial dosage of doxazosin tablets are 1 mg given once daily either in the morning or evening.

Depending on the individual patient's urodynamics and BPH symptomatology, the dose may be titrated at 1 to 2 week intervals to 2 mg, and thereafter to 4 mg and 8 mg once daily. The maximum recommended dose for BPH is 8 mg once daily.

Routinely monitor blood pressure in these patients.

### **2.3 Hypertension**

The initial dosage of doxazosin tablets is 1 mg given once daily. Daily dosage may be doubled up to 16 mg once daily, as needed, to achieve the desired reduction in blood pressure.

## **3 DOSAGE FORMS AND STRENGTHS**

Doxazosin Tablets, USP are available containing doxazosin mesylate, USP equivalent to 1 mg, 2 mg, 4 mg or 8 mg of doxazosin.

- The 1 mg tablets are White to off white, capsule shaped, debossed 'D' '1' on one side and break line on other side.
- The 2 mg tablets are White to off white, capsule shaped, debossed 'D' '2' on one side and break line on other side.
- The 4 mg tablets are White to off white, capsule shaped, debossed 'D' '4' on one side and break line on other side.
- The 8 mg tablets are White to off white, capsule shaped, debossed 'D' '8' on one side and break line on other side.

## **4 CONTRAINDICATIONS**

The use of doxazosin tablets are contraindicated in patients with a hypersensitivity to doxazosin, other quinazolines (e.g., prazosin, terazosin), or any of its components.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Postural Hypotension**

Postural hypotension with or without symptoms (e.g., dizziness) may develop within a few hours following administration of doxazosin tablets. However, infrequently, symptomatic postural hypotension has also been reported later than a few hours after dosing. As with other alpha-blockers, there is a potential for syncope, especially after the initial dose or after an increase in dosage strength. Advise patient how to avoid symptoms resulting from postural hypotension and what measures to take should they develop.

Concomitant administration of doxazosin tablets with a PDE-5 inhibitor can result in additive blood pressure lowering effects and symptomatic hypotension.

### **5.2 Cataract Surgery**

Intraoperative Floppy Iris Syndrome (IFIS) has been observed during cataract surgery in some patients on or previously treated with alpha<sub>1</sub> blockers. This variant of small pupil syndrome is characterized by the combination of a flaccid iris that billows in response to

intraoperative irrigation currents, progressive intraoperative miosis despite preoperative dilation with standard mydriatic drugs, and potential prolapse of the iris toward the phacoemulsification incisions. The patient's surgeon should be prepared for possible modifications to their surgical technique, such as the utilization of iris hooks, iris dilator rings, or viscoelastic substances. There does not appear to be a benefit of stopping alpha<sub>1</sub> blocker therapy prior to cataract surgery.

### 5.3 Prostate Cancer

Carcinoma of the prostate causes many of the symptoms associated with BPH and the two disorders frequently co-exist.

Carcinoma of the prostate should therefore be ruled out prior to commencing therapy with doxazosin tablets for the treatment of BPH.

### 5.4 Priapism

Alpha<sub>1</sub> antagonists, including doxazosin, have been associated with priapism (painful penile erection, sustained for hours and unrelieved by sexual intercourse or masturbation). This condition can lead to permanent impotence if not promptly treated.

## 6 ADVERSE REACTIONS

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### *Benign Prostatic Hyperplasia (BPH)*

The incidence of adverse events has been ascertained from worldwide clinical trials in 965 BPH patients. The incidence rates presented below (Table 2) are based on combined data from seven placebo-controlled trials involving once-daily administration of doxazosin tablets in doses of 1 to 16 mg in hypertensives and 0.5 to 8 mg in normotensives. Adverse reactions occurring more than 1% more frequently in BPH patients treated with doxazosin tablets vs placebo are summarized in Table 1.

**Table 1. Adverse Reactions Occurring more than 1% More Frequently in BPH Patients Treated with Doxazosin tablets Versus Placebo**

<b>BODY SYSTEM</b>	<b>Doxazosin Tablets N = 665</b>	<b>Placebo N = 300</b>
<b>NERVOUS SYSTEM DISORDERS</b>		
Dizziness*	15.6%	9.0%
Somnolence	3.0%	1.0%
<b>CARDIAC DISORDERS</b>		
Hypotension	1.7%	0%
<b>RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS</b>		

Dyspnoea	2.6%	0.3%
<b>GASTROINTESTINAL DISORDERS</b>		
Dry Mouth	1.4%	0.3%
<b>GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS</b>		
Fatigue	8.0%	1.7%
Oedema	2.7%	0.7%

\* Includes vertigo

Other adverse reactions occurring less than 1% more frequently in BPH patients treated with doxazosin tablets vs placebo but plausibly related to doxazosin tablets include: palpitations.

### *Hypertension*

Doxazosin tablets has been administered to approximately 4000 hypertensive patients in clinical trials, of whom 1679 were included in the hypertension clinical development program. In placebo-controlled studies, adverse events occurred in 49% and 40% of patients in the doxazosin and placebo groups, respectively, and led to discontinuation in 2% of patients in each group.

Adverse reactions occurring more than 1% more frequently in hypertensive patients treated with doxazosin tablets vs placebo are summarized in Table 1. Postural effects and edema appeared to be dose-related. The prevalence rates presented below are based on combined data from placebo-controlled studies involving once-daily administration of doxazosin at doses ranging from 1 to 16 mg.

**Table 2. Adverse Reactions Occurring more than 1% More Frequently in Hypertensive Patients Treated with Doxazosin tablets versus Placebo**

<b>BODY SYSTEM</b>	<b>Doxazosin Tablets N = 339</b>	<b>Placebo N = 336</b>
<b>NERVOUS SYSTEM DISORDERS</b>		
Dizziness	19%	9%
Somnolence	5%	1%
<b>RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS</b>		
Rhinitis	3%	1%
<b>RENAL AND URINARY DISORDERS</b>		
Polyuria	2%	0%
<b>REPRODUCTIVE SYSTEM AND BREAST DISORDERS</b>		
<b>GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS</b>		
Fatigue / Malaise	12%	6%

Other adverse reactions occurring less than 1% more frequently in hypertensive

patients treated with doxazosin tablets vs placebo but plausibly related to doxazosin tablets use include vertigo, hypotension, hot flushes, epistaxis and oedema.

Doxazosin tablets has been associated with decreases in white blood cell counts.

*Laboratory changes observed in clinical studies*

Leukopenia/Neutropenia: Decreases in mean white blood cell (WBC) and mean neutrophil count were observed in controlled clinical trials of hypertensive patients receiving doxazosin tablets. In cases where follow-up was available, WBC and neutrophil counts returned to normal after discontinuation of doxazosin tablets. No patients became symptomatic as a result of the low WBC or neutrophil counts.

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post-approval use of doxazosin tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

In post-marketing experience, the following additional adverse reactions have been reported:

*Blood and Lymphatic System Disorders:* leukopenia, thrombocytopenia;

*Immune System Disorders:* allergic reaction;

*Nervous System Disorders:* hypoesthesia;

*Eye Disorders:* Intraoperative Floppy Iris Syndrome [see Warnings and precautions (5.4)];

*Cardiac Disorders:* bradycardia;

*Respiratory, Thoracic and Mediastinal Disorders:* bronchospasm aggravated;

*Gastrointestinal Disorders:* vomiting;

*Hepatobiliary Disorders:* cholestasis, hepatitis cholestatic;

*Skin and Subcutaneous Tissue Disorders:* urticaria;

*Musculoskeletal and Connective Tissue Disorders:* muscle cramps, muscle weakness;

*Renal and Urinary Disorders:* hematuria, micturition disorder, micturition frequency, nocturia;

*Reproductive System and Breast Disorders:* gynecomastia, priapism.

## **7 DRUG INTERACTIONS**

### **7.1 CYP 3A Inhibitors**

*In vitro* studies suggest that doxazosin is a substrate of CYP 3A4. Strong CYP3A inhibitors may increase exposure to doxazosin. Monitor blood pressure and for symptoms of hypotension when doxazosin tablets are used concomitantly with strong CYP3A inhibitors [see Clinical Pharmacology (12.3)].

## 7.2 Phosphodiesterase-5 (PDE-5) inhibitors

Concomitant administration of doxazosin tablets with a phosphodiesterase-5 (PDE-5) inhibitor can result in additive blood pressure lowering effects and symptomatic hypotension. Monitor blood pressure and for symptoms of hypotension [*see Warnings and Precautions (5.1)*].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

The limited available data with doxazosin tablets in pregnant women are not sufficient to inform a drug-associated risk for major birth defects and miscarriage. However, untreated hypertension during pregnancy can result in increased maternal risks [*see Clinical Considerations*]. In animal reproduction studies, no adverse developmental effects were observed when doxazosin was orally administered to pregnant rabbits and rats during the period of organogenesis at doses of up to 41 and 20 mg/kg, respectively (exposures in rabbits and rats were 10 and 4 times, respectively, the human AUC exposures with a 12 mg/day therapeutic dose). A dosage regimen of 82 mg/kg/day in the rabbit was associated with reduced fetal survival [*see Data*].

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 - 4% and 15 - 20%, respectively.

#### Clinical Considerations

##### *Disease-associated maternal and/or embryo/fetal risk*

Hypertension in pregnancy increases the maternal risk for pre-eclampsia, gestational diabetes, premature delivery, and delivery complications (e.g., need for cesarean section, and post-partum hemorrhage). Hypertension increases the fetal risk for intrauterine growth restriction and intrauterine death.

#### Data

##### *Animal Data*

Radioactivity was found to cross the placenta following oral administration of labelled doxazosin to pregnant rats. Studies in pregnant rabbits and rats at daily oral doses of up to 41 and 20 mg/kg, respectively (plasma drug concentrations of 10 and 4 times, respectively, the human AUC exposures with a 12 mg/day therapeutic dose), have revealed no evidence of adverse developmental effects. A dosage regimen of 82 mg/kg/day in the rabbit was associated with reduced fetal survival. In peri- and postnatal studies in rats, postnatal development at maternal doses of 40 or 50 mg/kg/day of doxazosin (about 8 times human AUC exposure with a 12 mg/day therapeutic dose) was delayed, as evidenced by slower body weight gain and slightly later appearance of anatomical features and reflexes.

### 8.2 Lactation



## Risk Summary

There is limited information on the presence of doxazosin tablets in human milk [see *Data*]. There is no information on the effects of doxazosin tablets on the breastfeed infant or the effects on milk production.

## Data

A single case study reports that doxazosin tablets are present in human milk, which resulted in an infant dose of less than 1% of the maternal weight-adjusted dosage and a milk/plasma ratio of 0.1. However, these data are insufficient to confirm the presence of doxazosin tablets in human milk.

## **8.4 Pediatric Use**

The safety and effectiveness of doxazosin tablets have not been established in children.

## **8.5 Geriatric Use**

### *Benign Prostatic Hyperplasia (BPH)*

The safety and effectiveness profile of doxazosin tablets was similar in the elderly (age  $\geq$  65 years) and younger (age < 65 years) patients.

### *Hypertension*

Clinical studies of doxazosin tablets did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

## **8.6 Hepatic Impairment**

Doxazosin tablets are extensively metabolized in the liver. Hepatic impairment is expected to increase exposure to doxazosin. Use of doxazosin tablets in patients with severe hepatic impairment (Child-Pugh Class C) is not recommended. Monitor blood pressure and for symptoms of hypotension in patients with lesser degrees of hepatic impairment (Child-Pugh Class A and B) [see *Clinical Pharmacology (12.3)*].

## **10 OVERDOSAGE**


Experience with doxazosin tablets overdosage is limited. Two adolescents, who each intentionally ingested 40 mg doxazosin tablets with diclofenac or acetaminophen, were treated with gastric lavage with activated charcoal and made full recoveries. A two-year-old child who accidentally ingested 4 mg doxazosin tablets was treated with gastric lavage and remained normotensive during the five-hour emergency room observation period. A six-month-old child accidentally received a crushed 1 mg tablet of doxazosin tablets and was reported to have been drowsy. A 32-year-old female with chronic renal failure, epilepsy, and depression intentionally ingested 60 mg doxazosin tablets (blood level = 0.9 mcg/mL; normal values in hypertensives = 0.02 mcg/mL); death was attributed to a

grand mal seizure resulting from hypotension. A 39-year-old female who ingested 70 mg doxazosin tablets, alcohol, and Dalmane<sup>®</sup> (flurazepam) developed hypotension which responded to fluid therapy.

The oral LD<sub>50</sub> of doxazosin is greater than 1000 mg/kg in mice and rats. The most likely manifestation of overdose would be hypotension, for which the usual treatment would be intravenous infusion of fluid. As doxazosin is highly protein bound, dialysis would not be indicated.

## 11 DESCRIPTION

Doxazosin mesylate, USP is a quinazoline compound that is a selective inhibitor of the  $\alpha_1$  subtype of alpha-adrenergic receptors. The chemical name of doxazosin mesylate is 1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-(1,4-benzodioxan-2-ylcarbonyl) piperazine methanesulfonate. The empirical formula for doxazosin mesylate is  $C_{23}H_{25}N_5O_5 \cdot CH_4O_3S$  and the molecular weight is 547.58. It has the following structure:



Doxazosin mesylate, USP is freely soluble in formic acid, very slightly soluble in water and Methanol. Doxazosin tablets are available as colored tablets for oral use and contains 1 mg (white), 2 mg (white), 4 mg (white) and 8 mg (white) of doxazosin as the free base.

The inactive ingredients for all tablets are: lactose monohydrate, microcrystalline cellulose, sodium starch glycolate, sodium lauryl sulfate, Colloidal Silicon dioxide and magnesium stearate.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

#### *Benign Prostatic Hyperplasia (BPH)*

The symptoms associated with benign prostatic hyperplasia (BPH), such as urinary frequency, nocturia, weak stream, hesitancy, and incomplete emptying are related to two components, anatomical (static) and functional (dynamic). The static component is related to an increase in prostate size caused, in part, by a proliferation of smooth muscle cells in the prostatic stroma. However, the severity of BPH symptoms and the degree of urethral obstruction do not correlate well with the size of the prostate. The dynamic component of BPH is associated with an increase in smooth muscle tone in the prostate and bladder neck. The degree of tone in this area is mediated by the  $\alpha_1$  adrenoceptor, which is present in high density in the prostatic stroma, prostatic capsule

and bladder neck. Blockade of the  $\alpha_1$  receptor decreases urethral resistance and may relieve the obstruction and BPH symptoms and improve urine flow.

### *Hypertension*

The mechanism of action of doxazosin tablets are selective blockade of the  $\alpha_1$  (postjunctional) subtype of adrenergic receptors. Studies in normal human subjects have shown that doxazosin competitively antagonized the pressor effects of phenylephrine (an  $\alpha_1$  agonist) and the systolic pressor effect of norepinephrine. Doxazosin and prazosin have similar abilities to antagonize phenylephrine. The antihypertensive effect of doxazosin tablets results from a decrease in systemic vascular resistance. The parent compound doxazosin is primarily responsible for the antihypertensive activity. The low plasma concentrations of known active and inactive metabolites of doxazosin (2-piperazinyloxy, 6'- and 7'-hydroxy and 6- and 7-O-desmethyl compounds) compared to parent drug indicate that the contribution of even the most potent compound (6'-hydroxy) to the antihypertensive effect of doxazosin in man is probably small. The 6'- and 7'-hydroxy metabolites have demonstrated antioxidant properties at concentrations of 5  $\mu\text{M}$ , *in vitro*.

## **12.2 Pharmacodynamics**

### *Benign Prostatic Hyperplasia (BPH)*

Administration of doxazosin tablets to patients with symptomatic BPH resulted in a statistically significant improvement in maximum urinary flow rate [see *Clinical Studies (14.1)*].

### *Effect on Normotensive Patients with Benign Prostatic Hyperplasia (BPH)*

Although blockade of  $\alpha_1$  adrenoceptors also lowers blood pressure in hypertensive patients with increased peripheral vascular resistance, doxazosin tablets treatment of normotensive men with BPH did not result in a clinically significant blood pressure lowering effect (Table 4). The proportion of normotensive patients with a sitting systolic blood pressure less than 90 mmHg and/or diastolic blood pressure less than 60 mmHg at any time during treatment with doxazosin tablets 1 - 8 mg once daily was 6.7% with doxazosin and not significantly different (statistically) from that with placebo (5%).

### *Hypertension*

Administration of doxazosin tablets results in a reduction in systemic vascular resistance. In patients with hypertension, there is little change in cardiac output. Maximum reductions in blood pressure usually occur 2 - 6 hours after dosing and are associated with a small increase in standing heart rate. Like other  $\alpha_1$ -adrenergic blocking agents, doxazosin has a greater effect on blood pressure and heart rate in the standing position.

## **12.3 Pharmacokinetics**

### Absorption

After oral administration of therapeutic doses, peak plasma levels of doxazosin tablets occur at about 2 - 3 hours. Bioavailability is approximately 65%, reflecting first-pass metabolism of doxazosin by the liver. The effect of food on the pharmacokinetics of doxazosin tablets was examined in a crossover study with twelve hypertensive subjects. Reductions of 18% in mean maximum plasma concentration ( $C_{\text{max}}$ ) and 12% in the area

under the concentration-time curve (AUC) occurred when doxazosin tablets were administered with food. Neither of these differences is clinically significant.

In a crossover study in 24 normotensive subjects, the pharmacokinetics and safety of doxazosin were shown to be similar with morning and evening dosing regimens. The AUC after morning dosing was, however, 11% less than that after evening dosing and the time to peak concentration after evening dosing occurred significantly later than that after morning dosing (5.6 vs. 3.5 hours).

### Distribution

At the plasma concentrations achieved by therapeutic doses, approximately 98% of the circulating drug is bound to plasma proteins.

### Metabolism

Doxazosin tablets are extensively metabolized in the liver, mainly by O-demethylation of the quinazoline nucleus or hydroxylation of the benzodioxan moiety. *In vitro* studies suggest that the primary pathway for elimination is via CYP 3A4; however, CYP 2D6 and CYP 2C9 metabolic pathways are also involved to a lesser extent. Although several active metabolites of doxazosin have been identified, the pharmacokinetics of these metabolites have not been characterized.

### Excretion

Plasma elimination of doxazosin is biphasic, with a terminal elimination half-life of about 22 hours. Steady-state studies in hypertensive patients given doxazosin doses of 2 to 16 mg once daily showed linear kinetics and dose proportionality. In two studies, following the administration of 2 mg orally once daily, the mean accumulation ratios (steady-state AUC vs. first-dose AUC) were 1.2 and 1.7. Enterohepatic recycling is suggested by secondary peaking of plasma doxazosin concentrations.

In a study of two subjects administered radiolabelled doxazosin 2 mg orally and 1 mg intravenously on two separate occasions, approximately 63% of the dose was eliminated in the feces and 9% of the dose was found in the urine. On average only 4.8% of the dose was excreted as unchanged drug in the feces and only a trace of the total radioactivity in the urine was attributed to unchanged drug.

### Specific Populations

#### *Geriatric*

The pharmacokinetics of doxazosin tablets in young (<65 years) and elderly (≥65 years) subjects were similar for plasma half-life values and oral clearance.

#### *Renal Impairment*

Pharmacokinetic studies in elderly patients and patients with renal impairment have shown no significant alterations compared to younger patients with normal renal function.

#### *Hepatic Impairment*

Administration of a single 2 mg dose to patients with cirrhosis (Child-Pugh Class A) showed a 40% increase in exposure to doxazosin. The impact of moderate (Child-Pugh Class B) or severe (Child-Pugh Class C) hepatic impairment on the pharmacokinetics of doxazosin is not known [see *Use in Specific Populations (8.6)*].

## Drug Interactions

There are only limited data on the effects of drugs known to influence the hepatic metabolism of doxazosin (e.g., cimetidine).

*Cimetidine:* In healthy volunteers, the administration of a single 1 mg dose of doxazosin on day 1 of a four-day regimen of oral cimetidine (400 mg twice daily) resulted in a 10% increase in mean AUC of doxazosin, and a slight but not significant increase in mean  $C_{max}$  and mean half-life of doxazosin.

*In vitro* data in human plasma indicate that doxazosin tablets have no effect on protein binding of digoxin, warfarin, phenytoin, or indomethacin.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

*Carcinogenesis and Mutagenesis:* Chronic dietary administration (up to 24 months) of doxazosin mesylate at maximally tolerated doses of 40 mg/kg/day in rats and 120 mg/kg/day in mice revealed no evidence of carcinogenic potential. The highest doses evaluated in the rat and mouse studies are associated with AUCs (a measure of systemic exposure) that are 8 times and 4 times, respectively, the human AUC at a dose of 16 mg/day.

Mutagenicity studies revealed no drug- or metabolite-related effects at either chromosomal or subchromosomal levels.

*Fertility in Males:* Studies in rats showed reduced fertility in males treated with doxazosin at oral doses of 20 (but not 5 or 10) mg/kg/day, about 4 times the AUC exposures obtained with a 12 mg/day human dose. This effect was reversible within two weeks of drug withdrawal. There have been no reports of any effects of doxazosin on male fertility in humans.

### **13.2 Animal Toxicology and Pharmacology**

An increased incidence of myocardial necrosis or fibrosis was observed in long-term (6 - 12 months) studies in rats and mice (exposure 8 times human AUC exposure in rats and somewhat equivalent to human  $C_{max}$  exposure in mice). Findings were not seen at lower doses. In dogs no cardiotoxicity was observed following 12 months of oral dosing at doses that resulted in maximum plasma concentrations ( $C_{max}$ ) 14 times the  $C_{max}$  exposure in humans receiving a 12 mg/day therapeutic dose or in Wistar rats at  $C_{max}$  exposures 15 times human  $C_{max}$  exposure. There is no evidence that similar lesions occur in humans.

## **14 CLINICAL STUDIES**

### **14.1 Benign Prostatic Hyperplasia (BPH)**

The efficacy of doxazosin tablets was evaluated extensively in over 900 patients with BPH in double-blind, placebo-controlled trials. Doxazosin tablets treatment was superior to placebo in improving patient symptoms and urinary flow rate. Significant relief with doxazosin tablets were seen as early as one week into the treatment regimen, with

doxazosin tablets-treated patients (N=173) showing a significant (p<0.01) increase in maximum flow rate of 0.8 mL/sec compared to a decrease of 0.5 mL/sec in the placebo group (N=41). In long-term studies, improvement was maintained for up to 2 years of treatment. In 66 - 71% of patients, improvements above baseline were seen in both symptoms and maximum urinary flow rate.

In three placebo-controlled studies of 14 - 16 weeks' duration, obstructive symptoms (hesitation, intermittency, dribbling, weak urinary stream, incomplete emptying of the bladder) and irritative symptoms (nocturia, daytime frequency, urgency, burning) of BPH were evaluated at each visit by patient-assessed symptom questionnaires. The bothersomeness of symptoms was measured with a modified Boyarsky questionnaire. Symptom severity/frequency was assessed using a modified Boyarsky questionnaire or an AUA-based questionnaire. Uroflowmetric evaluations were performed at times of peak (2 - 6 hours post-dose) and/or trough (24 hours post-dose) plasma concentrations of doxazosin tablets.


The results from the three placebo-controlled studies (N=609) showing significant efficacy with 4 mg and 8 mg doxazosin are summarized in Table 3. In all three studies, doxazosin tablets resulted in statistically significant relief of obstructive and irritative symptoms compared to placebo. Statistically significant improvements of 2.3 - 3.3 mL/sec in maximum flow rate were seen with doxazosin tablets in Studies 1 and 2, compared to 0.1 - 0.7 mL/sec with placebo.

**Table 3**

SUMMARY OF EFFECTIVENESS DATA IN PLACEBO-CONTROLLED TRIALS

	SYMPTOM SCORE*			MAXIMUM FLOW RATE (mL/sec)		
	N	MEAN	MEAN <sup>b</sup>	N	MEAN	MEAN <sup>b</sup>
		BASELINE	CHANGE		BASELINE	CHANGE
<b>STUDY 1 (Titration to maximum dose of 8 mg)<sup>a</sup></b>						
Placebo	47	15.6	-2.3	41	9.7	+0.7
Doxazosin tablets	49	14.5	-4.9**	41	9.8	+2.9**
<b>STUDY 2 (Titration to fixed dose -14 weeks)<sup>a</sup></b>						
Placebo	37	20.7	-2.5	30	10.6	+0.1
Doxazosin tablets 4 mg	38	21.2	-5.0**	32	9.8	+2.3*
Doxazosin tablets 8 mg	42	19.9	-4.2*	36	10.5	+3.3**
<b>STUDY 3 (Titration to fixed dose -12 weeks)</b>						
Placebo	47	14.9	-4.7	44	9.9	+2.1
Doxazosin tablets 4 mg	46	16.6	-6.1*	46	9.6	+2.6


\*AUA questionnaire (range 0-30) in studies 1 and 3.  
 Modified boyarsky Questionnaire (range 7-39) in study 2.  
<sup>b</sup>Change is to endpoint.  
<sup>c</sup>Change is to fixed-dose efficacy phase, 22-26 hours post for studies 1 and 3 and 2-6 hours post-dose for study 2.  
<sup>d</sup>Study in hypertensives with BPH  
<sup>e</sup>36 patients received a dose of 8 mg Doxazosin tablets<sup>e</sup>  
 \*(\*)P<0.05 (0.01) compared to placebo mean change.



In one fixed-dose study (Study 2), doxazosin tablets therapy (4 to 8 mg, once daily) resulted in a significant and sustained improvement in maximum urinary flow rate of 2.3 - 3.3 mL/sec (Table 3) compared to placebo (0.1 mL/sec). In this study, the only study in which weekly evaluations were made, significant improvement with doxazosin tablets vs. placebo was seen after one week. The proportion of patients who responded with a maximum flow rate improvement of ≥3 mL/sec was significantly larger with doxazosin tablets (34 - 42%) than placebo (13 - 17%). A significantly greater improvement was also

seen in average flow rate with doxazosin tablets (1.6 mL/sec) than with placebo (0.2 mL/sec). The onset and time course of symptom relief and increased urinary flow from Study 1 are illustrated in Figure 1.

**Figure 1 - Study 1**



## 14.2 Hypertension

In a pooled analysis of placebo-controlled hypertension studies with about 300 hypertensive patients per treatment group, doxazosin, at doses of 1 to 16 mg given once daily, lowered blood pressure at 24 hours by about 10/8 mmHg compared to placebo in the standing position and about 9/5 mmHg in the supine position. Peak blood pressure effects (1 - 6 hours) were larger by about 50 - 75% (i.e., trough values were about 55 - 70% of peak effect), with the larger peak-trough differences seen in systolic pressures. There was no apparent difference in the blood pressure response of Caucasians and blacks or of patients above and below age 65. In the same patient population, patients receiving doxazosin tablets gained a mean of 0.6 kg compared to a mean loss of 0.1 kg for placebo patients.

**Table 4. Mean Changes in Blood Pressure from Baseline to the Mean of the Final Efficacy Phase in Normotensives (Diastolic BP < 90 mmHg) in Two Double-blind, Placebo-controlled U.S. Studies with Doxazosin Tablets 1 to 8 mg once daily**

	PLACEBO (N = 85)		DOXAZOSIN TABLETS (N = 183)	
<b>Sitting BP (mmHg)</b>	<b>Baseline</b>	<b>Change</b>	<b>Baseline</b>	<b>Change</b>
Systolic	128.4	-1.4	128.8	-4.9*
Diastolic	79.2	-1.2	79.6	-2.4*
<b>Standing BP (mmHg)</b>	<b>Baseline</b>	<b>Change</b>	<b>Baseline</b>	<b>Change</b>
Systolic	128.5	-0.6	128.5	-5.3*
Diastolic	80.5	-0.7	80.4	-2.6*

\* p ≤ 0.05 compared to placebo

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Doxazosin Tablets, USP are available as scored tablets for oral administration. Each tablet contains doxazosin mesylate, USP equivalent to 1 mg (white), 2 mg (white), 4 mg (white) or 8 mg (white) of the active constituent, doxazosin.

The 1 mg tablets are White to off white, capsule shaped, debossed 'D' '1' on one side and break line on other side. They are supplied as follows:

Bottles of 100 tablets NDC 16571- 165-01

The 2 mg tablets are White to off white, capsule shaped, debossed 'D' '2' on one side and break line on other side. They are supplied as follows:

Bottles of 100 tablets NDC 16571- 166-01

The 4 mg tablets are White to off white, capsule shaped, debossed 'D' '4' on one side and break line on other side. They are supplied as follows:

Bottles of 100 tablets NDC 16571- 167-01

The 8 mg tablets are White to off white, capsule shaped, debossed 'D' '8' on one side and break line on other side. They are supplied as follows:

Bottles of 100 tablets NDC 16571- 168-01

Recommended Storage: Store at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Preserve in tight containers.

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

### *Postural Hypotension*

Advise patients of the possibility of syncopal and orthostatic symptoms, especially at the initiation of therapy, and urged to avoid driving or hazardous tasks for 24 hours after the first dose, after a dosage increase, and after interruption of therapy when treatment is resumed. Advise patients to report symptoms to their healthcare provider.

### *Priapism*

Advise patients of the possibility of priapism and to seek immediate medical attention if symptoms occur.

### **Manufactured for:**

Rising Pharma Holdings, Inc.  
East Brunswick, NJ 08816

Made in India

Code No. DRUGS/TS/23/2007

**Issued:** 04/2024



## **PATIENT INFORMATION**

### **DOXAZOSIN TABLETS, USP**

(dox az' oh sin mes' i late)

#### **What are doxazosin tablets?**

Doxazosin tablets are a prescription medicine that contains doxazosin mesylate and is called an "alpha-blocker". Doxazosin tablets are used to treat:

- the symptoms of benign prostatic hyperplasia (BPH)
- high blood pressure (hypertension)

It is not known if doxazosin tablets are safe and effective in children.

#### **Who should not take doxazosin tablets?**

##### **Do not take doxazosin tablets if you:**

- are allergic to doxazosin, other quinazolines, or any of the ingredients in doxazosin tablets. See the end of this Patient Information leaflet for a complete list of ingredients in doxazosin tablets.

#### **What should I tell my healthcare provider before taking doxazosin tablets?**

##### **Before taking doxazosin tablets, tell your healthcare provider about all of your medical conditions, including if you:**

- have had low blood pressure, especially after taking other medicine. Signs of low blood pressure include fainting, dizziness, and lightheadedness.
- have any planned eye surgery
- have prostate cancer or a history of prostate cancer. Your healthcare provider may have you checked for prostate cancer before you start taking and while you take doxazosin tablets.
- have liver problems
- are pregnant or plan to become pregnant. It is not known if doxazosin tablets will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if doxazosin passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take doxazosin tablets.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Doxazosin tablets may affect the way other medicines work, and other medicines may affect the way doxazosin tablets work causing side effects.

##### **Especially tell your healthcare provider if you take:**

- other medicine for high blood pressure or medicine to treat erectile dysfunction (ED) called a phosphodiesterase type 5 (PDE-5) inhibitor. The use of doxazosin tablets with PDE-5 inhibitors can lead to a drop in blood pressure or to fainting.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

## How should I take doxazosin tablets?

- Take doxazosin tablets exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much doxazosin tablets to take and when to take it.
- Your healthcare provider may need to change your dose of doxazosin tablets until it is the right dose for you.

## What should I avoid while taking doxazosin tablets?

Do not drive or perform any hazardous task until at least 24 hours after you have taken doxazosin tablets if you are taking:

- your first dose of doxazosin tablets
- Doxazosin tablets for the first time **after** your healthcare provider has increased your dose of doxazosin tablets
- Doxazosin tablets for the first time **after** any breaks (interruptions) in your treatment with doxazosin tablets

## What are the possible side effects of doxazosin tablets?

### Doxazosin tablets may cause serious side effects, including:

- **A sudden drop in blood pressure**, especially when you first start treatment or when there is an increase in your dose of doxazosin tablets, is common but can also be serious. This may cause you to faint, or to feel dizzy or lightheaded. Your risk of having this problem may be increased if you take doxazosin tablets with certain other medicines that lower blood pressure including PDE-5 inhibitors. Your healthcare provider may monitor your blood pressure while you take doxazosin tablets. See "What should I avoid while taking doxazosin tablets?"
- **Eye problems during cataract surgery.** A condition called Intraoperative Floppy Iris Syndrome (IFIS) can happen during cataract surgery if you take or have taken alpha-blockers such as doxazosin tablets. If you need to have cataract surgery, be sure to tell your healthcare provider if you take or have taken doxazosin tablets.
- **A painful erection that will not go away.** Doxazosin tablets can cause a painful erection (priapism), which cannot be relieved by having sex. If this happens, get medical help right away. If priapism is not treated, you may not be able to get an erection in the future.

The most common side effects of doxazosin tablets are:

- weakness or lack of energy (asthenia)
- dizziness

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of doxazosin tablets. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

## General information about the safe and effective use of doxazosin tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use doxazosin tablets for a condition for which it was not prescribed. Do not give doxazosin tablets to other people, even if they have the same

symptoms you have. It may harm them.

This Patient Information leaflet summarizes the most important information about doxazosin tablets. For more information, ask your healthcare provider. You can ask your healthcare provider or pharmacist for information that is written for healthcare professionals.

### **What are the ingredients in Doxazosin Tablets?**

The size, shape and appearance of the tablet that you receive may vary, all the tablets have the same active ingredient, and this will not affect the way that the medicine works. You can identify the tablet that you have from the following information.

<b>Strength</b>	<b>Description</b>
1 mg	White to off white, capsule shaped, debossed 'D' '1' on one side and break line on other side.
2 mg	White to off white, capsule shaped, debossed 'D' '2' on one side and break line on other side.
4 mg	White to off white, capsule shaped, debossed 'D' '4' on one side and break line on other side.
8 mg	White to off white, capsule shaped, debossed 'D' '8' on one side and break line on other side.

**Active ingredient:** doxazosin

**Inactive ingredients:** lactose monohydrate, microcrystalline cellulose, sodium starch glycolate, sodium lauryl sulfate, colloidal silicon dioxide and magnesium stearate.

This Patient Information has been approved by the U.S. Food and Drug Administration

**Manufactured for:**

Rising Pharma Holdings, Inc.  
East Brunswick, NJ 08816

Made in India

Code No. DRUGS/TS/23/2007

**Issued:** 04/2024

PILR16801-00

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL 1 mg**


**NDC16571-165-01**

**Doxazosin Tablets, USP**

**1 mg**

**100 Tablets**

**Rx only**



**Rising**  
PHARMACEUTICALS

NDC 16571-165-01

**Doxazosin  
Tablets, USP**

**1 mg**

100 Tablets

**Rx only**

**Each tablet contains:**  
Doxazosin mesylate, USP.....equivalent  
to 1mg doxazosin.

**DOSAGE AND USE:**  
See accompanying package insert for  
full prescribing information.

**Store at 25°C (77°F) excursions  
permitted to 15°C to 30°C (59°F to  
86°F)** [see USP Controlled Room  
Temperature]. Preserve in tight  
containers.

**KEEP THIS AND ALL MEDICATIONS  
OUT OF THE REACH OF CHILDREN.**

**Manufactured for:**  
Rising Pharma Holdings, Inc.  
East Brunswick, NJ 08816

Made in India  
DRUGS/TS/23/2007  
**Issued:** 04/2024  
LR16501-00  
DMBLR004/00

Varnish free area  
for batch details  
will be printed online  
40 x 20 mm

3 16571 16501 8

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL 2 mg**


**NDC 16571-166-01**

**Doxazosin Tablets, USP**

**2 mg**

**100 Tablets**

**Rx only**



**Rising**  
PHARMACEUTICALS

NDC 16571-166-01

**Doxazosin  
Tablets, USP**

**2 mg**

100 Tablets

**Rx only**

**Each tablet contains:**  
Doxazosin mesylate, USP.....equivalent  
to 2mg doxazosin.

**DOSAGE AND USE:**  
See accompanying package insert for  
full prescribing information.

**Store at 25°C (77°F) excursions  
permitted to 15°C to 30°C (59°F to  
86°F)** [see USP Controlled Room  
Temperature]. Preserve in tight  
containers.

**KEEP THIS AND ALL MEDICATIONS  
OUT OF THE REACH OF CHILDREN.**

**Manufactured for:**  
Rising Pharma Holdings, Inc.  
East Brunswick, NJ 08816

Made in India  
DRUGS/TS/23/2007  
**Revised:** 05/2024  
LR16601-01  
DMBLR005/01

Varnish free area  
for batch details  
will be printed online  
40 x 20 mm

3 16571 16601 5

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL 4 mg**

**NDC16571-167-01**

**Doxazosin Tablets, USP**

**4 mg**

**100 Tablets**

**Rx only**

**Rising**<sup>®</sup> PHARMACEUTICALS NDC 16571-167-01

**Doxazosin Tablets, USP**

**4 mg**

100 Tablets **Rx only**

**Each tablet contains:**  
Doxazosin mesylate, USP.....equivalent to 4mg doxazosin.

**DOSAGE AND USE:**  
See accompanying package insert for full prescribing information.

**Store at 25°C (77°F) excursions permitted to 15°C to 30°C (59°F to 86°F)** [see USP Controlled Room Temperature]. Preserve in tight containers.

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

**Manufactured for:**  
Rising Pharma Holdings, Inc.  
East Brunswick, NJ 08816

Made in India  
DRUGS/TS/23/2007  
Revised: 05/2024  
LR16701-01  
DMBLR006/01

Varnish free area for batch details will be printed online 40 x 20 mm

3 16571 16701 2

**NDC16571-168-01**

**Doxazosin Tablets, USP**

**8 mg**

**100 Tablets**

**Rx only**

**Rising**<sup>®</sup> PHARMACEUTICALS NDC 16571-168-01

**Doxazosin Tablets, USP**

**8 mg**

100 Tablets **Rx only**

**Each tablet contains:**  
Doxazosin mesylate, USP.....equivalent to 8mg doxazosin.

**DOSAGE AND USE:**  
See accompanying package insert for full prescribing information.

**Store at 25°C (77°F) excursions permitted to 15°C to 30°C (59°F to 86°F)** [see USP Controlled Room Temperature]. Preserve in tight containers.

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

**Manufactured for:**  
Rising Pharma Holdings, Inc.  
East Brunswick, NJ 08816

Made in India  
DRUGS/TS/23/2007  
Revised: 05/2024  
LR16801-01  
DMBLR007/01

Varnish free area for batch details will be printed online 40 x 20 mm

3 16571 16801 9

## DOXAZOSIN

doxazosin tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:16571-165
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>DOXAZOSIN MESYLATE</b> (UNII: 86P6PQK0MU) (DOXAZOSIN - UNII:NW1291F1W8)	DOXAZOSIN	1 mg		
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>		<b>Strength</b>		
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MICROCRYSTALLINE CELLULOSE 102 (UNII: PNR0YF693Y)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
<b>Product Characteristics</b>				
<b>Color</b>	WHITE	<b>Score</b>	2 pieces	
<b>Shape</b>	CAPSULE	<b>Size</b>	9mm	
<b>Flavor</b>		<b>Imprint Code</b>	D;1	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:16571-165-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/20/2024	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANDA	ANDA212727	05/20/2024		

<b>DOXAZOSIN</b>			
doxazosin tablet			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:16571-166
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>
DOXAZOSIN MESYLATE (UNII: 86P6PQK0MU) (DOXAZOSIN - UNII:NW1291F1W8)		DOXAZOSIN	2 mg
<b>Inactive Ingredients</b>			

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE 102 (UNII: PNR0YF693Y)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

### Product Characteristics

Color	white	Score	2 pieces
Shape	CAPSULE	Size	9mm
Flavor		Imprint Code	D;2
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16571-166-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/20/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA212727	05/20/2024	

## DOXAZOSIN

doxazosin tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:16571-167
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOXAZOSIN MESYLATE (UNII: 86P6PQK0MU) (DOXAZOSIN - UNII:NW1291F1W8)	DOXAZOSIN	4 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE 102 (UNII: PNR0YF693Y)	

**SODIUM LAURYL SULFATE** (UNII: 368GB5141J)

**SODIUM STARCH GLYCOLATE TYPE A** (UNII: H8AV0SQX4D)

**SILICON DIOXIDE** (UNII: ETJ7Z6XBU4)

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	D;4
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16571-167-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/20/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA212727	05/20/2024	

## DOXAZOSIN

doxazosin tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:16571-168
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DOXAZOSIN MESYLATE</b> (UNII: 86P6PQK0MU) (DOXAZOSIN - UNII:NW1291F1W8)	DOXAZOSIN	8 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE 102</b> (UNII: PNR0YF693Y)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	



## Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	D;8
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16571-168-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/20/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA212727	05/20/2024	

**Labeler** - Rising Pharma Holdings, Inc. (116880195)

**Registrant** - Rising Pharma Holdings, Inc. (116880195)

## Establishment

Name	Address	ID/FEI	Business Operations
COHANCE LIFESCIENCES LIMITED		677637710	manufacture(16571-165, 16571-166, 16571-167, 16571-168)

Revised: 6/2024

Rising Pharma Holdings, Inc.