

**PROPARACAINE HYDROCHLORIDE- proparacaine hydrochloride solution/
drops
Sportpharm LLC**

DESCRIPTION


Proparacaine hydrochloride ophthalmic solution USP, 0.5% is a topical local anesthetic for ophthalmic use. The active ingredient is represented by the structural formula:

Established name:

Proparacaine Hydrochloride

Chemical name:

Benzoic acid, 3-amino-4-propoxy-,2-(diethylamino) ethyl ester, monohydrochloride



Molecular weight: 330.86

Each mL contains:

Active:proparacaine hydrochloride 5 mg (0.5%). **Inactives:**glycerin and purified water. The pH may be adjusted with hydrochloric acid and/or sodium hydroxide.

Preservative:benzalkonium chloride (0.01%).

CLINICAL PHARMACOLOGY

Proparacaine hydrochloride ophthalmic solution is a rapidly-acting topical anesthetic, with induced anesthesia lasting approximately 10-20 minutes.

INDICATIONS AND USAGE

Proparacaine hydrochloride ophthalmic solution is indicated for procedures in which a topical ophthalmic anesthetic is indicated; corneal anesthesia of short duration, e.g. tonometry, gonioscopy, removal of corneal foreign bodies and for short corneal and conjunctival procedures.

CONTRAINDICATIONS

Proparacaine hydrochloride ophthalmic solution is contraindicated in patients with known hypersensitivity to any of the ingredients of this preparation.

WARNINGS

NOT FOR INJECTION INTO THE EYE - FOR TOPICAL OPHTHALMIC USE ONLY

Prolonged use of a topical ocular anesthetic is not recommended. It may produce permanent corneal opacification with accompanying visual loss.

PRECAUTIONS

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed to evaluate carcinogenic potential, mutagenicity or possible impairment of fertility in males or females.

Pregnancy

Animal reproduction studies have not been conducted with proparacaine hydrochloride ophthalmic solution USP, 0.5%. It is also not known whether proparacaine hydrochloride can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Proparacaine hydrochloride should be administered to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when proparacaine hydrochloride is administered to a nursing woman.

Pediatric Use

Safety and effectiveness of proparacaine hydrochloride ophthalmic solution in pediatric patients have been established. Use of proparacaine hydrochloride is supported by evidence from adequate and well-controlled studies in adults and children over the age of twelve, and safety information in neonates and other pediatric patients.

Geriatric Use

No overall clinical differences in safety or effectiveness have been observed between the elderly and other adult patients.

ADVERSE REACTIONS

Occasional temporary stinging, burning and conjunctival redness may occur with the use of proparacaine. A rare, severe, immediate-type, apparently hyperallergic corneal reaction characterized by acute, intense and diffuse epithelial keratitis, a gray, ground glass appearance, sloughing of large areas of necrotic epithelium, corneal filaments and sometimes iritis with descemetitis has been reported.

Allergic contact dermatitis from proparacaine with drying and fissuring of the fingertips has been reported.

To report SUSPECTED ADVERSE REACTIONS, contact Bausch & Lomb Incorporated at 1-800-553-5340 or FDA at 1-800-FDA-1088 or

www.fda.gov/medwatch.

DOSAGE AND ADMINISTRATION

Usual Dosage: Removal of foreign bodies and sutures, and for tonometry: 1 to 2 drops (in single instillations) in each eye before operating.

Short corneal and conjunctival procedures: 1 drop in each eye every 5 to 10 minutes for 5 to 7 doses.

NOTE:Proparacaine hydrochloride ophthalmic solution USP, 0.5% should be clear, colorless to faint yellow color. If the solution becomes darker, discard the solution.

FOR TOPICAL OPHTHALMIC USE ONLY

HOW SUPPLIED

Proparacaine hydrochloride ophthalmic solution USP, 0.5% is supplied in a plastic bottle with a controlled drop tip and a white polypropylene cap in the following size:

NDC 85766-057-15 15 mL bottle (reabeled from NDC 24208-730-06)

Storage:

Refrigerate at 2°C to 8°C (36°F to 46°F). Protect from light. Keep tightly closed.

DO NOT USE IF IMPRINTED NECKBAND IS NOT INTACT.
--

Keep out of reach of children.

Distributed by:

Sportpharm LLC

379 Van Ness Ave 1401,

Torrance, CA 90501

Relabeled by:

Enovachem PHARMACEUTICALS

Torrance, CA 90501

PRINCIPAL DISPLAY PANEL

Relabeled For:



2237 N Commerce Parkway,
STE 1,
Weston Florida 33326

Proparacaine Hydrochloride Ophthalmic Solution, USP 0.5%

NDC: 85766-057-15

Qty: 15

Distributed By: Bausch & Lomb Americas Inc.

Source NDC: 24208-730-06

Description: 15mL, proparacaine hydrochloride ophthalmic sterile solution USP, 0.5% in a plastic bottle

Lot #: 00000000

Exp:

Batch #: 00000000

Drug Status: RX



(01) 0 0385766 05715 5

(17)

(10) 00000000

(21)

Proparacaine Hydrochloride Ophthalmic Solution, USP 0.5%

NDC: 85766-057-15

S/N:

Qty: 15

Proparacaine Hydrochloride Ophthalmic Solution, USP 0.5%

NDC: 85766-057-15

S/N:

Qty: 15

Proparacaine Hydrochloride Ophthalmic Solution, USP 0.5%

NDC: 85766-057-15

S/N:

Qty: 15

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

PROPARACAINE HYDROCHLORIDE

proparacaine hydrochloride solution/ drops

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:85766-057(NDC:24208-730)
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PROPARACAINE HYDROCHLORIDE (UNII: U96OL57GOY) (PROPARACAINE - UNII:B40B0JHI1X)	PROPARACAINE HYDROCHLORIDE	5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85766-057-15	1 in 1 CARTON	08/22/2025	
1		15 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

ANDA

ANDA040074

09/29/1995

Labeler - Sportpharm LLC (125298538)

Revised: 4/2026

Sportpharm LLC