

MINADCIN WART REMOVER PATCHES- salicylic acid 17% wart remover patch
Jiangxi Yudexi Pharmaceutical Co., LTD

85248-201

Active Ingredient

Salicylic acid 17%

Purpose

Wart Remover

Use

For the removal of common warts. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface

Warnings

For external use only.

Do not use

On irritated,infected, or reddened skin; on genital warts and warts on the face; on moles, birthmarks, and warts with hair growing from them;on mucous membranes.

When Using

Ask a doctor before use if you have diabetes, poor blood circulation

Stop Use

Ask a doctor before use if you have diabetes, poor blood circulation

Ask Doctor

Ask a doctor before use if you have diabetes, poor blood circulation

Keep Out Of Reach Of Children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

wash affected area: may soak wart in warm water for 5 minutes dry area thoroughly,remove bandage strips by peeling backing off.If necessary,cut medicated patch to fit wart carefully apply medicated plaster, repeat procedure every 48 hours as needed (until wart is removed) for up to 12 week

Other information

Store between 59° and 86°F(15°and 30°C).

Inactive ingredients

hard paraffin,lanolin,phenol.

PRINCIPAL DISPLAY PANEL

MINADCIN
WART REMOVER PATCHES

144 pads

WART REMOVER PATCHES

17% SALICYLIC ACID

TARGETS CORNS & WARTS

EFFECTIVE TREATMENT

CUT-TO-FIT PADS

HAND USE | FOOT USE

MEDICATED PAD

144 PADS

9mm | 12mm

MINADCIN[®]

Drug Facts	Purpose
Active ingredients	Salicylic acid 17%.....Wart Remover

USE
For the removal of common warts. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface

WARNINGS
For external use only.
DO NOT USE
on irritated, infected or reddened skin
on genital warts and warts on the face
on moles, birthmarks and warts with hair growing from them
on mucous membranes
KEEP OUT OF REACH OF CHILDREN If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS
wash affected area: may soak wart in warm water for 5 minutes dry area thoroughly, remove bandage strips by peeling backing off
If necessary, cut medicated patch to fit wart
carefully apply medicated plaster, repeat procedure every 48 hours as needed (until wart is removed) for up to 12 week

WHEN USING
Ask a doctor before use if you have diabetes poor blood circulation
OTHER INFORMATION Store between 59° and 86°F (15° and 30°C).

INACTIVE INGREDIENTS
hard paraffin, lanolin, phenol.

Manufacturer:Jiangxi Yudexi Pharmaceutical Co., LTD
Address:No.1998, Jingkai West 5th Road, Chengbei Jingkai District, Zhangshu City, Yichun City, Jiangxi Province 331200, China (CN)

MINADCIN WART REMOVER PATCHES

salicylic acid 17% wart remover patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85248-201
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	17 g in 100

Inactive Ingredients

Ingredient Name	Strength
PARAFFIN (UNII: I9O0E3H2ZE)	
LANOLIN (UNII: 7EV65EAW6H)	
PHENOL (UNII: 339NCG44TV)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85248-201-01	144 in 1 BOX; Type 0: Not a Combination Product	04/30/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M028	04/30/2026	

Labeler - Jiangxi Yudexi Pharmaceutical Co., LTD (455662836)

Establishment

Name	Address	ID/FEI	Business Operations
Jiangxi Zhongletang Pharmaceuticals Co.,Ltd.		411295811	manufacture(85248-201)

Revised: 4/2026

Jiangxi Yudexi Pharmaceutical Co., LTD