### NASAL- oxymetazoline hcl spray Rite Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### **Rite Aid Corporation Nasal Spray Drug Facts**

### Active ingredient

Oxymetazoline hydrochloride 0.05%

### Purpose

Nasal decongestant

### Uses

- temporarily relieves nasal congestion due to:
- common cold
- hay fever
- upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- shrinks swollen nasal membranes so you can breathe more freely

### Warnings

### Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland

### When using this product

- do not use more than directed
- do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- use of this container by more than one person may spread infection

### Stop use and ask a doctor if

symptoms persist

### If pregnant or breast-feeding,

ask a health professional before use.

### Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: ask a doctor

To spray, squeeze bottle quickly and firmly. Do not tilt head backward while spraying. Wipe nozzle clean after use. Replace cap tightly to maintain child resistance.

### Other information

- store at 20-25°C (68-77°F)
- retain carton for future reference on full labeling

### **Inactive ingredients**

benzalkonium chloride solution, benzyl alcohol, dibasic sodium phosphate, edetate disodium, glycerin, monobasic sodium phosphate, polyethylene glycol, povidone, propylene glycol, purified water

### Questions or comments?

1-800-719-9260

### **Principal Display Panel**

SEE NEW DIRECTIONS Compare to the active ingredient of Afrin® Nasal Spray EXTRA MOISTURIZING nasal relief nasal spray oxymetazoline HCl 0.05% nasal decongestant 12 HOUR fast, powerful congestion relief soothes and helps rehydrate dry noses for colds & allergies 1 FL OZ (30 mL)



SEE NEW DIRECTIONS

\*Compare to the active ingredient of Afrin® Nasal Spray

# nasal relief

oxymetazoline HCI 0.05%

nasal decongestant



fast, powerful congestion relief

soothes and helps rehydrate dry noses

for colds & allergies




nasal relief

oxymetazoline HCI 0.05%

> nasal decongestant

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

\*This product is not manufactured or distributed by MSD Consumer Care, Inc., distributor of Afrin® Nasal Spray.



### oxymetazoline HCI 0.05%

# 



Purpose

# Drug Facts

### Active ingredient

Oxymetazoline hydrochloride 0.05%....Nasal decongestant

### Uses

- temporarily relieves nasal congestion due to:
  - common cold hay fever
  - upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- shrinks swollen nasal membranes so you can breathe more freely

### Warnings

Ask a doctor before use if you have heart disease high blood pressure

- heart disease diabetes
  - thyroid disease
- trcuble urinating due to an enlarged prostate gland

### When using this product

- do not use more than directed
- do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- use of this container by more than one person may spread infection

Stop use and ask a doctor if symptoms persist

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.

children under 6 years of age: ask a doctor To spray, squeeze bottle quickly and firmly. Do not tilt head backward while spraying. Wipe nozzle clean after use. Replace cap tightly to maintain child resistance.

## Drug Facts (continued)

Inactive ingredients


benzalkonium chloride solution, benzyl alcohol, dibasic sodium phosphate, edetate disodium, glycerin, monobasic sodium phosphate, polyethylene glycol, povidone, propylene glycol, purified water


Questions or comments? 1-800-719-9260



IF YOU'RE NOT SATISFIED, WE'LL HAPPILY REFUND YOUR MONEY.

# DISTRIBUTED BY Rite Aid 30 Hunter Lane Camp Hill, PA 17011





NASAL						
oxymetazoline hc	l spray					
Product Inform	nation					
Product Type		HUMAN OTC DRUG	Item Code (Source)		NDC:11822-0065	
Route of Adminis	tration	NASAL				
Active Ingredi	ent/Active Moi	etv				
Active Ingredient/Active Moiety Ingredient Name Basis of Stre					ngth	Strength
OXYMETAZOLINE HYDRO CHLORIDE (UNII: K89MJ0S5VY) (OXYMETAZOLINE UNII:8VLN5B44ZY)			FAZOLINE -	_		0.05 g in 100 mL
Inactive Ingree	lients					
Inactive Ingree	lients	Ingredient Name				Strength
Inactive Ingree BENZYL ALCOHO						Strength
	L (UNII: LKG8494)	NBH)				Strength
BENZYL ALCOHO	L (UNII: LKG8494 J <b>M</b> (UNII: 7FLD91C	NBH)				Strength
BENZYL ALCOHO EDETATE DISODI	L (UNII: LKG8494 J <b>M</b> (UNII: 7FLD91C DC6A3C0OX)	₩ВН) 86К)				Strength
BENZYL ALCOHO EDETATE DISODI GLYCERIN (UNII: F	L (UNII: LKG8494) J <b>M</b> (UNII: 7FLD91C DC6A3C0OX) GLYCOL (UNII: 3W	₩ВН) 86К)				Strength
BENZYL ALCOHO EDETATE DISODI GLYCERIN (UNII: F POLYETHYLENE POVIDONE (UNII: 1 PROPYLENE GLY	L (UNII: LKG8494' J <b>M</b> (UNII: 7FLD91C DC6A3C0OX) GLYCOL (UNII: 3W 7Z989GH94E) COL (UNII: 6DC9Q	WBH) 86K) /JQ0SDW1A)				Strength
BENZYL ALCOHO EDETATE DISODI GLYCERIN (UNII: F POLYETHYLENE ( POVIDONE (UNII: T PROPYLENE GLY WATER (UNII: 059)	L (UNII: LKG8494' JM (UNII: 7FLD91C DC6A3C0OX) GLYCOL (UNII: 3W 'Z989GH94E) COL (UNII: 6DC9Q QF0K00R)	WBH) 86K) 7JQ0SDW1A) 167V3)				Strength
EDETATE DISODI GLYCERIN (UNII: F POLYETHYLENE POVIDONE (UNII: 1 PROPYLENE GLY WATER (UNII: 059 SODIUM PHOSPH	L (UNII: LKG8494' JM (UNII: 7FLD91C DC6A3C0OX) GLYCOL (UNII: 3W 7Z989GH94E) COL (UNII: 6DC9Q QF0KO0R) ATE, DIBASIC (UN	WBH) 86K) /JQ0SDW1A) 167V3) II: GR686LBA74)				Strength
BENZYL ALCOHO EDETATE DISODI GLYCERIN (UNII: F POLYETHYLENE POVIDONE (UNII: 1 PROPYLENE GLY WATER (UNII: 059 SODIUM PHO SPH	L (UNII: LKG8494' JM (UNII: 7FLD91C DC6A3C0OX) GLYCOL (UNII: 3W 7Z989GH94E) COL (UNII: 6DC9Q QF0KO0R) ATE, DIBASIC (UN	WBH) 86K) 7JQ0SDW1A) 167V3)				Strength
BENZYL ALCOHO EDETATE DISODI GLYCERIN (UNII: F POLYETHYLENE POVIDONE (UNII: 1 PROPYLENE GLY WATER (UNII: 059 SODIUM PHO SPH	L (UNII: LKG8494' JM (UNII: 7FLD91C DC6A3C0OX) GLYCOL (UNII: 3W 7Z989GH94E) COL (UNII: 6DC9Q QF0KO0R) ATE, DIBASIC (UN	WBH) 86K) /JQ0SDW1A) 167V3) II: GR686LBA74)				Strength
BENZYL ALCOHO EDETATE DISODI GLYCERIN (UNII: F POLYETHYLENE POVIDONE (UNII: F PROPYLENE GLY WATER (UNII: 0594 SODIUM PHOSPH SODIUM PHOSPH	L (UNII: LKG8494' JM (UNII: 7FLD91C DC6A3C0OX) GLYCOL (UNII: 3W 7Z989GH94E) COL (UNII: 6DC9Q QF0KO0R) ATE, DIBASIC (UN	WBH) 86K) /JQ0SDW1A) 167V3) II: GR686LBA74)				Strength
BENZYL ALCOHO EDETATE DISODI GLYCERIN (UNII: F POLYETHYLENE POVIDONE (UNII: 1 PROPYLENE GLY WATER (UNII: 059 SODIUM PHO SPH	L (UNII: LKG8494' JM (UNII: 7FLD91C DC6A3C0OX) GLYCOL (UNII: 3W 'Z989GH94E) COL (UNII: 6DC9Q QF0K00R) ATE, DIBASIC (UN	WBH) 86K) /JQ0SDW1A) 167V3) II: GR686LBA74)	Man	•keting Start Date		
BENZYL ALCOHO EDETATE DISODI GLYCERIN (UNII: F POLYETHYLENE POVIDONE (UNII: T PROPYLENE GLY WATER (UNII: 059 SODIUM PHOSPH SODIUM PHOSPH	L (UNII: LKG8494' JM (UNII: 7FLD91C DC6A3C0OX) GLYCOL (UNII: 3W 7Z989GH94E) COL (UNII: 6DC9Q QF0KO0R) ATE, DIBASIC (UN ATE, MONOBASIC	WBH) 86K) 7JQ0SDW1A) 167V3) II: GR686LBA74) 2: (UNII: 3980JIH2SW)		<b>*keting Start Date</b> 5/1996		

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
OTC monograph final	part341	12/05/1996					

Labeler - Rite Aid Corporation (014578892)

Revised: 12/2018

Rite Aid Corporation