UP AND UP DAYTIME MULTI SYMPTOM SEVERE COLD NIGHTTIME SEVERE COLD AND COUGH- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, diphenhydramine hydrochloride Target Corporation

Target Corporation Daytime Multi-Symptom Severe Cold/Nighttime Severe Cold and Cough Drug Facts

Active ingredients (in each packet) - Daytime

Acetaminophen 500 mg

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer Cough suppressant Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
- minor aches and pains
- minor sore throat pain
- headache
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 6 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and	one packet
over	
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg and sodium 25 mg
- phenylketonurics: contains phenylalanine 22 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?

1-888-547-7400

Active ingredients (in each packet) Nighttime

Acetaminophen 650 mg
Diphenhydramine HCl 25 mg
Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Antihistamine/cough suppressant

Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
- minor aches and pains
- minor sore throat pain
- nasal and sinus congestion
- headache
- itchy nose or throat
- runny nose
- sneezing
- itchy, watery eyes due to hay fever
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- thyroid disease
- high blood pressure
- liver disease
- heart disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control

Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and	one packet
over	
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg and sodium 25 mg
- **phenylketonurics:** contains phenylalanine 13 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?

1-888-547-7400

Package/Label Principal Display Panel

Compare to active ingredients in Theraflu® Severe Cold

Daytime Multi-Symptom Severe Cold

Acetaminophen (Pain reliever / Fever reducer)

Dextromethorphan HBr (Cough suppressant)

Phenylephrine HCI (Nasal decongestant)

• Nasal congestion, cough, body ache, sore throat pain, headache, fever

Serve hot

Green Tea & Honey Lemon Flavors

up&up™

6 Packets

6 MULTI-SYMPTOM PACKETS

Compare to active ingredients in Theraflu ${\rm \rlap{\$}}$ Nighttime Severe Cold ${\rm \rlap{\&}}$ Cough

Nighttime Severe Cold & Cough

Acetaminophen (Pain reliever / Fever reducer)

Diphenhydramine HCI (Antihistamine / Cough suppressant)

Phenylephrine HCI (Nasal decongestant)

- Nasal congestion, cough, body ache, sore throat pain, headache, fever
- Serve hot

Honey Lemon Infused with White Tea Flavors

6 Packets

6 NIGHTTIME PACKETS



Compare to active ingredients in Theraflu® Severe Cold*

Daytime

Multi-Symptom Severe Cold

Acetaminophen (Pain reliever / Fever reducer) Dextromethorphan HBr (Cough suppressant) Phenylephrine HCI (Nasal decongestant)

- Nasal congestion, cough, body ache, sore throat pain, headache, fever
- · Serve hot





Green Tea & Honey Lemon Flavors

> 6 Packets

6 MULTI-SYMPTOM PACKETS

Compare to active ingredients in Theraflu[®] Nighttime Severe Cold & Cough*

Nighttime

Severe Cold & Cough

Acetaminophen
(Pain reliever / Fever reducer)
Diphenhydramine HCI
(Antihistamine / Cough
suppressant)
Phenylephrine HCI
(Nasal decongestant)


- Nasal congestion, cough, body ache, sore throat pain, headache, fever
- Serve hot



Honey Lemon Infused with White Tea Flavors

> 6 Packets

6 NIGHTTIME PACKETS





UP AND UP DAYTIME MULTI SYMPTOM SEVERE COLD NIGHTTIME SEVERE COLD AND COUGH

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, diphenhydramine hydrochloride kit

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:11673-877 Packaging

#	Item Code	Package Description	Date	Marketing End Date
	NDC:11673-877- 55	1 in 1 CARTON; Type 0: Not a Combination Product	07/16/2021	08/01/2026

Quant	Quantity of Parts		
Part #	Package Quantity	Total Product Quantity	
Part 1	6 PACKET	6	
Part 2	6 PACKET	6	

Part 1 of 2

UP AND UP DAYTIME MULTI SYMPTOM SEVERE COLD

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride powder, for solution

Product Information	
Item Code (Source)	NDC:82442-122
Route of Administration	ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	500 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg	

Inactive Ingredients		
Ingredient Name	Strength	
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
ASPARTAME (UNII: Z0H242BBR1)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
MALTODEXTRIN (UNII: 7CVR7L4A2D)		
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)		
SUCROSE (UNII: C151H8M554)		
TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)		

Product Characteristics

Color		Score
Shape		Size
Flavor	HONEY (green tea) , LEMON (green tea)	Imprint Code
Contains		

ı	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:82442-122-	1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information				
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	OTC Monograph Drug	M012		

Part 2 of 2

UP AND UP NIGHTTIME SEVERE COLD AND COUGH

acetaminophen, diphenhydramine hydrochloride, phenylephrine hydrochloride powder, for solution

Product Information	
Item Code (Source)	NDC:82442-964
Route of Administration	ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg	
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg	

Inactive Ingredients		
Ingredient Name	Strength	
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
ASPARTAME (UNII: Z0H242BBR1)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		

FD&C RED NO. 40 (UNII: WZB9127XOA)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SUCROSE (UNII: C151H8M554)	
TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)	

Product Characteristics			
Color	WHITE (mixture of white, light yellow-orange particles) , ORANGE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82442-964- 00	1 in 1 PACKET; Type 1: Convenience Kit of Co-Package		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012			

Marketing In	Marketing Information		
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/16/2021	08/01/2026

Labeler - Target Corporation (006961700)

Revised: 11/2025 Target Corporation