

**MULTI-SYMPTOM FLU HBP- acetaminophen, chlorpheniramine maleate,
dextromethorphan hbr tablet, film coated
CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED**

CVS 44-738 MS

Active ingredients (in each caplet)

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg

Purpose

Pain reliever/fever reducer
Antihistamine
Cough suppressant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - sore throat
 - sneezing
 - headache
 - runny nose
 - cough
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from excessive moisture
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, FD&C red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

♥CVS™

NDC 51316-738-08

actual size

**Maximum Strength†
MULTI-SYMPTOM
FLU HBP**

Acetaminophen

Pain Reliever, Fever Reducer

Chlorpheniramine Maleate

Antihistamine

Dextromethorphan HBr

Cough Suppressant

**Compare to Coricidin® HBP Maximum Strength
Multi-Symptom Flu active ingredients***

Relieves: Body aches, Pains, Headache, Fever,
Cough, Runny nose, Sore throat, Sneezing

24 CAPLETS

For people with

HIGH BLOOD PRESSURE

†Under OTC monograph.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Multi-Symptom Flu. 50844 ORG101973808

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2025 CVS/pharmacy
CVS.com® 1-800-SHOP CVS

V-19849

100% money back guaranteed.
CVS.com/returnpolicy

24 CAPLETS

MULTI-SYMP TOM FLU HBP

Maximum Strength

CVS

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

<p>Drug Facts</p> <p>Active ingredients (in each caplet)</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer</p> <p>Chlorpheniramine maleate 2 mg Antihistamine</p> <p>Dextromethorphan HBr 10 mg Cough suppressant</p> <p>Uses</p> <p>temporarily relieves these common cold and flu symptoms:</p> <ul style="list-style-type: none"> ■ minor aches and pains ■ sore throat ■ sneezing ■ headache ■ runny nose ■ temporarily reduces fever <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> ■ blisters ■ rash ■ skin reddening <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ difficulty in urination due to enlargement of the prostate gland ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin 	<p>Purpose</p> <p>When using this product</p> <ul style="list-style-type: none"> ■ marked drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ pain or cough gets worse or lasts more than 7 days ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed ■ adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours. ■ children under 12 years: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from excessive moisture ■ see end flap for expiration date and lot number <p>Inactive ingredients</p> <p>corn starch, croscopolone, FD&C red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide</p> <p>Questions or comments? 1-800-426-9391</p> <p><small>*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Multi-Symptom Flu. 50844 ORG101973808</small></p>
---	---

NDC 51316-738-08

CVS

actual size

Maximum Strength[†]

MULTI-SYMP TOM FLU HBP

Acetaminophen

Pain Reliever, Fever Reducer

Chlorpheniramine Maleate

Antihistamine

Dextromethorphan HBr

Cough Suppressant

Compare to Coricidin® HBP Maximum Strength Multi-Symptom Flu active ingredients*

Relieves: Body aches, Pains, Headache, Fever, Cough, Runny nose, Sore throat, Sneezing

24 CAPLETS | For people with HIGH BLOOD PRESSURE

†Under OTC monograph.

Distributed by CVS Pharmacy, Inc.
Woonsocket, RI 02895
© 2025 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-19849
100% money back
CVS.com/returnpolicy

#551102

FPO 80%

UPC# 050428054635

X XXXXXX XXXXXX X

8-0231-738-08-J
ORG101973808

MULTI-SYMPTOM FLU HBP

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-738
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	44;738
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-738-09	2 in 1 CARTON	06/24/2022	

1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product	
2	NDC:51316-738-08	2 in 1 CARTON	11/17/2025
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/24/2022	

Labeler - CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED (062312574)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(51316-738) , pack(51316-738)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(51316-738)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(51316-738)

Revised: 11/2025

CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED