

**ACETAMINOPHEN, ASPIRIN AND CAFFEINE- acetaminophen, aspirin and caffeine tablet**

**Rising Pharma Holdings, Inc.**

-----  
**Compare to active ingredients in EXCEDRIN® EXTRA STRENGTH\***

**Acetaminophen,  
Aspirin (NSAID\*) and  
Caffeine Tablets USP  
250 mg/250 mg/65 mg**

**Pain Reliever/ Pain Reliever Aid**

**100 Tablets**

**DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED**

**READ ALL PRODUCT INFORMATION BEFORE USING. KEEP BOX FOR  
IMPORTANT INFORMATION.**

***Drug Facts***

***Active ingredients (in each tablet)***

***Purposes***

Acetaminophen USP, 250 mg.....Pain reliever  
Aspirin USP, 250 mg (NSAID\*).....Pain reliever  
Caffeine USP, 65 mg.....Pain reliever aid

\*nonsteroidal anti-inflammatory drug

***Uses***

- temporarily relieves minor aches and pains due to:
- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual & menstrual cramps

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash. If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product.
- take more or for a longer time than directed.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for:
- diabetes
- gout
- arthritis
- taking any other drug, or are under a doctor's care for any serious condition
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer.

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away. you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- fever gets worse or lasts for more than 3 days
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs.

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not use more than directed**
- adults: take 2 tablets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 8 tablets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

### ***Other information***

- store at controlled room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using.

### ***Inactive ingredients***

Colloidal silicon dioxide , hypromellose, microcrystalline cellulose , polyethylene glycol, povidone, pregelatinized starch , sodium starch glycolate , stearic acid, talc

**Questions and comments? Call 1-844-474-7464**

**Distributed by:**

Rising Pharma Holdings, Inc.

East Brunswick, NJ 08816

Made in India

Mfg. Lic. No.: G/25/1362

**Issued:** 06/2025

S-066

\*This product is not manufactured or distributed by the owner of the registered trademark Excedrin<sup>®</sup> Extra Strength

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**Acetaminophen, Aspirin (NSAID\*) and Caffeine Tablets USP,  
250mg/250mg/65mg**

**NDC 57237-348-01**

**100 Tablets**



Inactive Ingredients	
Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

Product Characteristics			
<b>Color</b>	white (Off White)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	S7
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57237-348-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/09/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/09/2025	

**Labeler** - Rising Pharma Holdings, Inc. (116880195)

**Registrant** - Elysium Pharmaceuticals Ltd (915664486)

Establishment			
Name	Address	ID/FEI	Business Operations
Elysium Pharmaceuticals Ltd		863182240	label(57237-348) , pack(57237-348)

Establishment			
Name	Address	ID/FEI	Business Operations
Elysium Pharmaceuticals Ltd		915664486	manufacture(57237-348) , analysis(57237-348)