

**TYLAN 200- tylosin injection, solution**  
**Elanco US Inc.**

-----  
**Elanco™**

**Tylan™ 200 Injection**

**(tylosin injection) 200 mg per mL**

**For Use In Cattle and Swine Only**

**An Antibiotic**

**Indications:**

In Beef Cattle and Non-lactating Dairy Cattle, Tylan 200 Injection is indicated for use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Arcanobacterium pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Arcanobacterium pyogenes*.

In Swine, Tylan 200 Injection is indicated for use in the treatment of swine arthritis caused by *Mycoplasma hyosynoviae*; swine pneumonia caused by *Pasteurella* spp.; swine erysipelas caused by *Erysipelothrix rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.

Each mL contains 200 mg of tylosin activity (as tylosin base) in 50 percent propylene glycol with 4 percent benzyl alcohol and water for injection.

**ADMINISTRATION AND DOSAGE:**

Tylan 200 Injection is administered intramuscularly.

**BEEF CATTLE AND NON-LACTATING DAIRY CATTLE**—Inject intramuscularly 8 mg per pound of body weight one time daily (1 mL per 25 pounds). Treatment should be continued 24 hours following remission of disease signs, not to exceed 5 days. Do not inject more than 10 mL per site.

**SWINE**—Inject intramuscularly 4 mg per pound of body weight (1 mL per 50 pounds) twice daily. Treatment should be continued 24 hours following remission of disease signs, not to exceed 3 days. Do not inject more than 5 mL per site.

Read accompanying directions fully before use.

**CAUTION:**

Do not mix Tylan 200 Injection with other injectable solutions as this may cause a precipitation of the active ingredients.

**WARNINGS:**

NOT FOR HUMAN USE.

KEEP OUT OF REACH OF CHILDREN.

**Adverse reactions, including shock and death may result from overdose in baby pigs.**

Do not attempt injection into pigs weighing less than 25 pounds (0.5 mL) with the common syringe. It is recommended that Tylan 50 Injection be used in pigs weighing less than 25 pounds.

Do not administer to horses or other equines. Injection of tylosin in equines has been fatal.

**RESIDUE WARNING: Swine:**

Swine intended for human consumption must not be slaughtered within 14 days of the last use of this drug product.

**RESIDUE WARNING: Cattle:**

Cattle intended for human consumption must not be slaughtered within 21 days of the last use of this drug product. This drug product is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. This product is not approved for use in calves intended to be processed for veal. A withdrawal period has not been established in pre-ruminating calves.

If tylosin medicated drinking water is used as a follow-up treatment for swine dysentery, the animal should thereafter receive feed containing 40 to 100 grams of tylosin per ton for 2 weeks to assure depletion of tissue residues.

**STORAGE CONDITIONS:**

Store at or below 25°C (77°F). Use this product within 90 days of the first puncture. Puncture the 250 mL vial a maximum of 35 times. If more than 35 punctures are anticipated, the use of multi-dosing equipment is recommended. When using a draw-off spike or needle with bore diameter larger than 16 gauge, discard any product remaining in the vial immediately after use.

Tylan, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2025 Elanco or its affiliates

**Approved by FDA under NADA # 012-965**

For additional information about reporting adverse drug experiences for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

**Manufactured for:**

**Elanco US Inc.**

**Greenfield, IN 46140, USA**

Product of Ireland

AH0206

YL231482X

Product of Brazil  
AH0206  
YL301935C

**Elanco™**

**Tylan™ 200 Inyectable**

**(tilosina inyectable) 200 mg por ml**

**Para uso exclusivo en ganado vacuno y cerdos**

**Un antibiótico**

### **Indicaciones:**

En ganado vacuno y vacas lecheras no lactantes, Tylan 200 Inyectable se indica para el tratamiento del complejo respiratorio bovino (fiebre de embarque, neumonía), generalmente asociado con *Pasteurella multocida* y *Arcanobacterium pyogenes*, pietín (pododermatitis necrótica), difteria de los terneros provocada por *Fusobacterium necrophorum* y metritis provocada por *Arcanobacterium pyogenes*.

En cerdos, Tylan 200 Inyectable se indica para el tratamiento de artritis en cerdos provocada por *Mycoplasma hyosynoviae*, neumonía porcina causada por *Pasteurella* spp., erisipelas porcinas provocadas por *Erysipelothrix rhusiopathiae*, disentería porcina asociada con *Treponema hyodysenteriae* cuando es tratada con el medicamento apropiado a través del alimento y/o el agua para beber.

Cada ml contiene 200 mg de actividad de tilosina (como tilosina base) en propilenglicol al 50 por ciento, alcohol bencílico al 4 por ciento y agua para inyección.

### **POSOLOGÍA Y ADMINISTRACIÓN:**

Tylan 200 Inyectable se administra por vía intramuscular.

**GANADO VACUNO Y VACAS LECHERAS NO LACTANTES**—Inyectar por vía intramuscular 8 mg por libra de peso corporal una vez al día (1 ml cada 25 libras). El tratamiento debe continuarse durante 24 horas luego de la remisión de los signos de la enfermedad sin extenderse más de 5 días. No aplicar más de 10 ml por lugar de inyección.

**CERDOS**—Inyectar por vía intramuscular 4 mg por libra de peso corporal (1 ml cada 50 libras) dos veces al día. El tratamiento debe continuarse durante 24 horas luego de la remisión de los signos de la enfermedad sin extenderse más de 3 días. No aplicar más de 5 ml por lugar de inyección.

Leer todas las instrucciones adjuntas antes de usar.

### **PRECAUCIÓN:**

No mezclar la inyección Tylan 200 con otras soluciones inyectables ya que esto puede ocasionar la precipitación de los principios activos.

## **ADVERTENCIAS:**

ESTE PRODUCTO NO DEBE UTILIZARSE EN SERES HUMANOS.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.

**Pueden ocurrir reacciones adversas, incluidos shock y muerte, en caso de sobredosis en crías de cerdos.**

No administrar la inyección a cerdos que pesen menos de 25 libras (0.5 ml) con la jeringa común. Se recomienda usar la inyección Tylan 50 en cerdos que pesen menos de 25 libras.

No administrar a caballos u otros equinos. La inyección de tilosina en equinos ha resultado mortal.

## **ADVERTENCIA ACERCA DE RESIDUOS:**

**Ganado porcino:** el ganado porcino previsto para consumo humano no se debe faenar durante los 14 días posteriores al último uso de este producto farmacológico.

## **ADVERTENCIA ACERCA DE RESIDUOS:**

**Ganado bovino:** el ganado bovino previsto para consumo humano no se debe faenar durante los 21 días posteriores al último uso de este producto farmacológico. Este producto farmacológico no está aprobado para su uso en ganado bovino lechero hembra de 20 meses de edad o más, incluidas las vacas lecheras secas. El uso en este ganado bovino puede producir residuos farmacológicos en la leche y/o en los terneros nacidos de estas vacas. Este producto no está aprobado para el uso en terneros que se procesarán para carne de ternera. No se ha establecido un período de retiro del fármaco en terneros prerrumiantes.

Si se suministra agua para beber con tilosina como tratamiento de seguimiento para la disentería porcina, el animal debe recibir posteriormente alimento que contenga entre 40 y 100 gramos de tilosina por tonelada durante 2 semanas para garantizar la depleción de los residuos de tejidos.

## **CONDICIONES DE ALMACENAMIENTO:**

Almacenar a una temperatura de 25 °C (77 °F) o inferior. Utilice este producto dentro de los 90 días posteriores a la primera punción. Perfore el vial de 250 ml un máximo de 35 veces. Si se anticipan más de 35 punciones, se recomienda el uso de equipo de dosificación. Cuando utilice una punta o aguja desprendible con un diámetro interior mayor que el calibre 16, deseche cualquier producto restante en el vial inmediatamente después de usarlo.

Tylan, Elanco y el logo de la barra diagonal son marcas de Elanco o sus afiliadas.

**Aprobado por la FDA bajo NADA # 012-965**

Para obtener información adicional sobre cómo informar experiencias adversas de medicamentos para medicamentos animales, comuníquese con la FDA al 1-888-FDA-VETS o <http://www.fda.gov/reportanimalae>.

**Fabricado por: Elanco US Inc.  
Greenfield, IN 46140, USA**

Producto de Irlanda  
AH0206  
YL231483X

Producto de Brasil  
AH0206  
YL301935C

## **Principal Display Panel - Tylan 200 Injection 250 mL Carton Label**

**Elanco™**

AH0206

**Tylan™ 200  
Injection**

™

**200 mg per mL**

**(tylosin injection)**

**For Use In Cattle and Swine Only**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**The Original Tylan 200 Injection**

**An Antibiotic  
For Cattle Diseases**

- **Pneumonia**
- **Shipping Fever**
- **Metritis**
- **Diphtheria**
- **Foot Rot**

**For Swine Diseases**

- **Pneumonia**
- **Mycoplasmal Arthritis**
- **Dysentery**
- **Erysipelas**

**250 mL**



SH-23 1422X

Overprint area  
35 mm x 25 mm  
Center line of overprint



Exp. Date  
Lot  
**Tylan 200**  
Injection  
200 mg per mL  
**Elanco**  
AH0206

NOTE: Text size 8 pt  
on 1.5 pt to allow 4 mm  
spacing between lines to  
match up with overprint.

Shaded Areas  
= No text

**Elanco** AH0206  
**Tylan 200**  
Injection  
200 mg per mL  
**(tylosin injection)**

Manufactured for:  
Elanco US Inc.  
Greenfield, IN 46140, USA  
  
For additional information about reporting  
adverse drug experiences for animal drugs,  
contact FDA at 1-888-FDA-VETS or  
<http://www.fda.gov/reportanimalae>.

**CAUTION:**  
Do not mix Tylan 200 Injection with other  
injectable solutions as this may cause a  
precipitation of the active ingredients.  
**WARNINGS:**  
NOT FOR HUMAN USE.  
KEEP OUT OF REACH OF CHILDREN.  
Adverse reactions, including shock and  
death may result from overdosage in  
baby pigs.  
Do not attempt injection into pigs weighing  
less than 25 pounds (0.5 mL) with the  
common syring. It is recommended that  
Tylan 50 Injection be used in pigs weighing  
less than 25 pounds.  
Do not administer to horses or other equines.  
Injection of tylosin in equines has been fatal.

**RESIDUE WARNING: Swine:**  
Swine intended for human  
consumption must not be  
slaughtered within 14 days of the  
last use of this drug product.  
**RESIDUE WARNING: Cattle:**  
Cattle intended for human  
consumption must not be  
slaughtered within 21 days of  
the last use of this drug product.  
This drug product is not approved  
for use in female dairy cattle  
20 months of age or older,  
including dry dairy cows. Use  
in these cattle may cause drug  
residue in milk and/or in calves  
born to these cows. This product  
is not approved for use in calves  
intended to be processed for veal.  
A withdrawal period has not been  
established in pre-ruminating  
calves.

If tylosin medicated drinking water is used as  
a follow-up treatment for swine dysentery,  
the animal should thereafter receive feed  
containing 40 to 100 grams of tylosin per  
ton for 2 weeks to assure depletion of  
tissue residue.

**Elanco** AH0206  
**Tylan 200**  
Injection  
200 mg per mL  
**(tylosin injection)**

For Use In Cattle and Swine Only

**CAUTION:** Federal law restricts  
this drug to use by or on the order  
of a licensed veterinarian.

The Original Tylan 200 Injection

- An Antibiotic  
For Cattle Diseases
- Pneumonia
  - Shipping Fever
  - Metritis
  - Diphtheria
  - Foot Rot

- For Swine Diseases
- Pneumonia
  - Mycoplasmal Arthritis
  - Dysentery
  - Erysipelas

250 mL

**Dosage and Administration:**  
**Beef Cattle and Non-lactating Dairy Cattle—**  
Inject intramuscularly 8 mg per pound of body  
weight one time daily (1 mL per 25 pounds).  
Treatment should be continued 24 hours  
following remission of disease signs, not to  
exceed 5 days.  
Do not inject more than 10 mL per site.  
**Swine—**  
Inject intramuscularly 4 mg per pound of  
body weight (1 mL per 50 pounds) twice daily.  
Treatment should be continued 24 hours  
following remission of disease signs, not to  
exceed 3 days.  
Do not inject more than 5 mL per site.

**In Beef Cattle and  
Non-lactating Dairy Cattle:**  
Treatment of bovine respiratory complex  
(shipping fever, pneumonia), foot rot (necrotic  
pododermatitis), calf diphtheria and metritis.

**In Swine:**  
Treatment of mycoplasmal swine arthritis,  
swine pneumonia, swine erysipelas and swine  
dysentery when followed by appropriate  
medication in the drinking water and/or feed.

Each mL contains 200 mg of tylosin activity  
(as tylosin base) in 50 percent propylene  
glycol with 4 percent benzyl alcohol and water  
for injection.

Read enclosed directions fully before use.  
Store at or below 25°C (77°F)  
Product of Ireland

Approved by FDA under NADA # 012-965





## Principal Display Panel - Tylan 200 Injection 250 mL Bottle Label

**Elanco™**

AH0206

**Tylan™ 200**

# Injection

TM

250 mL

(tylosin injection)

For Use In Cattle and Swine Only

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian

200 mg per mL

An Antibiotic

**Elanco™** AH0206

**Tylan™ 200**  
Injection

250 mL

**(tylosin injection)**

For Use In Cattle and Swine Only

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

200 mg per mL

An Antibiotic

Indications: In Beef Cattle and Non-lactating Dairy Cattle,

Tylan 200 Injection is indicated for use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Arcanobacterium pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Arcanobacterium pyogenes*.

In Swine, Tylan 200 Injection is indicated for use in the treatment of swine arthritis caused by *Mycoplasma hyosynoviae*; swine pneumonia caused by *Pasteurella* spp.; swine erysipelas caused by *Erysipelothrix rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.

Each mL contains 200 mg of tylosin activity (as tylosin base) in 50 percent propylene glycol with 4 percent benzyl alcohol and water for injection.

**ADMINISTRATION AND DOSAGE:**  
Tylan 200 Injection is administered intramuscularly.

**Manufactured for:**  
**Elanco US Inc.**  
**Greenfield, IN 46140, USA**  
Product of Ireland

YL231482X

7 27804 20114 2

**Elanco™** AH0206

**Tylan™ 200**  
Injection

250 mL

**(tylosin injection)**

For Use In Cattle and Swine Only

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

200 mg per mL

An Antibiotic

Indications: In Beef Cattle and Non-lactating Dairy Cattle,

Tylan 200 Injection is indicated for use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Arcanobacterium pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Arcanobacterium pyogenes*.

In Swine, Tylan 200 Injection is indicated for use in the treatment of swine arthritis caused by *Mycoplasma hyosynoviae*; swine pneumonia caused by *Pasteurella* spp.; swine erysipelas caused by *Erysipelothrix rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.

Each mL contains 200 mg of tylosin activity (as tylosin base) in 50 percent propylene glycol with 4 percent benzyl alcohol and water for injection.

**ADMINISTRATION AND DOSAGE:**  
Tylan 200 Injection is administered intramuscularly.

**Manufactured for:**  
**Elanco US Inc.**  
**Greenfield, IN 46140, USA**  
Product of Brazil

YL301935C

4023362

Lot: / Lote:  
Exp. Date: /  
Fecha de vencimiento:

Principal Display Panel - Tylan 200 Injection 250 mL Spanish Bottle Label

Elanco™

AH0206

## Tylan™ 200 Inyectable

™

250 mL

(tilosina inyectable)

Para uso exclusivo en ganado  
vacuno y cerdos

**PRECAUCIÓN:** Las leyes federales establecen que el uso de este fármaco se restrinja a veterinarios con licencia o bajo la indicación de estos.

200 mg por mL

Un antibiótico

Elanco™ AH0206  
**Tylan™ 200**  
**Inyectable**™  
250 mL

(tilosina inyectable)

Para uso exclusivo en ganado  
vacuno y cerdos

**PRECAUCIÓN:** Las leyes federales establecen que el uso de este fármaco se restrinja a veterinarios con licencia o bajo la indicación de estos.

200 mg por ml

Un antibiótico

Indicaciones: En ganado vacuno

y vacas lecheras no lactantes, Tylan 200 Inyectable se indica para el tratamiento del complejo respiratorio bovino (fiebre de embarque, neumonía), generalmente asociado con *Pasteurella multocida* y *Arcanobacterium pyogenes*, pietin (pododermatitis necrótica), difteria de los terneros provocada por *Fusobacterium necrophorum* y metritis provocada por *Arcanobacterium pyogenes*.

En cerdos, Tylan 200 Inyectable se indica para el tratamiento de artritis en cerdos provocada por *Mycoplasma hyosynoviae*, neumonía porcina causada por *Pasteurella* spp., erisipelas porcinas provocadas por *Erysipelothrix rhusiopathiae*, disentería porcina asociada con *Treponema hyodysenteriae* cuando

es tratada con el medicamento apropiado a través del alimento y/o el agua para beber.

Cada ml contiene 200 mg de actividad de tilosina (como tilosina base) en propilenglicol al 50 por ciento, alcohol bencílico al 4 por ciento y agua para inyección.

**POSOLOGÍA Y ADMINISTRACIÓN:**

Tylan 200 inyectable se administra por vía intramuscular.

**Fabricado por:**

**Elanco US Inc.**  
**Greenfield, IN 46140, USA**

Producto de Irlanda

YL231482X

<p><b>Elanco™</b> AH0206</p> <p><b>Tylan™ 200</b> <b>Inyectable</b> <sup>TM</sup></p> <p><b>250 mL</b></p>	<p>y vacas lecheras no lactantes, Tylan 200 Inyectable se indica para el tratamiento del complejo respiratorio bovino (fiebre de embarque, neumonía), generalmente asociado con <i>Pasteurella multocoda</i> y <i>Arcanobacterium pyogenes</i>, piétin (pododermatitis necrótica), difteria de los terneros provocada por <i>Fusobacterium necrophorum</i> y metritis provocada por <i>Arcanobacterium pyogenes</i>.</p> <p>En cerdos, Tylan 200 Inyectable se indica para el tratamiento de artritis en cerdos provocada por <i>Mycoplasma hyosynoviae</i>, neumonía porcina causada por <i>Pasteurella</i> spp., erisipelas porcinas provocadas por <i>Erysipelothrix rhusiopathiae</i>, disentería porcina asociada con <i>Treponema hyodysenteriae</i> cuando es tratada con el medicamento apropiado a través del alimento y/o el agua para beber.</p>	<p>Cada ml contiene 200 mg de actividad de tilosina (como tilosina base) en propilenglicol al 50 por ciento, alcohol bencílico al 4 por ciento y agua para inyección.</p> <p><b>POSOLOGÍA Y ADMINISTRACIÓN:</b> Tylan 200 inyectable se administra por vía intramuscular.</p> <p><b>Fabricado por:</b> <b>Elanco US Inc.</b> <b>Greenfield, IN 46140, USA</b> Producto de Brasil</p>
<p><b>(tilosina inyectable)</b> Para uso exclusivo en ganado vacuno y cerdos</p> <p><b>PRECAUCIÓN:</b> Las leyes federales establecen que el uso de este fármaco se restrinja a veterinarios con licencia o bajo la indicación de estos.</p> <p><b>200 mg por ml</b> <b>Un antibiótico</b> Indicaciones: En ganado vacuno</p>		

YL301935C

## Principal Display Panel - Tylan 200 Injection 500 mL Bottle Label

**Elanco™**

AH0206

**Tylan™ 200**  
**Injection**

<sup>TM</sup>

**Professional Size**  
**500 mL**

**(tylosin injection)**

**For Use In Cattle and Swine Only**

**200 mg per mL**


**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**An Antibiotic**


**Elanco™** AH0206  
**Tylan™ 200**  
**Injection**™  
 Professional Size  
**500 mL**  
**(tylosin injection)**  
 For Use In Cattle and Swine Only  
 200 mg per mL  
 CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.  
 An Antibiotic  
 Use automatic syringe equipment only  
 Indications: In Beef Cattle and Non-lactating Dairy Cattle,

Tylan 200 Injection is indicated for use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Arcanobacterium pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Arcanobacterium pyogenes*.  
 In Swine, Tylan 200 Injection is indicated for use in the treatment of swine arthritis caused by *Mycoplasma hyosynoviae*; swine pneumonia caused by *Pasteurella* spp.; swine erysipelas caused by *Erysipelothrix rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.  
 Each mL contains 200 mg of tylosin activity (as tylosin base) in 50 percent propylene glycol with 4 percent benzyl alcohol and water for injection.

**ADMINISTRATION AND DOSAGE:**  
 Tylan 200 Injection is administered intramuscularly.  
**Manufactured for:**  
 Elanco US Inc.  
 Greenfield, IN 46140, USA  
 Product of Ireland

YL231483X  
  
 7 27804 20115 9  


**Elanco™** AH0206  
**Tylan™ 200**  
**Injection**™  
 Professional Size  
**500 mL**  
**(tylosin injection)**  
 For Use In Cattle and Swine Only  
 200 mg per mL  
 CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.  
 An Antibiotic

Indications: In Beef Cattle and Non-lactating Dairy Cattle, Tylan 200 Injection is indicated for use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Arcanobacterium pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Arcanobacterium pyogenes*.  
 In Swine, Tylan 200 Injection is indicated for use in the treatment of swine arthritis caused by *Mycoplasma hyosynoviae*; swine pneumonia caused by *Pasteurella* spp.; swine erysipelas caused by *Erysipelothrix rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.  
 Each mL contains 200 mg of tylosin activity (as tylosin base) in 50 percent propylene glycol with 4 percent benzyl alcohol and water for injection.

**ADMINISTRATION AND DOSAGE:**  
 Tylan 200 Injection is administered intramuscularly.  
**Manufactured for:**  
 Elanco US Inc.  
 Greenfield, IN 46140, USA  
 Product of Brazil

YL301934C  
  
 7 27804 56445 2  
 4021346  
 Lot: / Lote:  
 Exp. Date: /  
 Fecha de vencimiento:

## Principal Display Panel - Tylan 200 Injection 500 mL Spanish Bottle Label

**Elanco™**

AH0206

**Tylan™ 200**  
**Inyectable**

™

**Tamaño profesional**  
**500 ml**

**(tilosina inyectable)**

**Para uso exclusivo en ganado**  
**vacuno y cerdos**

**200 mg por ml**

**PRECAUCIÓN:** Las leyes federales

establecen que el uso de este fármaco se restrinja a veterinarios con licencia o bajo la indicación de estos.

## Un antibiótico

<p><b>Elanco™</b> AH0206</p> <p><b>Tylan™ 200</b> <b>Inyectable</b>™</p> <p>Tamaño profesional <b>500 ml</b></p> <p><b>(tilosina inyectable)</b></p> <p>Para uso exclusivo en ganado vacuno y cerdos 200 mg por ml</p> <p><b>PRECAUCIÓN:</b> Las leyes federales establecen que el uso de este fármaco se restrinja a veterinarios con licencia o bajo la indicación de estos.</p> <p>Un antibiótico</p>	<p>Utilizar sólo equipos automáticos de jeringas</p> <p>Indicaciones: En ganado vacuno y vacas lecheras no lactantes, Tylan 200 Inyectable se indica para el tratamiento del complejo respiratorio bovino (fiebre de embarque, neumonía), generalmente asociado con <i>Pasteurella multocida</i> y <i>Arcanobacterium pyogenes</i>, piétin (pododermatitis necrótica), difteria de los terneros provocada por <i>Fusobacterium necrophorum</i> y metritis provocada por <i>Arcanobacterium pyogenes</i>.</p> <p>En cerdos, Tylan 200 Inyectable se indica para el tratamiento de artritis en cerdos provocada por <i>Mycoplasma hyosynoviae</i>, neumonía porcina causada por <i>Pasteurella</i> spp., erisipelas porcinas provocadas por <i>Erysipelothrix rhusiopathiae</i>, disentería porcina</p>	<p>asociada con <i>Treponema hyodysenteriae</i> cuando es tratada con el medicamento apropiado a través del alimento y/o el agua para beber.</p> <p>Cada ml contiene 200 mg de actividad de tilosina (como tilosina base) en propilenglicol al 50 por ciento, alcohol bencílico al 4 por ciento y agua para inyección.</p> <p><b>POSOLOGÍA Y ADMINISTRACIÓN:</b> Tylan 200 Inyectable se administra por vía intramuscular.</p> <p><b>Fabricado por:</b> <b>Elanco US Inc.</b> Greenfield, IN 46140, USA</p> <p>Producto de Irlanda</p> <p>YL231483X</p>
--	---	---

<p><b>Elanco™</b> AH0206</p> <p><b>Tylan™ 200</b> <b>Inyectable</b>™</p> <p>Tamaño profesional <b>500 ml</b></p> <p><b>(tilosina inyectable)</b></p> <p>Para uso exclusivo en ganado vacuno y cerdos 200 mg por ml</p> <p><b>PRECAUCIÓN:</b> Las leyes federales establecen que el uso de este fármaco se restrinja a veterinarios con licencia o bajo la indicación de estos.</p> <p>Un antibiótico</p>	<p>Indicaciones: En ganado vacuno y vacas lecheras no lactantes, Tylan 200 Inyectable se indica para el tratamiento del complejo respiratorio bovino (fiebre de embarque, neumonía), generalmente asociado con <i>Pasteurella multocida</i> y <i>Arcanobacterium pyogenes</i>, piétin (pododermatitis necrótica), difteria de los terneros provocada por <i>Fusobacterium necrophorum</i> y metritis provocada por <i>Arcanobacterium pyogenes</i>.</p> <p>En cerdos, Tylan 200 Inyectable se indica para el tratamiento de artritis en cerdos provocada por <i>Mycoplasma hyosynoviae</i>, neumonía porcina causada por <i>Pasteurella</i> spp., erisipelas porcinas provocadas por <i>Erysipelothrix rhusiopathiae</i>, disentería porcina asociada con <i>Treponema hyodysenteriae</i> cuando es tratada con el medicamento</p>	<p>apropiado a través del alimento y/o el agua para beber.</p> <p>Cada ml contiene 200 mg de actividad de tilosina (como tilosina base) en propilenglicol al 50 por ciento, alcohol bencílico al 4 por ciento y agua para inyección.</p> <p><b>POSOLOGÍA Y ADMINISTRACIÓN:</b> Tylan 200 Inyectable se administra por vía intramuscular.</p> <p><b>Fabricado por:</b> <b>Elanco US Inc.</b> Greenfield, IN 46140, USA</p> <p>Producto de Brasil</p> <p>YL301934C</p>
--	--	--

## TYLAN 200

tylosin injection, solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:58198-0206
<b>Route of Administration</b>	INTRAMUSCULAR		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
tylosin (UNII: YEF4JXN031) (tylosin - UNII:YEF4JXN031)	tylosin	200 mg in 1 mL

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:58198-0206-3	1 in 1 CARTON		
1		250 mL in 1 BOTTLE, GLASS		
2	NDC:58198-0206-5	500 mL in 1 BOTTLE, GLASS		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NADA	NADA012965	07/08/2013	

**Labeler** - Elanco US Inc. (966985624)

Revised: 7/2025

Elanco US Inc.