

**ANTACID ANTIGAS- aluminum hydroxide, magnesium hydroxide, dimethicone suspension**  
**Chain Drug Consortium, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active Ingredients**

Aluminum hydroxide (equivalent to Dried Gel, USP) 200 mg  
Magnesium hydroxide 200 mg  
Simethicone 20 mg

**Uses**

relieves: • heartburn • sour stomach  
• acid indigestion • upset stomach due to these symptoms  
• symptoms of gas

**Warnings**

**Ask a doctor before use if you have**

• kidney disease • a magnesium restricted diet

**Ask a doctor or pharmacist before use if you are**

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if**

symptoms last for more than 2 weeks

**If pregnant or breast-feeding**

ask a health professional before use.

**Keep this and all drugs out of the reach of children.**

In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

**Directions**

• shake well before use  
• do not exceed 24 teaspoons (120 mL) in a 24 hour period or use the maximum dosage for more than 2 weeks  
• adults and children 12 years of age and older: take 2 to 4 teaspoons (10-20 mL) between meals and at bedtime or as directed by a doctor

- children under 12 years of age: ask a doctor

**Other Information**

- do not freeze
- store at room temperature tightly closed
- each teaspoon contains:  
potassium 2 mg, magnesium 83 mg

**Inactive Ingredients**

benzyl alcohol, flavors, glycerin, hydroxyethyl cellulose,  
potassium citrate, propylene glycol, purified water, sorbitol

**Principal Display Panel**

This product is not manufactured or distributed by Johnson and Johnson Merck, distributor of Mylanta

## Drug Facts

<b>Active ingredients</b> (in each 5 mL = 1 teaspoon)	<b>Purpose</b>
Aluminum hydroxide (equivalent to Dried Gel, USP) 200 mg.....	Antacid
Magnesium hydroxide 200 mg.....	Antacid
Simethicone 20 mg.....	Antigas

**Uses** relieves: • heartburn • sour stomach  
• acid indigestion • upset stomach due to these symptoms  
• symptoms of gas

### Warnings

**Ask a doctor before use if you have**

• kidney disease • a magnesium restricted diet

**Ask a doctor or pharmacist before use if you are** presently taking a prescription drug. Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if** symptoms last for more than 2 weeks

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep this and all drugs out of the reach of children.**

In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

### Directions

• shake well before use  
• do not exceed 24 teaspoons (120 mL) in a 24 hour period or use the maximum dosage for more than 2 weeks  
• adults & children 12 years of age and older: take 2 to 4 teaspoons (10-20 mL) between meals and at bed time or as directed by a doctor  
• children under 12 years of age: ask a doctor

### Other information

• do not freeze  
• store at room temperature tightly closed  
• **each teaspoon contains:**  
potassium 2 mg, magnesium 83 mg

**Inactive ingredients** benzyl alcohol, flavors, glycerin, hydroxyethyl cellulose, potassium citrate, propylene glycol, purified water, sorbitol

**TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Johnson & Johnson Merck, distributor of Mylanta®

Distributed by: Chain Drug Consortium, LLC.,  
2300 NW Corporate Blvd., Suite 115  
Boca Raton, FL 33431

Questions or Comments? 1-866-534-4631

68-01372B



## Purposes

Antacid

Antacid

Antigas

## ANTACID ANTIGAS

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-004
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ALUMINUM HYDRO XIDE</b> (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDRO XIDE	200 mg in 5 mL
<b>MAGNESIUM HYDRO XIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDRO XIDE	200 mg in 5 mL
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	20 mg in 5 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYDROXYETHYL CELLULOSE (2000 MPAS AT 1%)</b> (UNII: S38J6RZN16)	
<b>POTASSIUM CITRATE</b> (UNII: EE90ONI6FF)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	LEMON (Maalox Type Lemon Flavor)	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-004-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part331	06/03/2009	

**Labeler** - Chain Drug Consortium, LLC (101668460)

