

TYLENOL REGULAR STRENGTH- acetaminophen tablet, film coated
Kenvue Brands LLC

TYLENOL Regular Strength

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg) in 24 hours for adults or 5 caplets (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 4 to 6 hours while symptoms last▪ do not take more than 10 caplets in 24 hours, unless directed by a doctor▪ do not use for more than 10 days unless directed by a doctor
children 6 years to under 12 years	<ul style="list-style-type: none">▪ take 1 caplet every 4 to 6 hours while symptoms last▪ do not take more than 5 caplets in 24 hours▪ do not use for more than 5 days unless directed by a doctor
children under 6 years	ask a doctor

Other information

- store between 20-25°C (68-77°F). Avoid high humidity. Protect from light.
- **do not use if any individual blister unit is open or torn**

Inactive ingredients

carnauba wax, crospovidone, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

325mg

NDC 50580-600-02

To re-order reference the NDC code

TYLENOL®

Acetaminophen Pain Reliever - Fever Reducer

Regular Strength

EZ DOSE™

100 Caplets

325 mg each

10 Blister Cards

with

10 Individual Blisters


SMART PACKAGING

DESIGNED FOR HEALTH

CARE PROFESSIONALS

FOR HOSPITAL AND GOVERNMENT USE ONLY

313493



Regular Strength

Acetaminophen Pain Reliever - Fever Reducer

TYLENOL

325mg

NDC 50580-600-02

To re-order reference the NDC code

TYLENOL

Acetaminophen Pain Reliever - Fever Reducer

Regular Strength

TYLENOL 325
HOSPITAL

100 Caplets
325 mg each

10 Blister Cards
with
10 Individual Blisters

EZ DOSE™

SMART PACKAGING
DESIGNED FOR HEALTH
CARE PROFESSIONALS

FOR HOSPITAL AND GOVERNMENT USE ONLY



NDC 50580-600-02

325mg


EZ DOSE™

Blister pack
configuration for
ease of inventory

Clear dosing
information for
ease of identification

Perforation and
rounded corners for
easy dispensing

Scannable LOT/EXP
information on
bottom of carton



30035500/313493

313493

325mg

TYLENOL®

Acetaminophen Pain Reliever - Fever Reducer

Regular Strength

Drug Facts

Active ingredient (in each caplet) **Purpose**
Acetaminophen 325 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg) in 24 hours for adults or 5 caplets (1,625 mg) in 24 hours for children. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Drug Facts (continued)

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
 - pain gets worse or lasts more than 5 days in children under 12 years
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)

adults and
children
12 years
and over

- take 2 caplets every 4 to 6 hours while symptoms last
- do not take more than 10 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children 6
years to under
12 years

- take 1 caplet every 4 to 6 hours while symptoms last
- do not take more than 5 caplets in 24 hours
- do not use for more than 5 days unless directed by a doctor

children
under 6 years

ask a doctor

Other information

- store between 20-25°C (68-77°F). Avoid high humidity. Protect from light.
- do not use if any individual blister unit is open or torn

Inactive ingredients camellia wax, croscollidone, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

©J&J 2016

Visit us at www.tylenol.com or call toll-free 1-877-TYLENOL (1-877-895-3665)



Care To Recycle™

TYLENOL REGULAR STRENGTH

acetaminophen tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-600
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name		Basis of Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		325 mg

Inactive Ingredients	
Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SHELLAC (UNII: 46N107B71O)	

Product Characteristics			
Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	TYLENOL;325;HOSPITAL
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-600-01	50 in 1 BOTTLE; Type 0: Not a Combination Product	08/31/2017	

2	NDC:50580-600-02	10 in 1 CARTON	05/31/2017	10/22/2021
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:50580-600-03	1000 in 1 BOTTLE; Type 0: Not a Combination Product	08/31/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/31/2017	

Labeler - Kenvue Brands LLC (118772437)

Revised: 11/2024

Kenvue Brands LLC