

# LIPIODOL- ethiodized oil injection

## Guerbet LLC

-----

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LIPIODOL safely and effectively. See full prescribing information for LIPIODOL.

### LIPIODOL® (Ethiodized Oil) Injection

Initial U.S. Approval: 1954

**WARNING: FOR INTRALYMPHATIC, INTRAUTERINE AND SELECTIVE HEPATIC INTRA-ARTERIAL USE ONLY**

*See Full Prescribing Information for complete Boxed Warning*

**Pulmonary and cerebral embolism can result from inadvertent intravascular injection or intravasation of Lipiodol. Inject Lipiodol slowly with radiologic monitoring; do not exceed recommended dose (5.1).**

### RECENT MAJOR CHANGES

Indications and Usage (1) 4/2014

Dosage and Administration, Dosage Guidelines (2.1) 4/2014

Contraindications (4) 4/2014

Warnings and Precautions (5) 4/2014

### INDICATIONS AND USAGE

Lipiodol is an oil-based radiopaque contrast agent indicated for: (1)

- hysterosalpingography in adults
- lymphography in adult and pediatric patients
- selective hepatic intra-arterial use for imaging tumors in adults with known hepatocellular carcinoma (HCC)

### DOSAGE AND ADMINISTRATION

Use a disposable syringe to draw and inject Lipiodol. (2)

- **Hysterosalpingography**

Inject increments of 2 mL of Lipiodol into the endometrial cavity until tubal patency is determined; stop the injection if the patient develops excessive discomfort. Inject with radiologic monitoring.

- **Lymphography**

Inject Lipiodol into a lymphatic vessel with radiologic monitoring.

Adults:

- unilateral lymphography of the upper extremities: 2 to 4 mL
- unilateral lymphography of the lower extremities: 6 to 8 mL
- penile lymphography: 2 to 3 mL
- cervical lymphography: 1 to 2 mL

Pediatric patients:

- Inject a minimum of 1 mL to a maximum of 6 mL according to the anatomical area to be visualized. Do not exceed 0.25 mL/kg.
- **Selective Hepatic Intra-arterial Use**  
Inject 1.5 to 15 mL of Lipiodol slowly under continuous radiologic monitoring. Do not exceed 20 mL total dosage.

### DOSAGE FORMS AND STRENGTHS

Each mL of Lipiodol contains 480 mg Iodine organically combined with ethyl esters of fatty acids of poppy seed oil. (3)

### CONTRAINDICATIONS

Hypersensitivity to Lipiodol, hyperthyroidism, traumatic injuries, recent hemorrhage or bleeding. (4)

- **Lipiodol Hysterosalpingography is contraindicated in:**

pregnancy, acute pelvic inflammatory disease, marked cervical erosion, endocervicitis and intrauterine bleeding, in the immediate pre-or postmenstrual phase, or within 30 days of curettage or conization.

- **Lipiodol Lymphography is contraindicated in:**

right to left cardiac shunt, advanced pulmonary disease, tissue trauma or hemorrhage, advanced neoplastic disease with expected lymphatic obstruction, previous surgery interrupting the lymphatic system, or radiation therapy to the examined area.

- **Lipiodol Selective Hepatic Intra-arterial Injection is contraindicated in:**

the presence of dilated bile ducts unless external biliary drainage was performed before injection.

----- **WARNINGS AND PRECAUTIONS** -----

- Pulmonary and cerebral embolism: avoid use in patients with severely impaired lung function, cardiorespiratory failure or right-sided cardiac overload (5.1)
- Hypersensitivity reactions: avoid use in patients with a history of sensitivity to other iodinated contrast agents, bronchial asthma or allergic disorders because of an increased risk of a hypersensitivity reaction to Lipiodol (5.2)
- Exacerbation of chronic liver disease (5.3)
- Thyroid dysfunction (5.4)

----- **ADVERSE REACTIONS** -----

Adverse reactions caused by Lipiodol include hypersensitivity reactions, pulmonary embolism, pulmonary dysfunction, exacerbation of liver disease, procedural complications, abdominal pain, fever, nausea, vomiting, and thyroid dysfunction. (6)

To report SUSPECTED ADVERSE REACTIONS, contact GUERBET LLC at 1-877-729-6679 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch) (6)

Revised: 5/2018

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

2.1 Dosing Guidelines

2.2 Drug Handling

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Pulmonary and Cerebral Embolism

5.2 Hypersensitivity Reactions

5.3 Exacerbation of Chronic Liver Disease

5.4 Thyroid Dysfunction

**6 ADVERSE REACTIONS**

6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

7.1 Interference with Iodine-Based Diagnostic Tests and Iodine-Based Radiotherapy

**8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Nursing Mothers

8.3 Pediatric Use

8.4 Geriatric Use

8.5 Renal Impairment

**10 OVERDOSAGE**

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**16 HOW SUPPLIED/STORAGE AND HANDLING**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### **⚠WARNING: FOR INTRALYMPHATIC, INTRAUTERINE AND SELECTIVE HEPATIC INTRA-ARTERIAL USE ONLY**

**Pulmonary and cerebral embolism can result from inadvertent intravascular injection or intravasation of Lipiodol.**

**Inject Lipiodol slowly with radiologic monitoring; do not exceed recommended dose (5.1).**

## 1 INDICATIONS AND USAGE

Lipiodol is an oil-based radio-opaque contrast agent indicated for:

- hysterosalpingography in adults
- lymphography in adult and pediatric patients
- selective hepatic intra-arterial use for imaging tumors in adults with known hepatocellular carcinoma (HCC)

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Dosing Guidelines

Draw Lipiodol into a disposable syringe.

Use the smallest possible amount of Lipiodol according to the anatomical area to be visualized.

#### **Hysterosalpingography**

Using aseptic technique inject Lipiodol into the endometrial cavity with fluoroscopic control. Inject increments of 2 mL of Lipiodol until tubal patency is determined; stop the injection if patient develops excessive discomfort. Re-image after 24 hours to establish whether Lipiodol has entered the peritoneal cavity.

Before using Lipiodol exclude the presence of these conditions: pregnancy, uterine bleeding and endocervicitis, acute pelvic inflammatory disease, the immediate pre- or postmenstrual phase or within 30 days of curettage or conization.

#### **Lymphography**

Inject Lipiodol into a lymphatic vessel under radiologic guidance to prevent inadvertent venous administration or intravasation.

Adults:

- unilateral lymphography of the upper extremities 2 to 4 mL
- unilateral lymphography of the lower extremities 6 to 8 mL
- penile lymphography 2 to 3 mL
- cervical lymphography 1 to 2 mL

Pediatric patients:

- Inject a minimum of 1 mL to a maximum of 6 mL according to the anatomical area to be visualized. Do not exceed 0.25 mL/kg.

The following method is recommended for lymphography of the upper or lower extremities. Start the injection of Lipiodol into a lymphatic channel at a rate not to exceed 0.2 mL per minute. Inject the total dose of Lipiodol in no less than 1.25 hours. Use frequent radiologic monitoring to determine the appropriate injection rate and to follow the progress of Lipiodol within the lymphatics. Interrupt the injection if the patient experiences pain. Terminate the injection if lymphatic blockage is present to

minimize introduction of Lipiodol into the venous circulation via lymphovenous channels. Terminate the injection as soon as Lipiodol is radiographically evident in the thoracic duct to minimize entry of Lipiodol into the subclavian vein and pulmonary embolization. Obtain immediate post-injection images. Re-image at 24 or 48 hours to evaluate nodal architecture.

### **Selective Hepatic Intra-arterial Injection**

Determine the dose depending on the tumor size, local blood flow in the liver and in the tumor(s).

- Inject from 1.5 to 15 mL slowly under continuous radiologic monitoring. Stop the injection when stagnation or reflux is evident. Limit the dose to only the quantity required for adequate visualization. The total dose of Lipiodol administered should not exceed 20 mL.

## **2.2 Drug Handling**

Inspect Lipiodol visually for particulate matter and discoloration before administration. Do not use the solution if particulate matter is present or if the container appears damaged. Lipiodol is a clear, pale yellow to amber colored oil; do not use if the color has darkened.

Draw Lipiodol into a disposable syringe and use promptly. Discard any unused portion of Lipiodol.

## **3 DOSAGE FORMS AND STRENGTHS**

Each milliliter of Lipiodol contains 480 mg/mL of Iodine organically combined with ethyl esters of fatty acids of poppy seed oil.

## **4 CONTRAINDICATIONS**

Lipiodol is contraindicated in patients with hypersensitivity to Lipiodol, hyperthyroidism, traumatic injuries, recent hemorrhage or bleeding.

### **Hysterosalpingography**

Lipiodol hysterosalpingography is contraindicated in pregnancy, acute pelvic inflammatory disease, marked cervical erosion, endocervicitis and intrauterine bleeding, in the immediate pre-or postmenstrual phase, or within 30 days of curettage or conization.

### **Lymphography**

Lipiodol Lymphography is contraindicated in patients with a right to left cardiac shunt, advanced pulmonary disease, tissue trauma or hemorrhage advanced neoplastic disease with expected lymphatic obstruction, previous surgery interrupting the lymphatic system, radiation therapy to the examined area.

### **Selective Hepatic Intra-arterial Use Patients with HCC**

Lipiodol use is contraindicated in areas of the liver where the bile ducts are dilated unless external biliary drainage was performed before injection.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Pulmonary and Cerebral Embolism**

Pulmonary embolism may occur immediately or after a few hours to days from inadvertent systemic vascular injection or intravasation of Lipiodol and cause decreased pulmonary diffusing capacity and pulmonary blood flow, pulmonary infarction, acute respiratory distress syndrome and fatalities. Embolization of Lipiodol to brain and other major organs may occur. Avoid use of Lipiodol in patients with severely impaired lung function, cardiorespiratory failure, or right-sided cardiac overload. Perform radiological monitoring during the Lipiodol injection. Do not exceed the recommended maximum dose and rate of injection of Lipiodol. During lymphography to minimize the risk of pulmonary embolism obtain radiographic confirmation of intralymphatic (rather than venous) injection,

and terminate the procedure when Lipiodol becomes visible in the thoracic duct or lymphatic obstruction is observed.

## 5.2 Hypersensitivity Reactions

Anaphylactoid and anaphylactic reactions with cardiovascular, respiratory or cutaneous manifestations, ranging from mild to severe, including death, have uncommonly occurred following Lipiodol administration. Avoid use in patients with a history of sensitivity to other iodinated contrast agents, bronchial asthma or allergic disorders because of an increased risk of a hypersensitivity reaction to Lipiodol. Administer Lipiodol only in situations where trained personnel and therapies are promptly available for the treatment of hypersensitivity reactions, including personnel trained in resuscitation; ensure continuous medical monitoring and maintain an intravenous access line. Most hypersensitivity reactions to Lipiodol occur within half an hour after administration. Delayed reactions can occur up to several days after administration. Observe patients for signs and symptoms of hypersensitivity reactions during and for at least 30 minutes following Lipiodol administration.

## 5.3 Exacerbation of Chronic Liver Disease

Lipiodol hepatic intra-arterial administration can exacerbate the following conditions: portal hypertension and cause variceal bleeds due to obstruction of the intrahepatic portal channels by opening a pre sinusoidal anastomosis; hepatic ischemia and cause liver enzyme elevations, fever and abdominal pain; hepatic failure and cause ascites and encephalopathy. Hepatic vein thrombosis, irreversible liver insufficiency and fatalities have been reported. Procedural risks include vascular complications and infections.

## 5.4 Thyroid Dysfunction

Iodinated contrast media can affect thyroid function because of the free iodine content and can cause hyperthyroidism or hypothyroidism in predisposed patients. Patients at risk are those with latent hyperthyroidism and those with Hashimoto thyroiditis, or history of thyroid irradiation. As Lipiodol may remain in the body for several months, thyroid diagnostic results can be affected for up to two years after lymphography.

## 6 ADVERSE REACTIONS

### 6.2 Postmarketing Experience

The following adverse reactions (Table 1) have been identified during post approval use of Lipiodol. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The following adverse reactions are described in more detail in other sections of the prescribing information:

Pulmonary and cerebral embolism [see *Warnings and Precautions (5.1)*]

Hypersensitivity reactions [see *Warnings and Precautions (5.2)*]

Exacerbation of chronic liver disease [see *Warnings and Precautions (5.3)*]

**Table 1: Adverse Reactions in the Postmarketing Experience**

System Organ Class	Adverse Reaction
Endocrine disorders	hypothyroidism, hyperthyroidism, thyroiditis
Eye disorders	retinal vein thrombosis
Gastrointestinal disorders	nausea, vomiting, diarrhea
General disorders and administration site	

conditions	fever, pain, granuloma
Hepatobiliary disorders	hepatic vein thrombosis
Immune system disorders	hypersensitivity, anaphylactic reaction, anaphylactoid reaction
Nervous system disorders	cerebral embolism
Respiratory, thoracic and mediastinal disorders	pulmonary embolism, dyspnea, cough, acute respiratory distress syndrome
Urinary system disorders	renal insufficiency

### **Hysterosalpingography**

Abdominal pain, foreign body reactions, exacerbation of pelvic inflammatory disease.

### **Lymphography**

Cardiovascular collapse, lymphangitis, thrombophlebitis, edema or exacerbation of preexisting lymphedema, dyspnea and cough, fever, iodism (headache, soreness of mouth and pharynx, coryza and skin rash), allergic dermatitis, lipogranuloma, delayed healing at the site of incision.

### **Selective Hepatic Intra-arterial Injection**

Fever, abdominal pain, nausea, and vomiting are the most common reactions; other reactions include hepatic ischemia, liver enzymes abnormalities, transitory decrease in liver function, liver decompensation and renal insufficiency. Procedural risks include vascular complications and infections.

## **7 DRUG INTERACTIONS**

### **7.1 Interference with Iodine-Based Diagnostic Tests and Iodine-Based Radiotherapy**

Following Lipiodol administration, ethiodized oil remains in the body for several months, and may interfere with thyroid function testing for up to two years. Ethiodized oil interferes with radioactive iodine uptake by thyroid tissue for several weeks to months and may impair visualization of thyroid scintigraphy and reduce effectiveness of iodine 131 treatment.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### **Pregnancy Category C**

#### **Risk Summary**

There are no adequate and well-controlled studies of Lipiodol effects in pregnant women. Use Lipiodol during pregnancy only if clearly needed.

#### **Human Data**

It is not known whether Lipiodol can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. The use of Lipiodol during pregnancy causes iodine transfer which may interfere with the thyroid function of the fetus and result in brain damage and permanent hypothyroidism. Institute thyroid function testing and careful medical monitoring of the neonate exposed to Lipiodol in utero.

#### **Animal Data**

Animal reproduction studies have not been conducted using the indicated routes of administration of Lipiodol. Lipiodol was not embryotoxic or teratogenic in rats after oral administration of up to 110 mg Iodine/kg each day between gestation days 6 to 17, or in rabbits after 4-5 intermittent (once every three

days) oral administrations of 12.5 mg Iodine/kg between gestation days 6 to 18.

## **8.2 Nursing Mothers**

No nonclinical lactation studies of Lipiodol have been reported.

Lipiodol is excreted in human milk. Avoid use of Lipiodol in a nursing woman because of risk of hypothyroidism in nursing infants. If breastfeeding is continued the neonate's thyroid function should be monitored.

## **8.3 Pediatric Use**

For lymphography use a dose of minimum of 1 mL to a maximum of 6 mL according to the anatomical area to be visualized. Do not exceed 0.25 mL/kg. Administer the smallest possible amount of Lipiodol according to the anatomical area to be visualized.

## **8.4 Geriatric Use**

There are no studies conducted in geriatric patients.

## **8.5 Renal Impairment**

Prior to an intra-arterial administration of Lipiodol screen all patients for renal dysfunction by obtaining history and/or laboratory tests. Consider follow-up renal function assessments for patients with a history of renal dysfunction.

## **10 OVERDOSAGE**

Overdose may lead to respiratory, cardiac or cerebral complications, which can potentially be fatal. Microembolisms to multiple organs may occur more frequently after overdose. Promptly initiate symptomatic treatment and support of vital functions.

## **11 DESCRIPTION**

Lipiodol, ethiodized oil injection, is a sterile injectable radio-opaque agent. Each milliliter contains 480 mg of Iodine organically combined with ethyl esters of fatty acids of poppy seed oil. The precise structure of Lipiodol is unknown.

Lipiodol is a sterile, clear, pale yellow to amber colored oil. Lipiodol has a viscosity of 34 – 70 mPa·s at 20°C, and a density of 1.28 g/cm<sup>3</sup> at 20°C.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Ethiodized oil is an iodinated poppy seed oil based contrast agent.

### **12.3 Pharmacokinetics**

Following intra-arterial administration of Lipiodol, ethiodized oil retained in normal hepatic parenchyma is phagocytized by the Kupffer cells of the liver and washed out via the hepatic lymphatic system in about 2 to 4 weeks. In HCC, retention in the liver tumor is prolonged, allowing re-imaging of the tumor for four weeks or longer.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term studies in animals have not been performed to evaluate carcinogenic potential, or whether Lipiodol can affect fertility in males or females. Lipiodol did not demonstrate mutagenic potential in bacterial reverse mutation assays (*in vitro*), in a chromosomal aberration test in the mouse lymphoma assay (*in vitro*), and was negative in an *in vivo* micronucleus test in rats after intravenous injection of 479 mg I/kg.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Lipiodol is supplied in a box of one 10 mL ampoule, NDC 67684-1901-1.

Lipiodol is supplied in a box of one 10 mL vial, NDC 67684-1901-2.

Each vial is closed with a rubber stopper and sealed with an aluminum cap.

Store at controlled room temperature 15°-30°C (59°-86°F) [see USP, Controlled Room Temperature (CRT)]. Protect from light. Remove from carton only upon use.

Rx Only

Guerbet LLC

821 Alexander Road, Suite 204

Princeton, New Jersey 08540, USA

For further information or ordering, call

1-877-729-6679

Ampoules manufactured for Guerbet LLC by:

Jubilant HollisterStier General Partnership, Canada

Vials manufactured for Guerbet LLC by: Recipharm Monts, France

**Revised 05/2018**



# Lipiodol®

(Ethiodized Oil) Injection

NDC 67684-1901-1

Manufactured for Guerbet LLC by:  
Jubilant HollisterStier General Partnership  
16751 Trans-Canada Highway  
Kirkland, Quebec, Canada H9H 4J4

Distributed by Guerbet LLC  
821 Alexander Road, Suite 204  
Princeton, New Jersey 08540,  
USA



**Lipiodol®**  
(Ethiodized Oil) Injection  
4.8 g Iodine per 10 mL  
(480 mg Iodine per mL)  
**STERILE LIQUID FOR INJECTION**  
10 mL ampoule

**R** only

**Lipiodol®**  
(Ethiodized Oil) Injection  
4.8 g Iodine per 10 mL  
(480 mg Iodine per mL)

**STERILE LIQUID  
FOR INJECTION**

For Intrathecal,  
Intralymphatic and  
Selective Hepatic  
Intra-arterial  
Administration Only

For Single Use Only,  
Discard Unused Portion.

**Guerbet** |

10 mL ampoule

4.8 g Iodine per 10 mL  
(480 mg Iodine per mL)

USUAL DOSAGE: See package insert.

Do not open carton until ready for use.

Store between 15 and 30°C (59-86°F).

Store in the original packaging protected from light.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to use.

**R** only

**Lipiodol®**  
(Ethiodized Oil) Injection  
4.8 g Iodine per 10 mL  
(480 mg Iodine per mL)

**STERILE LIQUID  
FOR INJECTION**

For Intrathecal,  
Intralymphatic and  
Selective Hepatic  
Intra-arterial  
Administration Only

For Single Use Only,  
Discard Unused Portion.

**Guerbet** |

10 mL ampoule



Lot:  
Exp.:

#####

**Rx** only

NDC 67684-1901-1

# Lipiodol<sup>®</sup>

(Ethiodized Oil) Injection

**4.8 g Iodine per 10 mL**  
**(480 mg Iodine per mL)**

##  
##  
##  
##

For Intrauterine, Intralymphatic  
and Selective Hepatic  
Intra-arterial Administration Only

USUAL DOSAGE: See package insert.  
Protect from light. Single Use Only,  
Discard Unused Portion.

**Guerbet LLC** - 821 Alexander Rd,  
Ste 204 - Princeton, NJ 08540, USA



**10mL**



(01)00367684190114

Lot

Exp



# LIPIODOL

ethiodized oil injection

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:67684-1901
<b>Route of Administration</b>	INTRALYMPHATIC, INTRAUTERINE, INTRA-ARTERIAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ETHIODIZED OIL (UNII: KZW0R0686Q) (IODINE - UNII:9679TC07X4)	IODINE	480 mg in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
POPPY SEED OIL (UNII: 9G3Z76ES9J)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67684-1901-1	1 in 1 CARTON	03/21/2014	
1		10 mL in 1 AMPULE; Type 0: Not a Combination Product		
2	NDC:67684-1901-2	1 in 1 CARTON	09/28/2018	
2		10 mL in 1 VIAL; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA009190	03/21/2014	

**Labeler** - Guerbet LLC (037876096)

**Registrant** - Guerbet LLC (037876096)

## Establishment

Name	Address	ID/FEI	Business Operations
Simafex		381108273	api manufacture(67684-1901)

## Establishment

Name	Address	ID/FEI	Business Operations
Jubilant HollisterStier General Partnership		246762764	manufacture(67684-1901) , pack(67684-1901)

## Establishment

Name	Address	ID/FEI	Business Operations
RECIPHARM MONTS		777990813	manufacture(67684-1901) , pack(67684-1901)

## Establishment

Name	Address	ID/FEI	Business Operations
IFTS		270199391	analysis(67684-1901)

Revised: 1/2020

Guerbet LLC