

**CORICIDIN HBP NIGHTTIME MULTI-SYMPTOM COLD-  
acetaminophen,dextromethorphan hydrobromide, doxylamine  
succinate liquid  
Bayer HealthCare LLC.**

**Coricidin HBP Nighttime Multi-Symptom Cold Liquid**

***Drug Facts***

***Active ingredients (in each 15 mL)***

|                                     |                             |
|-------------------------------------|-----------------------------|
| Acetaminophen 325 mg                | Pain reliever/fever reducer |
| Dextromethorphan hydrobromide 15 mg | Cough suppressant           |
| Doxylamine succinate 6.25 mg        | Antihistamine               |

***Purpose***

|                                     |                             |
|-------------------------------------|-----------------------------|
| Acetaminophen 325 mg                | Pain reliever/fever reducer |
| Dextromethorphan hydrobromide 15 mg | Cough suppressant           |
| Doxylamine succinate 6.25 mg        | Antihistamine               |

***Uses***

temporarily relieves common cold/flu symptoms:

- headache
- sore throat
- runny nose
- minor aches and pains
- cough due to minor throat and bronchial irritation
- sneezing
- fever

***Warnings***

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 120 mL (4 doses) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

acetaminophen may cause severe skin or severe allergic reactions. Symptoms may

include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- glaucoma
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- redness or swelling is present

**If pregnant or breast-feeding**

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children**

**Keep out of reach of children.** In case of overdose, get medical help

or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed
- adults and children 12 years and over: take 30 mL in the dosing cup provided every 6 hours while symptoms last
- do not exceed 120 mL (4 doses) in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- **each 15 mL contains:** sodium 18 mg
- store at 20°-25°C (68°-77°F)

***Inactive ingredients***

alcohol 10%, anhydrous citric acid, FD&C blue no.1, FD&C red no.40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

***Questions or comments***

***Questions or comments?*** 1-800-317-2165 (Mon – Fri 9AM – 5PM EST)



### Product Characteristics

|          |        |              |  |
|----------|--------|--------------|--|
| Color    |        | Score        |  |
| Shape    |        | Size         |  |
| Flavor   | CHERRY | Imprint Code |  |
| Contains |        |              |  |

### Packaging

| # | Item Code        | Package Description                                            | Marketing Start Date | Marketing End Date |
|---|------------------|----------------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:11523-4358-1 | 355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 11/20/2015           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| OTC Monograph Drug | M012                                     | 11/20/2015           |                    |

**Labeler** - Bayer HealthCare LLC. (112117283)

Revised: 12/2024

Bayer HealthCare LLC.