

LIDOCAINE PAIN RELIEF GEL -PATCH- lidocaine patch
Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.

83602-230

Active ingredient

Lidocaine 4% Purpose: Topical Anesthetic

Purpose

Topical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Ask a doctor if:

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

Stop use if:

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

Pregnancy or breast-feeding

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children

Keep out of reach of children. If swallowed accidentally , get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application

Children under age of 12:Consult a doctor

Other information

Store at 20°C to 25°C (68°F - 77°F)

Store in a cool dry place away from direct sunlight

Inactive ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, PropyleneGlycol, Polyacrylic Acid, Povidone, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments?

supports@sumifunofficial.com

Package label. Principal display panel

Maximum Strength

CUT OPEN HERE

Child Resistant Packaging

Sumifun[®]

Lidocaine

Pain Relief Gel-Patch

Desensitizes Aggravated Nerves And Relieves Pain:

- NO Irritation
- Stay-put Flexible Patch
- Numbs Pain Away

150% LARGER*

*Than 3.94" x 5.5" Sumifun large patch



XXL PATCH

1 PATCH
9.84" X 5.5"
(25cm X 14cm)

Drug Facts

Active ingredient	Purpose
Lidocaine 4%	Topical Anesthetic

Uses
Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings
For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding,
ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.
If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions
Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application

Children under age of 12: Consult a doctor

Other Information


- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive ingredients
Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, PropyleneGlycol, Polyacrylic Acid, Povidone, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments:
supports@sumifunofficial.com

Business Cooperation:
business@sumifunofficial.com

Child Resistant Packaging



13
PATCHES

Lidocaine

www.sumifunofficial.com
YouTube: @sumifun
Facebook: @sumifunOfficial
Instagram: @sumifunofficial
Twitter: @sumifunOfficial

Maximum Strength

Sumifun®

Lidocaine

Pain Relief Gel-Patch

Desensitizes Aggravated Nerves And Relieves Pain:

- NO Irritation
- Stay-put Flexible Patch
- Numbs Pain Away

150%
LARGER*
*than 3.54" x 5.5" Sumifun large patch

XXL PATCH

13 PATCHES
9.84" X 5.5"
(25cm X 14cm)

Guangzhou Lidaquanniao Import & Export Trade Co., Ltd.
478 Baibeiqiang A, No. 6, Dongfeng 5th Road,
Zhuou, Taishan District, Guangzhou, Guangdong, China.

www.cphc.com

Usage



Remove patch from outer package.



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

Sumifun

Drug Facts

Active ingredient Purpose
Lidocaine 4% Topical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or sores
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding,

use a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application.

Children under age of 12: Consult a doctor

Other Information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive ingredients

Dihydroxyaluminum Aminoacetate, Ectoin, Disodium Glycinate, Kaolin, Methylparaben, Polyacrylate 80, Propylene Glycol, Polyacrylic Acid, Povidone, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments:

support@sumifunofficial.com
Business Cooperation:
business@sumifunofficial.com

Child Resistant Packaging



10
PATCHES

Lidocaine

www.sumifunofficial.com
YouTube: @sumifun
Facebook: @sumifunOfficial
Twitter: @sumifunOfficial
Instagram: @sumifunofficial
TikTok: @sumifunofficial

Maximum Strength

Sumifun® Lidocaine Pain Relief Gel-Patch

Desensitizes Aggravated
Nerves And Relieves Pain:

- NO Irritation
- Stay-put Flexible Patch
- Numbs Pain Away

**150%
LARGER***

*Than 3.94" x 5.5" Sumifun large patch

XXL PATCH

10 PATCHES
9.84" X 5.5"
(25cm X 14cm)

Guangzhou Lidaqianmai Import & Export Trade Co., Ltd.
412 Building A, No. 8, Dong Heng San Road,
Zhuocun, Tanhe District, Guangzhou, Guangdong, China.

©2020 Sumifun

Usage



Remove patch from
outer package



Partially peel back
protective film



Place on
affected area



Remove remaining
film while
pressing patch to skin

Sumifun

Drug Facts

Active ingredient Lidocaine 4% **Purpose** Topical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knee, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or tissues
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 6-hour application

Children under age of 12: Consult a doctor

Other information

- Store at 20°C to 25°C (68°F - 77°F)
- Store in a cool dry place away from direct sunlight

Inactive ingredients

Dihydroxyaluminum Ammoniate, Edidate Disodium, Glycerin, Kaolin, Methybaraben, Polybutene 88, Propylene Glycol, Polyacrylic Acid, Povidone, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water

Questions or Comments:

support@sumifunofficial.com
business@sumifunofficial.com

Child Resistant Packaging



6 9 2 4 9 0 1 3 2 3 3



LIDOCAINE PAIN RELIEF GEL -PATCH

lidocaine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83602-230
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.04 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
GLYCERIN (UNII: PDC6A3C0OX)	
POVIDONE K90 (UNII: RDH86HJV5Z)	

METHYLPARABEN (UNII: A2I8C7HI9T)
PROPYLPARABEN (UNII: Z8IX2SC1OH)
EDETATE DISODIUM (UNII: 7FLD91C86K)
KAOLIN (UNII: 24H4NWX5CO)
TARTARIC ACID (UNII: W4888I119H)
WATER (UNII: 059QF0KO0R)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JN12J)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83602-230-13	13 in 1 BOX	05/08/2026	
1		1 in 1 BAG		
1		30 g in 1 PATCH; Type 0: Not a Combination Product		
2	NDC:83602-230-16	18 in 1 BOX	05/08/2026	
2		1 in 1 BAG		
2		30 g in 1 PATCH; Type 0: Not a Combination Product		
3	NDC:83602-230-10	10 in 1 BOX	05/08/2026	
3		1 in 1 BAG		
3		30 g in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	05/08/2026	

Labeler - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Registrant - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Revised: 5/2026

Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.