

**MULTI SYMPTOM FLU RELIEF MAX STRENGTH DAYTIME NIGHTTIME COMBO PACK- acetaminophen, dextromethorphan hbr, and acetaminophen, chlorpheniramine maleate, dextromethorphan hbr  
CVS PHARMACY**

-----  
**Theraflu Flu Relief Max Strength Combo Pack *Drug Facts***

**Theraflu Flu Relief Max Strength\* Daytime Powder**

***Active ingredients (in each packet)***

Acetaminophen 1000 mg

Dextromethorphan HBr 30 mg

***Purposes***

Pain reliever/Fever reducer

Cough suppressant

***Uses***

- temporarily relieves these symptoms due to a common cold:
  - headache
  - minor aches and pains
  - cough due to minor throat and bronchial irritation
  - minor sore throat pain
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

### **Do not use**

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

### **When using this product**

- **do not exceed recommended dosage.**

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.  
These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not

notice any signs or symptoms.

### ***Directions***

- **do not use more than directed**
- take every 6 hours, while symptoms persist. Do not take more than 3 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume the entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

### ***Other information***

- **each packet contains:** potassium 6 mg
- store at room temperature. Protect from excessive heat and moisture.

### ***Inactive ingredients***

anhydrous citric acid, caramel, flavor, maltodextrin, potassium chloride, silica, sucralose, sucrose.

### ***Questions or comments?***

**1-866-467-2748**

## **Theraflu Flu Relief Max Strength Nighttime\* Powder**

### ***Drug Facts***

#### ***Active ingredients (in each packet)***

Acetaminophen 1000 mg

Chlorpheniramine maleate 4 mg

Dextromethorphan HBr 30 mg

## **Purposes**

Pain reliever/Fever reducer

Antihistamine

Cough suppressant

## **Uses**

- temporarily relieves these symptoms due to a common cold:
  - headache
  - minor aches and pains
  - cough due to minor throat and bronchial irritation
  - minor sore throat pain
  - runny nose
- temporarily reduces fever

## **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

## **Do not use**

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAO, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.  
These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not use more than directed**

- take every 6 hours, while symptoms persist. Do not take more than 3 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat

### ***Other information***

- **each packet contains:** potassium 6 mg
- store at room temperature. Protect from excessive heat and moisture.

### ***Inactive ingredients***

anhydrous citric acid, caramel, flavor, maltodextrin, potassium chloride, silica, sucralose, sucrose.

### ***Questions or comments?***

**1-866-467-2748**

### **Other Safety Information**

**READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE. KEEP CARTON FOR REFERENCE. DO NOT DISCARD.**

**DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME.**

**DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.**

**DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF THE DAYTIME PRODUCT UNLESS DIRECTED BY YOUR DOCTOR.**

**TAMPER-EVIDENT INNER UNIT. DO NOT USE IF NECKBAND PRINTED WITH "SEALED FOR SAFETY" IS TORN OR MISSING.**

**Distributed by:**

## **Principal Display Panel**

\*Compare to the active ingredients in Theraflu Multi-Symptom Flu Relief Max Strength\*\*  
Daytime & Nighttime.

**NDC 51316-559-06**

***FLU RELIEF***

***MAX STRENGTH\*\****

**6 x DAYTIME**

**Acetaminophen** Pain Reliever/Fever Reducer

**Dextromethorphan HBr** Cough Suppressant

**6 x NIGHTTIME**

**Acetaminophen** Pain Reliever/Fever Reducer

**Chlorpheniramine Maleate** Antihistamine

**Dextromethorphan HBr** Cough Suppressant

**Hot liquid therapy that relieves:**

- Fever
- Body ache
- Headache
- Sore throat pain
- Cough
- Runny nose (Nighttime only)

**Free from synthetic dyes**


**6 DAYTIME PACKETS**

**6 NIGHTTIME PACKETS**

**12 TOTAL PACKETS**

**Honey Lemon**

**Natural & Artificial Flavored**



# MULTI SYMPTOM FLU RELIEF MAX STRENGTH DAYTIME NIGHTTIME COMBO PACK

acetaminophen, dextromethorphan hbr, and acetaminophen, chlorpheniramine maleate, dextromethorphan hbr kit

## Product Information

**Product Type** HUMAN OTC DRUG      **Item Code (Source)** NDC:51316-559

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-559-06	1 in 1 CARTON; Type 1: Convenience Kit of Co-Package	04/21/2023	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 CARTON	6
Part 2	1 CARTON	6



## Part 1 of 2

### MS FLU RELIEF MAX STRENGTH DAYTIME

acetaminophen, dextromethorphan hbr powder

#### Product Information

Item Code (Source) NDC:51316-558

Route of Administration ORAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg

#### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CAMEL (UNII: T9D99G2B1R)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POTASSIUM CHLORIDE (UNII: 660YQ98I10)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SUCROSE (UNII: C151H8M554)	

#### Product Characteristics

Color	WHITE (to off white, yellow, beige, and brown color)	Score	
Shape		Size	
Flavor	HONEY (Lemon)	Imprint Code	
Contains			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-558-06	6 in 1 CARTON; Type 1: Convenience Kit of Co-Package		

#### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/21/2023	

## Part 2 of 2

### MS FLU RELIEF MAX STRENGTH NIGHTTIME

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr powder

#### Product Information

**Item Code (Source)** NDC:51316-560

**Route of Administration** ORAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg

#### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>CAMEL</b> (UNII: T9D99G2B1R)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>POTASSIUM CHLORIDE</b> (UNII: 660YQ98I10)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>SUCROSE</b> (UNII: C151H8M554)	

#### Product Characteristics

<b>Color</b>	WHITE (white to off-white, yellow, beige, and brown color)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	HONEY (lemon)	<b>Imprint Code</b>	
<b>Contains</b>			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-560-06	6 in 1 CARTON; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/21/2023	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/21/2023	

**Labeler** - CVS PHARMACY (062312574)

Revised: 3/2024

CVS PHARMACY