FEXOFENADINE HYDROCHLORIDE AND PSEUDOEPHEDRINE HYDROCHLORIDE - fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended release

HEB

Fexofenadine Hydrochloride and Pseudoephedrine Hydrochloride Extended-Release Tablets, 60 mg/120 mg

Drug Facts

Active ingredients (in each extended-release tablet)	Purpose
Fexofenadine HCl, USP 60 mg	Antihistamine
Pseudoephedrine HCl, USP 120 mg	Nasal Decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids

• do not take with fruit juices (see Directions)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

• do not divide, crush, chew or dissolve the tablet; swallow tablet whole

vears of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information

- do not use if carton is opened or if individual blister units are torn or opened
- store between 20° to 25°C (68° to 77°F)
- USP dissolution test is pending.

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, ethyl cellulose, ferric oxide yellow, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid.

Questions or comments?

Call toll-free **1-800-818-4555 weekdays**

PRINCIPAL DISPLAY PANEL - 30 Tablet Blister Pack Carton

Compare to Allegra-D® active ingredients*

NDC 37808-999-30

H-E-B_®

Allergy Relief Fexofenadine HCl 60 mg/antihistamine pseudoephedrine HCl 120 mg/nasal decongestant Extended-Release Tablets, USP Allergy & Congestion Indoor & Outdoor Allergies

12 Hour

Non-Drowsy Original Prescription Strength

12 Hour Relief of:

- Nasal and Sinus Congestion Due to Colds or Allergies
- Sneezing; Runny Nose; Itchy, Watery Eyes and Itchy Nose or Throat Due to Allergies

30 EXTENDED RELEASE TABLETS

actual size



fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended release

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-999	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
FEXO FENADINE HYDRO CHLO RIDE (UNII: 2S068B75ZU) (FEXOFENADINE - UNII: E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	60 mg	
PSEUDO EPHEDRINE HYDRO CHLO RIDE (UNII: 6 V9 V2RYJ8 N) (PSEUDO EPHEDRINE - UNII:7CUC9 DDI9 F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg	

Inactive Ingredients			
Ingredient Name	Strength		
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)			
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)			
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
PO VIDO NE K30 (UNII: U725QWY32X)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
DIBASIC CALCIUM PHO SPHATE DIHYDRATE (UNII: O7TSZ97GEP)			
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)			
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)			
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			

Product Characteristics			
Color	WHITE, YELLOW	Score	no score
Shape	CAPSULE (bilayer)	Size	17mm
Flavor		Imprint Code	724
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:37808-999- 30	1 in 1 CARTON	08/01/2018		
1		30 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090818	08/01/2018	

Labeler - HEB (007924756)

Establishment			
Name	Address	ID/FEI	Business Operations
Sun Pharmaceutical Industries Limited		650445203	ANALYSIS(37808-999), MANUFACTURE(37808-999)

Revised: 7/2018 HE B