

NIGHTTIME COLD AND FLU- acetaminophen,dextromethorphan,doxylamine capsule, liquid filled

Velocity Pharma

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Acetaminophen, Dextromethorphan and Doxylamine Night time Cold and Flu

Active Ingredient

(in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine Succinate 6.25 mg

Purpose

pain reliever

Cough Suppressant

Antihistamine

Uses

pain reliever, cough suppressant and Antihistamine

Warnings

Warnings Failure to follow these warnings could result in serious consequences.

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours which is maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

do not use:

- with any other drug containing acetaminophen (prescription or not prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (Certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- Heart disease
- Thyroid disease
- Diabetes
- High blood pressure
- Trouble urinating due to enlarged prostate gland

- **ask your doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product, do not use more than directed.**
- **Stop use and ask a doctor if:**
 - Redness or swelling is present
 - You get nervous, dizzy or sleepless
 - Fever gets worse or lasts more than 3 days
 - New symptoms occur
 - Symptoms do not get better within 7 days or are accompanied by a fever

Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use.

Overdose warning: Taking more than recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Tamper evident: this package is safety sealed and child resistant. Use only if blisters are intact. If difficult to open use scissors.

Direction

- **do not exceed 4 doses per 24 hours**
- **take only as directed – see overdose warning**
- **Adults and children 12 years and over:** 2 softgels with water every 4 hours
- **children under 12 years:** ask a doctor
- **Children under 4 years:** do not use

Other Information

- store at room temperature

Inactive Ingredients

FD&C Red No.40, FD&C Yellow No. 6, Gelatin, Glycerin, Poyethylene Glycol, Povidone, Propylene Glycol, Purified Water, Sorbitol Special, Titanium dioxide

Questions or Comments

Call toll free 1-855-314-1850

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

03

VELOCITY PHARMA LLC
226 B Sherwood Avenue, Farmingdale, NY 11735
Distributed by
This product is not manufactured or distributed by Procter & Gamble.
distributor of Vicksi[®] M[®]Q[®]



MULTI SYMPTOM RELIEF
✓ Aches and Fever - Acetaminophen
✓ Cough - Dextromethorphan HBr
✓ Sneezing, Runny Nose -
Doxylamine Succinate



NightTime

Cold and Flu

NightTime
Cold and Flu

NDC 76168-056-02

Compare to the active ingredients in Vicksi[®] M[®]Q[®]

Inactive ingredients: DMC, Yellow No. 10, FD&C Blue No. 1, Gelatin, Glycerin, Polyethylene Glycol, Povidone, Propylene Glycol, Purified Water, Sorbitol Special, Titanium Dioxide.
Other information: Store at room temperature.
Questions call: 1-855-314-1850

Directions

- Do not exceed 4 doses per 24 hours.
 - Take only as directed - see Overdose warning
- | | |
|---------------------------------------|-------------------------------------|
| Adults and Children 12 years and over | 2 softgels with water every 6 hours |
| Children 4 to under 12 years | ask a doctor |
| Children under 4 years | do not use |

Drug Facts

Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine Succinate 6.25mg	Antihistamine

Drug Facts (continued)

Warnings: Failure to follow these warnings could result in serious consequences.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- More than 4 doses in 24 hours, which is the maximum daily amount for this product
- With other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Do not use:

- With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- Liver disease
- Heart disease
- Thyroid disease
- Diabetes
- High blood pressure
- Trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood-thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if:

- Redness or swelling is present
- Fever gets worse or lasts more than 3 days
- Symptoms do not get better within 7 days or are accompanied by a fever

If pregnant or breast-feeding: ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

Tamper evident: This package is safety sealed and child resistant. Use only if blisters are intact. If difficult to open, use scissors.

C31 01

NIGHTTIME COLD AND FLU

acetaminophen,dextromethorphan,doxylamine capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76 168-056
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL 1000 (UNII: U076Q6Q621)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	22mm
Flavor		Imprint Code	215
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76 168-056-02	12 in 1 BLISTER PACK	08/01/2017	
1		1 in 1 CARTON; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	05/01/2014	

Labeler - Velocity Pharma (962198409)

Registrant - Velocity Pharma (962198409)

Establishment

Name	Address	ID/FEI	Business Operations
medgel		677385498	manufacture(76168-056)

Revised: 9/2017

Velocity Pharma