

**ALKA-SELTZER PLUS MAXIMUM STRENGTH DAY COLD AND FLU-  
acetaminophen, dextromethorphan hydrobromide, phenylephrine  
hydrochloride capsule, liquid filled  
Bayer HealthCare LLC.**

-----  
**Alka-Seltzer Plus® Maximum Strength Day Cold & Flu Liquid Gels**

***Drug Facts***

***Active ingredients (in each capsule)***

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Phenylephrine hydrochloride 5 mg

***Purposes***

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

***Uses***

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains · headache · cough
- sore throat · nasal and sinus congestion
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening · blisters · rash · hives
- facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

## **Ask a doctor before use if you have**

- liver disease ● heart disease ● high blood pressure
- thyroid disease ● diabetes
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

## **When using this product do not exceed recommended dosage**

### **Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

**Other information**

**Other information**

- store at room temperature. Avoid excessive heat above 40°C (104°F).

**Inactive ingredients** FD&C yellow #6, FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

**Questions or comments?** 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

**Carton 20**

Alka-Seltzer PLUS

MAXIMUM STRENGTH

Cold

& Flu

DAY NON-DROWSY

ACETAMINOPHEN / Pain Reliever-Fever Reducer

Dextromethorphan Hydrobromide / Cough Suppressant

Phenylephrine HCl / Nasal Decongestant

- Cough
- Nasal Congestion
- Headache & Body Ache
- Sore Throat
- Sinus Pressure

20 LIQUID GELS (Liquid Filled Capsules)

LIFT HERE

**Drug Facts (continued)**

**Drug Facts**  
**Active ingredients (in each capsule)**  
 Acetaminophen 325 mg ..... Pain reliever/fever reducer  
 Dextromethorphan hydrobromide 10 mg.....Cough suppressant  
 Phenylephrine hydrochloride 5 mg .....Nasal decongestant

**Uses**  
 • temporarily relieves these symptoms due to a cold or flu:  
 • minor aches and pains  
 • cough  
 • nasal and sinus congestion  
 • temporarily reduces fever

**Warnings**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours  
 • with other drugs containing acetaminophen  
 • 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:  
 • skin reddening  
 • hives  
 • rash  
 • blisters  
 • asthma (wheezing)  
 • shock  
 If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Purposes**

**Do not use**  
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 • if you have ever had an allergic reaction to this product or any of its ingredients  
 • in children under 12 years of age

**Ask a doctor before use if you have**  
 • liver disease  
 • heart disease  
 • high blood pressure  
 • thyroid disease  
 • diabetes  
 • cough with excessive phlegm (mucus)  
 • difficulty in urination due to enlargement of the prostate gland  
 • persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not exceed recommended dosage**

**Drug Facts (continued)**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

MAXIMUM STRENGTH



**Cold & Flu**

DAY NON-DROWSY

MAXIMUM STRENGTH



**Cold & Flu**

DAY NON-DROWSY

**ACETAMINOPHEN / Pain Reliever-Fever Reducer**  
 Dextromethorphan Hydrobromide / Cough Suppressant  
 Phenylephrine HCl / Nasal Decongestant

- ▶ Cough
- ▶ Nasal Congestion
- ▶ Headache & Body Ache
- ▶ Sore Throat
- ▶ Sinus Pressure

**20 LIQUID GELS** (Liquid Filled Capsules)



6256

6256

8 0 1650055530



**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**Do not use if blister  
is cut or broken.  
DOES NOT CONTAIN ASPIRIN**

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Children even if you do not notice any signs or symptoms.

**Other Information**

- store at room temperature. Avoid excessive heat above 40°C (104°F).

**Inactive Ingredients** FD&C yellow #6, FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

**Questions or comments?** 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

**Drug Facts (continued)**

**Directions**

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

**ALKA-SELTZER PLUS MAXIMUM STRENGTH DAY COLD AND FLU**  
acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0280-1590
<b>Route of Administration</b>	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

## Inactive Ingredients

Ingredient Name	Strength
<b>GLYCERIN</b> (UNII: PDC6A3C00X)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	AS;DC
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-1590-20	2 in 1 CARTON	06/19/2017	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/19/2017	

**Labeler** - Bayer HealthCare LLC. (112117283)

Revised: 12/2023

Bayer HealthCare LLC.