

**ALKA-SELTZER PLUS MAXIMUM STRENGTH DAY COLD AND FLU-
acetaminophen, dextromethorphan hydrobromide, phenylephrine
hydrochloride capsule, liquid filled
Bayer HealthCare LLC.**

Alka-Seltzer Plus® Maximum Strength Day Cold & Flu Liquid Gels

Drug Facts

Active ingredients (in each capsule)

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Phenylephrine hydrochloride 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains · headache · cough
- sore throat · nasal and sinus congestion
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening · blisters · rash · hives
- facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease ● heart disease ● high blood pressure
- thyroid disease ● diabetes
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Other information

Other information

- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients FD&C yellow #6, FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

Carton 20

Alka-Seltzer PLUS

MAXIMUM STRENGTH

Cold

& Flu

DAY NON-DROWSY

ACETAMINOPHEN / Pain Reliever-Fever Reducer

Dextromethorphan Hydrobromide / Cough Suppressant

Phenylephrine HCl / Nasal Decongestant

- Cough
- Nasal Congestion
- Headache & Body Ache
- Sore Throat
- Sinus Pressure

20 LIQUID GELS (Liquid Filled Capsules)

LIFT HERE

Drug Facts (continued)

Drug Facts
Active ingredients (in each capsule)
 Acetaminophen 325 mg Pain reliever/fever reducer
 Dextromethorphan hydrobromide 10 mg.....Cough suppressant
 Phenylephrine hydrochloride 5 mgNasal decongestant

Uses
 • temporarily relieves these symptoms due to a cold or flu:
 • minor aches and pains
 • cough
 • nasal and sinus congestion
 • temporarily reduces fever

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
 • with other drugs containing acetaminophen
 • 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:
 • skin reddening
 • hives
 • rash
 • blisters
 • asthma (wheezing)
 • shock
 If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Purposes

Do not use
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 • if you have ever had an allergic reaction to this product or any of its ingredients
 • in children under 12 years of age

Ask a doctor before use if you have
 • liver disease
 • heart disease
 • high blood pressure
 • thyroid disease
 • diabetes
 • cough with excessive phlegm (mucus)
 • difficulty in urination due to enlargement of the prostate gland
 • persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Drug Facts (continued)
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

MAXIMUM STRENGTH



Cold & Flu

DAY NON-DROWSY



MAXIMUM STRENGTH



Cold & Flu

DAY NON-DROWSY

ACETAMINOPHEN / Pain Reliever-Fever Reducer
 Dextromethorphan Hydrobromide / Cough Suppressant
 Phenylephrine HCl / Nasal Decongestant

- ▶ Cough
- ▶ Nasal Congestion
- ▶ Headache & Body Ache
- ▶ Sore Throat
- ▶ Sinus Pressure

20 LIQUID GELS (Liquid Filled Capsules)

6256

6256





PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

**Do not use if blister
is cut or broken.
DOES NOT CONTAIN ASPIRIN**

Drug Facts (continued)

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Children even if you do not notice any signs or symptoms.

Other Information

- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive Ingredients FD&C yellow #6, FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

ALKA-SELTZER PLUS MAXIMUM STRENGTH DAY COLD AND FLU
acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-1590
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C00X)	
WATER (UNII: 059QF0K00R)	
SHELLAC (UNII: 46N107B71O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
GELATIN (UNII: 2G86QN327L)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SORBITAN (UNII: 6O92ICV9RU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	AS;DC
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-1590-20	2 in 1 CARTON	06/19/2017	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/19/2017	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 12/2025

Bayer HealthCare LLC.