

**SEVERE COLD AND FLU PLUS CONGESTION- acetaminophen,  
dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl  
COSTCO WHOLESALE CORPORATION**

-----

**Kirkland 44-795796-81**

**Day**

**Active ingredients (in each caplet)**

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

**Purpose**

Pain reliever/fever reducer  
Cough suppressant  
Expectorant  
Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
  - nasal congestion
  - sore throat
  - minor aches and pains
  - headache
  - sinus congestion and pressure
  - cough due to minor throat and bronchial irritation
  - reduces swelling of nasal passages
- temporarily reduces fever
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- diabetes
- thyroid disease
- heart disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

### **When using this product**

**do not exceed recommended dosage.**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed**
- do not take more than 8 caplets in 24 hours
- adults and children 12 years and over: take 2 caplets with water every 4 hours
- children under 12 years: ask a doctor

### **Other information**

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- see end flap for expiration date and lot number
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

### ***Inactive ingredients***

benzyl alcohol, corn starch, croscopovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, propylene glycol, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

### ***Questions or comments?***

**1-800-426-9391**

### ***Night***

#### ***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Doxylamine succinate 6.25 mg  
Phenylephrine HCl 5 mg

### ***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Antihistamine  
Nasal decongestant

### ***Uses***

- temporarily relieves common cold and flu symptoms:
  - nasal congestion
  - sore throat
  - minor aches and pains
  - headache
  - sinus congestion and pressure
  - cough due to minor throat and bronchial irritation
  - cough to help you sleep
  - runny nose and sneezing
- reduces swelling of nasal passages
- temporarily reduces fever
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

### ***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may

include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed**
- do not take more than 8 caplets in 24 hours
- adults and children 12 years and over: take 2 caplets with water every 4 hours
- children under 12 years: ask a doctor

**Other information**

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- see end flap for expiration date and lot number
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients**

benzyl alcohol, black iron oxide, corn starch, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, propylene glycol, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

**Principal display panel**

**KIRKLAND**

*Signature*

**COMPARE TO VICKS® DAYQUIL® & NYQUIL® VAPOCOOL®  
SEVERE COLD & FLU + CONGESTION**

*active ingredients\**

**NDC 63981-795-81**

ITM. / ART. 1729556

**COOL BLAST  
SEVERE COLD & FLU  
PLUS CONGESTION**

**Acetaminophen 325 mg,**  
*Dextromethorphan HBr 10 mg, Guaifenesin 200 mg,  
Phenylephrine HCl 5 mg*  
*Pain Reliever • Fever Reducer • Cough Suppressant*  
*Expectorant • Nasal Decongestant*  
*Relieves:*  
*Headache, Fever, Sore Throat,*  
*Minor Aches & Pains*  
*Chest Congestion, Thins & Loosens Mucus*  
*Nasal Congestion, Sinus Pressure*

**Acetaminophen 325 mg,**  
*Dextromethorphan HBr 10 mg, Doxylamine succinate 6.25 mg,  
Phenylephrine HCl 5 mg*  
*Pain Reliever • Fever Reducer • Cough Suppressant*  
*Antihistamine • Nasal Decongestant*  
*Relieves:*  
*Headache, Fever, Sore Throat,*  
*Minor Aches & Pains*  
*Nasal Congestion,  
Sinus Pressure*  
*Sneezing, Runny Nose*

*Nasal Congestion, Sinus Pressure*

*Cough*

*Actual*

*Size*

*112 DAY COATED CAPLETS*

*Cough*

*Actual*

*Size*

*56 NIGHT COATED CAPLETS*

**168**

**TOTAL CAPLETS**

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

**Do Not Take Daytime and Nighttime Products at the Same Time.**

\*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® and NyQuil® VapoCOOL® Severe Cold & Flu + Congestion.

50844 REV0223A79579681

Manufactured by: LNK INTERNATIONAL, INC.


60 Arkay Drive, Hauppauge, NY 11788 USA

For: Costco Wholesale Corporation

999 Lake Drive, Issaquah, WA 98027 USA

[www.costco.com](http://www.costco.com)

23V08014a



E7263R1

LOT NO. / EXP. DATE

**PARENTS:**  
Look for this symbol on the packaging to help you identify products that are safe for children.

**Parental Involvement:**  
Parents should be involved in their child's health care decisions.

**Child-Resistant Packaging:**  
This product is child-resistant, but it is not child-proof. Children under 5 years of age may still be able to open this container.

**Keep Out of Reach of Children:**  
Always keep this product out of the reach of children. Use proper storage to help prevent access to this product by children.

**Use Only as Directed:**  
Do not use this product if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**Warnings:**  
Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**Directions:**  
Read the Drug Facts carefully before using this product. Use only as directed. Do not exceed the recommended dose. Do not use for more than 10 days unless directed by a doctor. Do not use if symptoms persist or worsen. Do not use if you experience dizziness, lightheadedness, or other symptoms. Do not use if you experience dizziness, lightheadedness, or other symptoms.

**Other information:**  
This product contains acetaminophen, dextromethorphan, guaifenesin, and phenylephrine. Do not use if you are allergic to any of these ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**Keep Out of Reach of Children:**  
Always keep this product out of the reach of children. Use proper storage to help prevent access to this product by children.

**Use Only as Directed:**  
Do not use this product if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**Warnings:**  
Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**Directions:**  
Read the Drug Facts carefully before using this product. Use only as directed. Do not exceed the recommended dose. Do not use for more than 10 days unless directed by a doctor. Do not use if symptoms persist or worsen. Do not use if you experience dizziness, lightheadedness, or other symptoms. Do not use if you experience dizziness, lightheadedness, or other symptoms.

**Other information:**  
This product contains acetaminophen, dextromethorphan, guaifenesin, and phenylephrine. Do not use if you are allergic to any of these ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**KIRKLAND Signature**  
**COOL BLAST**  
**SEVERE COLD & FLU PLUS CONGESTION**

**Acetaminophen 325 mg, Dextromethorphan HBr 10 mg, Guaifenesin 200 mg, Phenylephrine HCl 5 mg**  
Pain Reliever • Fever Reducer • Cough Suppressant  
Expectorant • Nasal Decongestant

**Acetaminophen 325 mg, Dextromethorphan HBr 10 mg, Doxylamine succinate 6.25 mg, Phenylephrine HCl 5 mg**  
Pain Reliever • Fever Reducer • Cough Suppressant  
Antihistamine • Nasal Decongestant

**112 DAY COATED CAPLETS**      **56 NIGHT COATED CAPLETS**      **168 TOTAL CAPLETS**

**KIRKLAND Signature**  
**COOL BLAST**  
**SEVERE COLD & FLU PLUS CONGESTION**

**Acetaminophen 325 mg, Dextromethorphan HBr 10 mg, Guaifenesin 200 mg, Phenylephrine HCl 5 mg**  
Pain Reliever • Fever Reducer • Cough Suppressant  
Expectorant • Nasal Decongestant

**Acetaminophen 325 mg, Dextromethorphan HBr 10 mg, Doxylamine succinate 6.25 mg, Phenylephrine HCl 5 mg**  
Pain Reliever • Fever Reducer • Cough Suppressant  
Antihistamine • Nasal Decongestant

**112 DAY COATED CAPLETS**      **56 NIGHT COATED CAPLETS**      **168 TOTAL CAPLETS**

**Relieves:**  
Headache, Fever, Sore Throat, Minor Aches & Pains, Chest Congestion, Thins & Loosens Mucus, Nasal Congestion, Sinus Pressure, Cough

**Relieves:**  
Headache, Fever, Sore Throat, Minor Aches & Pains, Nasal Congestion, Sinus Pressure, Sneezing, Runny Nose, Cough

**NO PRINT / NO VARIATION AREA**  
**LOT NO. / EXP. DATE**

**Drug Facts (continued)**  
Other information: Each caplet contains acetaminophen 325 mg, dextromethorphan HBr 10 mg, guaifenesin 200 mg, and phenylephrine HCl 5 mg. Do not use if you are allergic to any of these ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**Warnings:**  
Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**Directions:**  
Read the Drug Facts carefully before using this product. Use only as directed. Do not exceed the recommended dose. Do not use for more than 10 days unless directed by a doctor. Do not use if symptoms persist or worsen. Do not use if you experience dizziness, lightheadedness, or other symptoms. Do not use if you experience dizziness, lightheadedness, or other symptoms.

**Other information:**  
This product contains acetaminophen, dextromethorphan, guaifenesin, and phenylephrine. Do not use if you are allergic to any of these ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.


**Keep Out of Reach of Children:**  
Always keep this product out of the reach of children. Use proper storage to help prevent access to this product by children.

**Use Only as Directed:**  
Do not use this product if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**Warnings:**  
Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.

**Directions:**  
Read the Drug Facts carefully before using this product. Use only as directed. Do not exceed the recommended dose. Do not use for more than 10 days unless directed by a doctor. Do not use if symptoms persist or worsen. Do not use if you experience dizziness, lightheadedness, or other symptoms. Do not use if you experience dizziness, lightheadedness, or other symptoms.

**Other information:**  
This product contains acetaminophen, dextromethorphan, guaifenesin, and phenylephrine. Do not use if you are allergic to any of these ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease, high blood pressure, or other cardiovascular conditions. Do not use if you are taking any other medications, especially other pain relievers, cough suppressants, or decongestants.



**Kirkland 44-795796**

**SEVERE COLD AND FLU PLUS CONGESTION**  
acetaminophen, dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl kit

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63981-795	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63981-795-81	14 in 1 CARTON	06/30/2023	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination		

**Quantity of Parts**

Part #	Package Quantity	Total Product Quantity
Part 1		8
Part 2		4

**Part 1 of 2****SEVERE COLD AND FLU PLUS CONGESTION DAY**

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

**Product Information**

Item Code (Source)	NDC:63981-995
Route of Administration	ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

**Inactive Ingredients**

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>	MENTHOL	<b>Imprint Code</b>	44;795
<b>Contains</b>			

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/30/2023	

## Part 2 of 2

### SEVERE COLD AND FLU PLUS CONGESTION NIGHT

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl tablet, film coated

## Product Information

<b>Item Code (Source)</b>	NDC:63981-996
<b>Route of Administration</b>	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>FERROSO FERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>FD&amp;C YELLOW NO. 6 ALUMINUM LAKE</b> (UNII: GYP6Z2JR6Q)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

SUCRALOSE (UNII: 96K6UQ3ZD4)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>	MENTHOL	<b>Imprint Code</b>	44;796
<b>Contains</b>			

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/30/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/30/2023	

**Labeler** - COSTCO WHOLESALE CORPORATION (103391843)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63981-795) , pack(63981-795)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63981-795)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(63981-795)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63981-795)

Revised: 4/2025

COSTCO WHOLESALE CORPORATION