



**Drug Facts**

Active ingredients: Iodine Topical Solution (Iodine 7% iodine)  
Antiseptic

**Purpose:**

**WARNINGS:** For external use only. **Product should not be used on pregnant women or infants 4 8 weeks, or anyone with a thyroid condition.**

Do not ingest or apply to food. Do not use in the eyes, for longer than 1 week unless directed by a doctor, on individuals with sensitivity to iodine products, over large areas of the body.

Stop use and consult a doctor if irritation and redness develop, condition persists for more than 72 hours, in case of deep or puncture wounds, in case of animal bites or serious burns.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Inactive ingredients:**  
Polyethylene glycol, sodium iodide

**SAMPLE - NOT FOR SALE**



**HYPODINE**  
IODINE TOPICAL ANTISEPTIC SOLUTION

1 fl oz (30 mL)

FOR HOSPITAL OR PROFESSIONAL USE ONLY

**Use:**

- Healthcare antiseptic for preparation of the skin prior to surgery.
- First aid antiseptic to help prevent infection in minor cuts, scrapes and burns.

**Directions:** Clean the affected area. For preparation of the skin prior to surgery, apply product to the operative site prior to surgery. It is recommended for maximum benefit to apply the product at intervals of approximately 24 hours, 12 hours and 1 hour before surgery.

For first aid antiseptic apply a small amount of this product on the area 1 to 3 times daily. May be covered with a sterile bandage. If banded, let dry first.

**Other information:** Store at room temperature or in cool environment. Protected from freezing. Avoid excessive heat and direct sunlight. Not for consumer use.

NDC: 83465-521-01  
Manufactured by OPURE Corp.  
24 Old County Home Rd, Rahway NJ 07065  
MADE IN THE USA.

8L2P-0524

**Drug Facts**

Active ingredients: Iodine Topical Solution (Iodine 7% iodine)  
Antiseptic

**Purpose:**

**WARNINGS:** For external use only. **Product should not be used on pregnant women or infants 4 8 weeks, or anyone with a thyroid condition.**

Do not ingest or apply to food. Do not use in the eyes, for longer than 1 week unless directed by a doctor, on individuals with sensitivity to iodine products, over large areas of the body.

Stop use and consult a doctor if irritation and redness develop, condition persists for more than 72 hours, in case of deep or puncture wounds, in case of animal bites or serious burns.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Inactive ingredients:**  
Polyethylene glycol, sodium iodide

**SAMPLE - NOT FOR SALE**



**HYPODINE**  
IODINE TOPICAL ANTISEPTIC SOLUTION

1 fl oz (30 mL)

FOR HOSPITAL OR PROFESSIONAL USE ONLY

**Use:**

- Healthcare antiseptic for preparation of the skin prior to surgery.
- First aid antiseptic to help prevent infection in minor cuts, scrapes and burns.

**Directions:** Clean the affected area. For preparation of the skin prior to surgery, apply product to the operative site prior to surgery. It is recommended for maximum benefit to apply the product at intervals of approximately 24 hours, 12 hours and 1 hour before surgery.

For first aid antiseptic apply a small amount of this product on the area 1 to 3 times daily. May be covered with a sterile bandage. If banded, let dry first.

**Other information:** Store at room temperature or in cool environment. Protected from freezing. Avoid excessive heat and direct sunlight. Not for consumer use.

NDC: 83465-521-01  
Manufactured by OPURE Corp.  
24 Old County Home Rd, Rahway NJ 07065  
MADE IN THE USA.

8L2P-0524

**Drug Facts**

Active ingredients: Iodine Topical Solution (Iodine 7% iodine)  
Antiseptic

**Purpose:**

**WARNINGS:** For external use only. **Product should not be used on pregnant women or infants 4 8 weeks, or anyone with a thyroid condition.**

Do not ingest or apply to food. Do not use in the eyes, for longer than 1 week unless directed by a doctor, on individuals with sensitivity to iodine products, over large areas of the body.

Stop use and consult a doctor if irritation and redness develop, condition persists for more than 72 hours, in case of deep or puncture wounds, in case of animal bites or serious burns.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Inactive ingredients:**  
Polyethylene glycol, sodium iodide

**SAMPLE - NOT FOR SALE**



**HYPODINE**  
IODINE TOPICAL ANTISEPTIC SOLUTION

1 fl oz (30 mL)

FOR HOSPITAL OR PROFESSIONAL USE ONLY

**Use:**

- Healthcare antiseptic for preparation of the skin prior to surgery.
- First aid antiseptic to help prevent infection in minor cuts, scrapes and burns.

**Directions:** Clean the affected area. For preparation of the skin prior to surgery, apply product to the operative site prior to surgery. It is recommended for maximum benefit to apply the product at intervals of approximately 24 hours, 12 hours and 1 hour before surgery.

For first aid antiseptic apply a small amount of this product on the area 1 to 3 times daily. May be covered with a sterile bandage. If banded, let dry first.

**Other information:** Store at room temperature or in cool environment. Protected from freezing. Avoid excessive heat and direct sunlight. Not for consumer use.

NDC: 83465-521-01  
Manufactured by OPURE Corp.  
24 Old County Home Rd, Rahway NJ 07065  
MADE IN THE USA.

8L2P-0524

**Drug Facts**

Active ingredients: Iodine Topical Solution (Iodine 7% iodine)  
Antiseptic

**Purpose:**

**WARNINGS:** For external use only. **Product should not be used on pregnant women or infants 4 8 weeks, or anyone with a thyroid condition.**

Do not ingest or apply to food. Do not use in the eyes, for longer than 1 week unless directed by a doctor, on individuals with sensitivity to iodine products, over large areas of the body.

Stop use and consult a doctor if irritation and redness develop, condition persists for more than 72 hours, in case of deep or puncture wounds, in case of animal bites or serious burns.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Inactive ingredients:**  
Polyethylene glycol, sodium iodide

**SAMPLE - NOT FOR SALE**



**HYPODINE**  
IODINE TOPICAL ANTISEPTIC SOLUTION

1 fl oz (30 mL)

FOR HOSPITAL OR PROFESSIONAL USE ONLY

**Use:**

- Healthcare antiseptic for preparation of the skin prior to surgery.
- First aid antiseptic to help prevent infection in minor cuts, scrapes and burns.

**Directions:** Clean the affected area. For preparation of the skin prior to surgery, apply product to the operative site prior to surgery. It is recommended for maximum benefit to apply the product at intervals of approximately 24 hours, 12 hours and 1 hour before surgery.

For first aid antiseptic apply a small amount of this product on the area 1 to 3 times daily. May be covered with a sterile bandage. If banded, let dry first.

**Other information:** Store at room temperature or in cool environment. Protected from freezing. Avoid excessive heat and direct sunlight. Not for consumer use.

NDC: 83465-521-01  
Manufactured by OPURE Corp.  
24 Old County Home Rd, Rahway NJ 07065  
MADE IN THE USA.

8L2P-0524

**Drug Facts**

Active ingredients: Iodine Topical Solution (Iodine 7% iodine)  
Antiseptic

**Purpose:**

**WARNINGS:** For external use only. **Product should not be used on pregnant women or infants 4 8 weeks, or anyone with a thyroid condition.**

Do not ingest or apply to food. Do not use in the eyes, for longer than 1 week unless directed by a doctor, on individuals with sensitivity to iodine products, over large areas of the body.

Stop use and consult a doctor if irritation and redness develop, condition persists for more than 72 hours, in case of deep or puncture wounds, in case of animal bites or serious burns.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Inactive ingredients:**  
Polyethylene glycol, sodium iodide

**SAMPLE - NOT FOR SALE**



**HYPODINE**  
IODINE TOPICAL ANTISEPTIC SOLUTION

1 fl oz (30 mL)

FOR HOSPITAL OR PROFESSIONAL USE ONLY

**Use:**

- Healthcare antiseptic for preparation of the skin prior to surgery.
- First aid antiseptic to help prevent infection in minor cuts, scrapes and burns.

**Directions:** Clean the affected area. For preparation of the skin prior to surgery, apply product to the operative site prior to surgery. It is recommended for maximum benefit to apply the product at intervals of approximately 24 hours, 12 hours and 1 hour before surgery.

For first aid antiseptic apply a small amount of this product on the area 1 to 3 times daily. May be covered with a sterile bandage. If banded, let dry first.

**Other information:** Store at room temperature or in cool environment. Protected from freezing. Avoid excessive heat and direct sunlight. Not for consumer use.

NDC: 83465-521-01  
Manufactured by OPURE Corp.  
24 Old County Home Rd, Rahway NJ 07065  
MADE IN THE USA.

8L2P-0524

# I2PURE HYPODINE

hypodine liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83465-521
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IODINE</b> (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	3 g in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM IODIDE</b> (UNII: F5WR8N145C)	3 g in 30 mL
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	24 g in 30 mL

## Product Characteristics

<b>Color</b>	brown	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83465-521-30	30 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product	05/11/2026	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	05/11/2026	

**Labeler** - I2Pure Corp (117480499)

## Establishment

Name	Address	ID/FEI	Business Operations
I2Pure Corp		145009177	manufacture(83465-521)

Revised: 5/2026

I2Pure Corp