

PENNCHLOR 100G- chlortetracycline calcium granule
Pharmgate Animal Health

Pennchlor® 100G™

Chlortetracycline
 Type A Medicated Article

Active Drug Ingredients: Chlortetracycline as Chlortetracycline Calcium Complex equivalent to 100 grams Chlortetracycline Hydrochloride/lb.

Ingredients: Chlortetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.

For Use in the Manufacture of Medicated Feeds

CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS

Directions For Use

Indications For Use	Use Levels of Chlortetracycline G®	lbs. of Pennchlor100 per ton
Chickens		
For Chickens: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days)	100-200g/ton	1.0-2.0
WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.		
For Chickens: Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days)	200-400g/ton	2.0-4.0
WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.		
For Chickens: Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. (Feed for 5 days)	500g/ton	5.0
WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. WITHDRAW 24 HOURS PRIOR TO SLAUGHTER.		
Turkeys		
For Turkeys: Control of infectious synovitis caused		

by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days)	200g/ton	2.0
WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.		
For Turkeys: Control of hexamitiasis caused by <i>Hexamita meleagrides</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days)	400g/ton	4.0
WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.		
For Turkeys: Turkey poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.	400g/ton	4.0
For Turkeys: Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline. (Feed continuously for 7-14 days)	25 mg/lb body weight/day	
WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.		
Sheep		
For Breeding Sheep: Reducing the incidence of (vibrionic) abortion caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline.	80 mg/head/day	
WARNING: ZERO-DAY WITHDRAWAL PERIOD.		
Swine		
For Swine: Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by <i>Group E Streptococci</i> susceptible to chlortetracycline.	50-100g/ton	0.5-1.0
For Breeding Swine: Control of leptospirosis (reducing the instances of abortions and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. (Feed continuously for not more than 14 days)	400g/ton	4.0
For Swine: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Feed for not more than 14 days)	Feed approximately 400g/t, varying with body weight and feed consumption to provide 10mg/lb body weight/day	
WARNING: ZERO-DAY WITHDRAWAL PERIOD.		
CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR		

**THIS PRODUCT IN PRE-RUMINATING CALVES.
DO NOT USE IN CALVES TO BE PROCESSED
FOR VEAL.**

For Growing Cattle (over 400 lbs.): For the reduction of the incidence of liver abscesses.	70 mg/head/day
WARNING: ZERO-DAY WITHDRAWAL PERIOD.	
For Beef Cattle: For the control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline.	350 mg/head/day
WARNING: ZERO-DAY WITHDRAWAL PERIOD.	
For Beef Cattle (under 700 lbs.): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.	350 mg/head/day
WARNING: ZERO-DAY WITHDRAWAL PERIOD.	
For Beef Cattle (over 700 lbs.): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.	0.5 mg/lb body weight/day
WARNING: ZERO-DAY WITHDRAWAL PERIOD.	
For Calves, Beef, and Nonlactating Dairy Cattle: For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Treat for not more than 5 days)	10 mg/lb body weight/day
WARNING: ZERO-DAY WITHDRAWAL PERIOD.	

Approved by FDA under NADA # 138-935

Pennchlor 100 G[®] is a registered trademark of Pharmgate Inc.

Marketed by:

Pharmgate Animal Health, LLC

14040 Industrial Road

Omaha, Nebraska 68144

TAKE TIME



OBSERVE DIRECTIONS

PRINCIPAL DISPLAY PANEL - 22.68 Kg Bag Label

Pennchlor[®] 100 G[™]

Chlortetracycline

Type A Medicated Article

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a

licensed veterinarian.

See mixing directions, claims, cautions and warnings on back.

Restricted Drug (California) - Use Only as Directed.

Not For Human Use.

Net wt. 50 LB (22.68 Kg)

Pharmgate

ANIMAL HEALTH



Pharmgate
ANIMAL HEALTH

Pennchlor 100 G™

Pennchlor® 100 G™



Chlortetracycline Type A Medicated Article

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

See mixing directions, claims, cautions and warnings on back.
Restricted Drug (California) - Use Only as Directed.
Not For Human Use.

Net wt. 50 LB (22.68 Kg)



Pharmgate
ANIMAL HEALTH

Pennchlor 100 G™

Pharmgate
ANIMAL HEALTH

Pennchlor 100 G™

Chlortetracycline
Type A Medicated Article

Active Drug Ingredient		Chlortetracycline as Chlortetracycline Calcium Complex equivalent to 100 grams Chlortetracycline Hydrochloride (h)	
Ingredients		Chlortetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.	
For Use in the Manufacture of Medicated Feeds			
CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS			
Directions For Use			
Indications For Use	Use Level of Chlortetracycline	No. of Pounds/100^g per ton	Indications For Use
Chickens			
For Chickens: Control of infectious enteritis caused by <i>Mycobacterium avium</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	100-200g/ton	10-20	
For Chickens: Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	200-400g/ton	20-40	
For Chickens: Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. (Feed for 5 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. WITHDRAWAL NEEDED PRIOR TO SLAGGERS.	50g/ton	5.0	
Turkeys			
For Turkeys: Control of infectious enteritis caused by <i>Mycobacterium avium</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	200g/ton	2.0	
For Turkeys: Control of bacterial enteritis caused by <i>Haemophilus influenzae</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	400g/ton	4.0	
For Turkeys: Turkey profits not over 4 weeks of age. Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.	400g/ton	4.0	
For Turkeys: Control of enteric bacterial septicaemia associated with <i>Haemophilus paratyphus</i> enteritis, circumoral enteritis) susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	250g/lb body weight/day		
Sheep			
For Breeding Sheep: Reducing the incidence of (bacterial abortion caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline. WARNING: ZERO-DAY WITHDRAWAL PERIOD.	100g/head/day		
			Swine
			For Swine: Reducing the incidence of central lymphadenitis (swill abscess) caused by Group E <i>Streptococcus</i> susceptible to chlortetracycline.
			For Breeding Swine: Control of leptospirosis (reducing the incidence of abortions and shedding of leptospires) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. (Feed continuously for not more than 14 days)
			For Swine: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella typhimurium</i> and bacterial paratyphoid caused by <i>Haemophilus influenzae</i> susceptible to chlortetracycline. (Feed for not more than 14 days)
			Feed approximately 400g/t. varying with body weight and feed consumption to provide 10mg/lb body weight/day
			CATTLE, BEEF CATTLE AND NON-LACTATING DAIRY CATTLE
			WARNING: A WITHDRAWAL PERIOD IS REQUIRED. DO NOT BREED OR MILK FOR 28 DAYS. PRODUCT IS PRE-CONSUMPTION CATTLE. DO NOT USE IN CATTLE TO BE PRE-CARRIED FOR VEAL.
			For Growing Cattle (over 400 lbs.): For the reduction of the incidence of liver abscesses.
			WARNING: ZERO-DAY WITHDRAWAL PERIOD.
			For Beef Cattle: For the control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline.
			WARNING: ZERO-DAY WITHDRAWAL PERIOD.
			For Beef Cattle (over 700 lbs.): Control of active infection of <i>Streptococcus</i> caused by <i>Streptococcus</i> susceptible to chlortetracycline.
			WARNING: ZERO-DAY WITHDRAWAL PERIOD.
			For Beef Cattle (over 700 lbs.): Control of active infection of <i>Streptococcus</i> caused by <i>Streptococcus</i> susceptible to chlortetracycline.
			WARNING: ZERO-DAY WITHDRAWAL PERIOD.
			For Calves, Beef, and Nonlactating Dairy Cattle: For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial paratyphoid caused by <i>Haemophilus influenzae</i> susceptible to chlortetracycline. (Treat for not more than 5 days)
			WARNING: ZERO-DAY WITHDRAWAL PERIOD.

Approved by FDA under NADA # 138-935
Pennchlor 100 G[®] is a registered trademark of Pharmgate Inc.



Rev: 05-22
Bag ZG



Marketed by:
Pharmgate Animal Health, LLC
14040 Industrial Road
Omaha, Nebraska 68144



VFD-1

PENNCHLOR 100G

chlortetracycline calcium granule

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:51429-101
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Chlortetracycline Calcium (UNII: NR4B2SX17S) (Chlortetracycline - UNII:WCK1KIQ23Q)	Chlortetracycline Hydrochloride	220 g in 1 kg

Inactive Ingredients

Ingredient Name	Strength
Calcium Carbonate (UNII: H0G9379FGK)	
Mineral Oil (UNII: T5L8T28FGP)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51429-101-01	22.68 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA138935	01/08/2007	

Labeler - Pharmgate Animal Health (833270817)

Registrant - Pharmgate Inc. (079628671)

Revised: 2/2023

Pharmgate Animal Health