

LIDOCAINE PAIN RELIEF PATCH- lidocaine patch
GERI-CARE PHARMACEUTICAL CORP

GC 163 Lidocaine Pain Relief Patch

Active Ingredients

Lidocaine 4.0%

Purpose

Topical Anesthetic

Use

Temporarily relieves minor pain.

Warnings

For external use only.

Do not use ■ More than one patch on your body at a time ■ On cut, irritated or swollen skin

■ On puncture wounds ■ For more than one week without consulting a doctor

■ If you are allergic to any active or inactive ingredients

When using this product ■ Use only as directed

■ Read and follow all directions and warnings on this carton ■ Do not allow contact with the eyes

■ Do not use at the same time as other topical analgesics ■ Do not microwave

■ Do not bandage tightly or apply local heat (such as heating pads) to the area of use

■ Dispose of used patch in manner that always keeps product away from children and pets. Used

patches still contain the drug product that can produce serious adverse effects if a child or pet

chews or ingests this patch.

Stop use and consult a doctor if ■ Condition worsens ■ Redness is present ■

Irritation develops

■ Symptoms persist for more than 7 days or clear up and occur again within a few days

■ You

experience signs of skin injury, such as pain, swelling or blistering where the product was applied.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children and pets.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222)

right away.

Directions

Adults and children 12 years of age and over: Clean and dry affected area.

Carefully remove

backing from patch (see illustration). Apply sticky side of patch to affected area. Use one patch

for up to 12 hours. Discard patch after single use.

Children under 12 years of age: consult a physician.

Other Information

■ Store at room temperature ■ Avoid direct sunlight ■ Protect from excessive moisture

Inactive Ingredients

aluminum glycinate, benzyl alcohol, butylated hydroxyanisole, carboxymethylcellulose sodium,

edetate disodium, ethylhexylglycerin, glycerin, kaolin, methacrylate copolymer, oleic acid, phenoxyethanol, polyacrylic acid, polysorbate 80, polyvinyl alcohol, propylene glycol, PVP, sodium

polyacrylate, sorbitol, tartaric acid, titanium dioxide, water

Questions or comments:

Call 1-800-540-3765

GERI CARE®

CHILD-RESISTANT PACKAGING
NDC 57896-163-05

Compare to the active ingredient in
Aspercreme® Max Strength Lidocaine Patch.**

LIDOCAINE PAIN RELIEF PATCH

Lidocaine 4% Topical Anesthetic

Temporarily relieves pain

- Stay-put flexible patch
- Easy to apply and remove
- No-mess, single-use application
- Odor Free



5 Patches
3.93 X 5.51 IN (10cm X 14cm)

MAXIMUM STRENGTH*

*Lidocaine 4% is the maximum concentration available without a prescription.

GERI CARE®
LIDOCAINE PAIN RELIEF PATCH
Lidocaine 4% • Topical Anesthetic



Tamper Evident: Do not use if pouch is damaged or opened.

Drug Facts	
Active ingredient Lidocaine 4.0%	Purpose Topical anesthetic
Use Temporarily relieves minor pain.	
Warnings	
For external use only	
Do not use ■ More than one patch on your body at a time ■ On cut, irritated or swollen skin	
■ On puncture wounds ■ For more than one week without consulting a doctor	
■ If you are allergic to any active or inactive ingredients	
When using this product ■ Use only as directed	
■ Read and follow all directions and warnings on this carton ■ Do not allow contact with the eyes	
■ Do not use at the same time as other topical analgesics ■ Do not microwave	
■ Do not bandage tightly or apply local heat (such as heating pads) to the area of use	
■ Dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.	
Stop use and consult a doctor if ■ Condition worsens ■ Redness is present ■ Irritation develops	
■ Symptoms persist for more than 7 days or clear up and occur again within a few days ■ You experience signs of skin injury, such as pain, swelling or blistering where the product was applied.	
If pregnant or breastfeeding, ask a health professional before use.	
Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center 1-800-222-1222 right away.	
Directions	
Adults and children 12 years of age and over: Clean and dry affected area. Carefully remove backing from patch (see illustration). Apply sticky side of patch to affected area. Use one patch for up to 12 hours. Discard patch after single use.	
Children under 12 years of age: consult a physician.	
Other information	
■ Store at room temperature ■ Avoid direct sunlight ■ Protect from excessive moisture	
Inactive ingredients	
aluminum glycinate, benzyl alcohol, butylated hydroxyanisole, carboxymethylcellulose sodium, edetate disodium, ethylhexylglycerin, glycerin, kaolin, methacrylate copolymer, oleic acid, phenoxethanol, polyacrylic acid, polysorbate 80, polyvinyl alcohol, propylene glycol, PVP, sodium polyacrylate, sorbitol, tartaric acid, titanium dioxide, water	
Questions or comments: Call 1-800-540-3765	

**This product is not manufactured or distributed by the owner of the registered trademark Aspercreme®.
Distributed by:
GERI-CARE PHARMACEUTICALS CORP.
1295 Towbin Ave
Lakewood, NJ 08701
Made in India

REV 0825



LIDOCAINE PAIN RELIEF PATCH

lidocaine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-163
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 mg in 100 mg

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM POLYACRYLATE (8000 MW) (UNII: 285CYO341L)	
TARTARIC ACID (UNII: W4888I19H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	
DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	

POLYACRYLIC ACID (8000 MW) (UNII: 73861X4K5F)
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
SORBITOL (UNII: 506T60A25R)
BENZYL ALCOHOL (UNII: LKG8494WBH)
KAOLIN (UNII: 24H4NWX5CO)
OLEIC ACID (UNII: 2UMI9U37CP)
POLYSORBATE 80 (UNII: 6OZP39ZG8H)
2-ETHYLHEXYL ACRYLATE, METHACRYLATE, METHYL METHACRYLATE, OR BUTYL METHACRYLATE/HYDROXYPROPYL DIMETHICONE COPOLYMER (30000-300000 MW) (UNII: S7ZA3CCJ4M)
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)

Product Characteristics

Color		Score	
Shape	RECTANGLE	Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-163-05	5 in 1 CARTON	05/01/2026	
1	NDC:57896-163-01	350 mg in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	05/01/2026	

Labeler - GERI-CARE PHARMACEUTICAL CORP (611196254)

Registrant - GERI-CARE PHARMACEUTICAL CORP (611196254)

Revised: 5/2026

GERI-CARE PHARMACEUTICAL CORP