

**EASY EVERYTHING SUNSCREEN- zinc oxide cream**  
**VLT Ventures LLC**

-----  
Zinc Oxide, 16%

Sunscreen

Helps Prevent Sunburn. If used as directed with other sun protection measures (see "Directions"), decreases the risk of skin cancer and early skin aging caused by the sun.

For external use only. Do not use on broken or damaged skin. Stop use and ask a doctor if rash occurs. When using this product, keep out of eyes. Rinse with water to remove. Keep out of reach of children. If swallowed, get medical help or contact a poison control center right away.

Keep out of reach of children

Apply liberally 15 minutes before sun exposure. Use a water-resistant product if swimming or sweating. Reapply at least every 2 hours. Children under 6 months: Ask a doctor. Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad-Spectrum SPF value of 15 or higher and other sun protection measures including: Limit time in the sun, especially from 10 a.m. - 2 p.m. Wear long-sleeved shirts, pants, hats and sunglasses.

Capric/Caprylic Triglyceride, Ceramide 3, Cyclohexasiloxane, Cyclopentasiloxane, Dimethicone, Dimethicone Crosspolymer, Dimethicone/Vinyl Dimethicone Crosspolymer, Dimethiconol, Iron Oxide, Hydrogen Dimethicone, PEG-10 Dimethicone, Polyhydroxystearic Acid, Tetrahexyldecyl Ascorbate, Tocopheryl Acetate, Vinyl Dimethicone/Methicone Silsesquioxane Crosspolymer

- Protect this product from excessive heat and direct sun
- May stain some fabrics

## DRUG FACTS

**ACTIVE INGREDIENT:**  
Zinc Oxide 16%

**PURPOSE:**  
Sunscreen

### USES:

Helps prevent sunburn. If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early aging caused by the sun.

### WARNINGS:

For external use only. Do not use on damaged or broken skin. Stop use and ask a doctor if rash occurs. When using this product, keep out of eyes. Rinse with water to remove. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

### DIRECTIONS:

- Apply liberally 15 minutes before sun exposure
- Use a water resistant product if swimming or sweating.
- Reapply at least every 2 hours
- Children under 6 months: Ask a doctor
- **Sun Protection Measures:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including:
  - Limit time in the sun, especially from 10 am - 2 pm
  - Wear long-sleeved shirts, pants, hats, and sunglasses

### INACTIVE INGREDIENTS:

Capric/Caprylic Triglyceride, Ceramide 3, Cyclohexasiloxane, Cyclopentasiloxane, Dimethicone, Dimethicone Crosspolymer, Dimethicone/Vinyl Dimethicone Crosspolymer, Dimethiconol, Hydrogen Dimethicone, Iron Oxide, PEG-10 Dimethicone, Polyhydroxystearic Acid, Tetrahexyldecyl Ascorbate, Tocopheryl Acetate, Vinyl Dimethicone/Methicone Silsesquioxane Crosspolymer

### OTHER INFORMATION:

- Protect this product from excessive heat and direct sun
- May stain some fabrics.



MANUFACTURED FOR:  
Cinchy | Dallas, TX 75208  
www.cinchysun.com



# cinchy

Easy Everything Sunscreen

SPF 40

Lightly Tinted for All Skin Tones  
Broad Spectrum  
Daily Facial Mineral Sunscreen  
Silky Moisturizing Ceramides  
Antioxidants C & E

Net Wt. 1.75 oz / 50 g

## EASY EVERYTHING SUNSCREEN

zinc oxide cream

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:87670-208(NDC:58892-208)
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	160 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
CAPRYLIC/CAPRIC TRIGLYCERIDE (UNII: C9H2L21V7U)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIMETHICONE CROSSPOLYMER (UNII: UF7620L1W6)	
ALPHA-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	

<b>DIMETHICONOL (2000 CST)</b> (UNII: T74O12AN6Y)
<b>CI 77499</b> (UNII: XM0M87F357)
<b>PEG-10 DIMETHICONE (600 CST)</b> (UNII: 8PR7V1SVM0)
<b>CYCLOHEXASILOXANE</b> (UNII: XHK3U310BA)
<b>HYDROGEN DIMETHICONE (20 CST)</b> (UNII: 12Z59IF64N)
<b>POLYHYDROXYSTEARIC ACID (2300 MW)</b> (UNII: YXH47AOU0F)
<b>DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE)</b> (UNII: 9E4CO0W6C5)
<b>CYCLOPENTASILOXANE</b> (UNII: 0THT5PCI0R)
<b>TETRAHEXYLDECYL ASCORBATE</b> (UNII: 9LBV3F07AZ)
<b>CI 77492</b> (UNII: EX438O2MRT)
<b>CI 77491</b> (UNII: 1K09F3G675)
<b>VINYL DIMETHICONE/METHICONE SILSESQUIOXANE CROSSPOLYMER</b> (UNII: 9NH1UDD2RR)
<b>CERAMIDE 3</b> (UNII: 4370DF050B)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87670-208-50	50 g in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	07/10/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	07/10/2024	

### Labeler - VLT Ventures LLC (136296328)

### Establishment

Name	Address	ID/FEI	Business Operations
Fragrance Manufacturing Inc.		793406000	manufacture(87670-208)

### Establishment

Name	Address	ID/FEI	Business Operations
Custom Analytics LLC		144949372	analysis(87670-208)

### Establishment

Name	Address	ID/FEI	Business Operations
Fallien Cosmeceuticals Ltd.		958388357	relabel(87670-208)