

**FLURISO - isoflurane liquid**  
**MWI Veterinary Supply Co**


-----  
**CAUTION**

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**WARNING: Not for use in horses intended for food**

**DESCRIPTION**

Isoflurane, USP is a nonflammable, nonexplosive general inhalation anesthetic agent. Its chemical name is 1-chloro-2,2,2-trifluoroethyl difluoromethyl ether, and its structural formula is:



Each mL contains 99.9% isoflurane.

Some physical constants are:

Molecular weight	184.5	
Boiling point at 760 mm Hg	48.5° C (uncorr.)	
Refractive index $n_D^{20}$	1.2990-1.3005	
Specific gravity 25°/25° C	1.496	
Vapor pressure in mm Hg**	20° C	238
	25° C	295
	30° C	367
	35° C	450

\*\*Equation for vapor pressure calculation:

$$\log_{10} P_{vap} = A + B/T$$

where: A = 8.056

B = -1664.58

$$T = ^\circ\text{C} + 273.16 \text{ (Kelvin)}$$

Partition coefficients at 37° C:

Water/gas	0.61
Blood/gas	1.43
Oil/gas	90.8

Partition coefficients at 37° C:		
	Water/gas	0.61
	Blood/gas	1.43
	Oil/gas	90.8

Partition coefficients at 25° C - rubber and plastic

Conductive rubber/gas	62.0
Butyl rubber/gas	75.0
Polyvinyl chloride/gas	110.0
Polyethylene/gas	~2.0
Polyurethane/gas	~1.4
Polyolefin/gas	~1.1
Butyl acetate/gas	~2.5

Purity by gas chromatography >99.9%

Lower limit of flammability in oxygen or nitrous oxide at 9 joules/sec. and 23° C None

Lower limit of flammability in oxygen or nitrous oxide at 900 joules/sec. and 23° C Greater than useful concentration in anesthesia

MAC (Minimum Alveolar Concentration) is 1.31% in horses<sup>1</sup> and 1.28% in dogs.<sup>6</sup>

Isoflurane is a clear, colorless, stable liquid containing no additives or chemical stabilizers. Isoflurane has a mildly pungent, musty, ethereal odor. Samples stored in indirect sunlight in clear, colorless glass for five years, as well as samples directly exposed for 30 hours to a 2 amp, 115 volt, 60 cycle long wave U.V. light were unchanged in composition as determined by gas chromatography. Isoflurane in one normal sodium methoxide-methanol solution, a strong base, for over six months consumed essentially no alkali, indicative of strong base stability. Isoflurane does not decompose in the presence of soda lime (at normal operating temperatures), and does not attack aluminum, tin, brass, iron or copper.

## CLINICAL PHARMACOLOGY

Isoflurane, USP is an inhalation anesthetic. Induction and recovery from anesthesia with isoflurane are rapid.<sup>2,5</sup> The level of anesthesia may be changed rapidly with isoflurane. Isoflurane is a profound respiratory depressant. RESPIRATION MUST BE MONITORED CLOSELY IN THE HORSE AND

DOG AND SUPPORTED WHEN NECESSARY. As anesthetic dose is increased, both tidal volume and respiratory rate decrease.<sup>3,6</sup> This depression is partially reversed by surgical stimulation, even at deeper levels of anesthesia.

Blood pressure decreases with induction of anesthesia but returns toward normal with surgical stimulation. Progressive increases in depth of anesthesia produce corresponding decreases in blood pressure; however, heart rhythm is stable and cardiac output is maintained with controlled ventilation and normal PaCO<sub>2</sub> despite increasing depth of anesthesia. The hypercapnia which attends spontaneous ventilation during isoflurane anesthesia increases heart rate and raises cardiac output above levels observed with controlled ventilation.<sup>3</sup> Isoflurane does not sensitize the myocardium to exogenously administered epinephrine in the dog.

Muscle relaxation may be adequate for intra-abdominal operations at normal levels of anesthesia. However, if muscle relaxants are used to achieve greater relaxation, it should be noted that: **ALL COMMONLY USED MUSCLE RELAXANTS ARE MARKEDLY POTENTIATED WITH ISOFLURANE, THE EFFECT BEING MOST PROFOUND WITH THE NONDEPOLARIZING TYPE.** Neostigmine reverses the effect of nondepolarizing muscle relaxants in the presence of isoflurane but does not reverse the direct neuromuscular depression of isoflurane.

## **INDICATIONS & USAGE**

Isoflurane, USP is used for induction and maintenance of general anesthesia in horses and dogs.

## **CONTRAINDICATIONS**

Isoflurane, USP is contraindicated in horses and dogs with known sensitivity to isoflurane or to other halogenated agents.

## **WARNINGS**

Increasing depth of anesthesia with isoflurane may increase hypotension and respiratory depression. The electroencephalographic pattern associated with deep anesthesia is characterized by burst suppression, spiking, and isoelectric periods.<sup>4</sup>

Since levels of anesthesia may be altered easily and rapidly, only vaporizers producing predictable percentage concentrations of isoflurane should be used. (See **DOSAGE AND ADMINISTRATION**).

The action of nondepolarizing relaxants is augmented by isoflurane. Less than the usual amounts of these drugs should be used. If the usual amounts of nondepolarizing relaxants are given, the time for recovery from myoneural blockade will be longer in the presence of isoflurane than in the presence of other commonly used anesthetics.

Not for use in horses intended for food.

Keep out of reach of children.

## **PRECAUTIONS**

Isoflurane, USP like other inhalational anesthetics, can react with desiccated carbon dioxide (CO<sub>2</sub>) absorbents to produce carbon monoxide which may result in elevated carboxyhemoglobin levels in some patients. Case reports suggest that barium hydroxide lime and soda lime become desiccated when

fresh gases are passed through the CO<sub>2</sub> absorber canister at high flow rates over many hours or days. When a clinician suspects that CO<sub>2</sub> absorbent may be desiccated, it should be replaced before the administration of isoflurane.

## **PREGNANCY**

**Usage in Pregnancy:** Reproduction studies have been performed in mice and rats with no evidence of fetal malformation attributable to isoflurane. Adequate data concerning the safe use of isoflurane in pregnant and breeding horses and dogs have not been obtained.

## **ADVERSE REACTIONS**

Hypotension, respiratory depression, and arrhythmias have been reported.

## **OVERDOSAGE**

In the event of overdosage, or what may appear to be overdosage, the following action should be taken: Stop drug administration, establish that the airway is clear and initiate assisted or controlled ventilation with pure oxygen as circumstances dictate.

## **DOSAGE & ADMINISTRATION**

**Caution: Operating rooms should be provided with adequate ventilation to prevent the accumulation of anesthetic vapors.**

**Premedication:** A premedication regimen, which may be employed depending upon the patient status, to avert excitement during induction, might include an anticholinergic, a tranquilizer, a muscle relaxant, and a short-acting barbiturate.

**Inspired Concentration:** The delivered concentration of Isoflurane, USP should be known. Isoflurane may be vaporized using a flow-through vaporizer specifically calibrated for isoflurane. Vaporizers delivering a saturated vapor which is then diluted (e.g. Vernitrol<sup>®</sup> vaporizer) also may be used. The delivered concentration from such a vaporizer may be calculated using the formula:

$$\% \text{ Isoflurane} = \frac{100 P_V F_V}{F_T (P_A - P_V)}$$

where:  $P_A$  = Pressure of atmosphere

$P_V$  = Vapor pressure of isoflurane

$F_V$  = Flow of gas through vaporizer (mL/min)

$F_T$  = Total gas flow (mL/min)

Isoflurane contains no stabilizer. Nothing in the drug product alters calibration or operation of these vaporizers.

## **Induction:**

**Horses:** Inspired concentrations of 3.0% to 5.0% isoflurane alone with oxygen following a barbiturate anesthetic induction are usually employed to induce surgical anesthesia in the horse.

**Dogs:** Inspired concentrations of 2.0% to 2.5% isoflurane alone with oxygen following a barbiturate anesthetic induction are usually employed to induce surgical anesthesia in the dog.

These concentrations can be expected to produce surgical anesthesia in 5 to 10 minutes.

**Maintenance:**

The concentration of vapor necessary to maintain anesthesia is much less than that required to induce it.

**Horses:** Surgical levels of anesthesia in the horse may be sustained with a 1.5% to 1.8% concentration of isoflurane in oxygen.

**Dogs:** Surgical levels of anesthesia in the dog may be sustained with a 1.5% to 1.8% concentration of isoflurane in oxygen.

The level of blood pressure during maintenance is an inverse function of isoflurane concentration in the absence of other complicating problems. Excessive decreases, unless related to hypovolemia, may be due to depth of anesthesia and in such instances may be corrected by lightening the level of anesthesia. Recovery from isoflurane anesthesia is typically uneventful.<sup>2</sup>

**HOW SUPPLIED**

Isoflurane, USP is packaged in 100 mL (NDC 13985-528-40) and 250 mL (NDC 13985-528-60) amber-colored bottles.

**Storage**

**Storage:**

Store at room temperature 20° to 25°C (68° to 77° F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

**REFERENCES**

1. Steffey EP, Howland D Jr, Giri S, Eger El II. Enflurane, Halothane and Isoflurane Potency in Horses. *AM J Vet Res.* 1977; 38(7): 1037-1039.
2. Auer JA, Garner HE, Amend JF, Hutcheson DP, Salem CA. Recovery from Anesthesia in Ponies: A Comparative Study of the Effects of Isoflurane, Enflurane, Methoxyflurane and Halothane. *Equine Vet J.* 1978; 10(1): 18-23.
3. Steffey EP, Howland D Jr. Comparison of Circulatory and Respiratory Effects of Isoflurane and Halothane Anesthesia in Horses. *Am J Vet Res.* 1980; 41(5): 821-825.
4. Auer JA, Amend JF, Garner HE, Hutcheson DP, Salem CA. Electroencephalographic Responses During Volatile Anesthesia in Domestic Ponies: A Comparative Study of Isoflurane, Enflurane, Methoxyflurane, and Halothane. *Equine Practice.* 1979;3:130-134.
5. Klide AM. Cardiopulmonary Effects of Enflurane and Isoflurane in the Dog. *Am J Vet Res.* 1976; 37(2): 127-131.
6. Steffey EP, Howland D. Isoflurane Potency in the Dog and Cat. *Am J Vet Res.* 1977; 38(11): 1833-1936.

**Distributed By**

Distributed By: MWI  
Boise, ID 83705

(888) 694-8381  
www.vetone.net

## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Label 100 mL

NDC 13985-528-40

**100 mL**

**VET one**<sup>®</sup>

**Fluriso**<sup>™</sup>

(Isoflurane, USP)

**Liquid for Inhalation**

**ANESTHETIC**  
Nonflammable, Nonexplosive

For veterinary use in horses and dogs.

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

ANADA 200-237, Approved by FDA

**V1 501017** Net Contents:100 mL

3  
13985-00931  
0

**For Animal Use Only.**

**WARNING:** Not for use in horses intended for food.

**IMPORTANT:** See accompanying product information for dosage and directions for use.

**WARNING:** Keep out of reach of children.

**CAUTION:** Operating rooms should be provided with adequate ventilation to prevent the accumulation of anesthetic vapors.

Each mL contains 99.9% isoflurane.

**Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F)**  
**[See USP Controlled Room Temperature].**

Distributed by: MWI  
Boise, ID 83705  
(888) 694-8381  
www.vetone.net

TAKE TIME  OBSERVE LABEL DIRECTIONS

Code : AP/DRUGS/220/96 Rev.03/12 Product of India

To report adverse reactions or obtain MSDS Call (888) 822-8431 DNI 22135/1

Label 250 mL

Label 250 mL

NDC 13985-528-60



250 mL

**Fluriso**<sup>TM</sup>  
(Isoflurane, USP)

**Liquid for Inhalation**

**ANESTHETIC**

Nonflammable, Nonexplosive

For veterinary use in horses and dogs.

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

ANADA 200-237, Approved by FDA

V1 502017

Net Contents: 250 mL



**For Animal Use Only.**

**WARNING:** Not for use in horses intended for food.

**IMPORTANT:** See accompanying product information for dosage and directions for use.

**WARNING:** Keep out of reach of children.

**CAUTION:** Operating rooms should be provided with adequate ventilation to prevent the accumulation of anesthetic vapors.

Each mL contains 99.9% isoflurane.

**Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].**

Distributed by: MWI  
Boise, ID 83705  
(888) 694-8381  
www.vetone.net

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

Code : AP/DRUGS/220/96 Rev.03/12

Product of India

To report adverse reactions or obtain MSDS  
Call (888) 822-8431

DNI 22136/1

**FLURISO**

iso flurane liquid

**Product Information**

<b>Product Type</b>	PREScription ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:13985-528
<b>Route of Administration</b>	RESPIRATORY (INHALATION)		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ISO FLURANE (UNII: CYS9 AKD70 P) (ISOFLURANE - UNII:CYS9 AKD70 P)	ISOFLURANE	1 mL in 1 mL

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-528-40	100 mL in 1 BOTTLE, GLASS		
2	NDC:13985-528-60	250 mL in 1 BOTTLE, GLASS		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200237	02/27/2014	

**Labeler** - MWI Veterinary Supply Co (019926120)

**Registrant** - Piramal Enterprises Limited (919067108)

