

**SHISEIDO ULTIMATE SUN PROTECTOR- avobenzone, homosalate, octisalate, and octocrylene spray**  
**SHISEIDO AMERICAS CORPORATION**

---

**SHISEIDO ULTIMATE SUN PROTECTOR SPRAY**

**Drug Facts**

<b>Active ingredients</b>	<b>Purpose</b>
AVOBENZONE 2.3%	Sunscreen
HOMOSALATE 10.0%	Sunscreen
OCTISALATE 5.0%	Sunscreen
OCTOCRYLENE 5.0%	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

**For external use only**

**Do not use** on damaged or broken skin

**When using this product** keep out of eyes. Rinse with water to remove.

**Stop use and ask a doctor** if rash occurs.

**Keep out of reach of children.** If product is swallowed, get medical help or contact a Poison Control Center right away.

**Flammable**

Do not use near heat, flame, or while smoking

**Directions**

For sunscreen use:

- spray liberally and spread evenly by hand 15 minutes before sun exposure
- hold container 4 to 6 inches from the skin to apply.
- do not spray directly into face. Spray on hands then apply to face.
- do not apply in windy conditions.
- use in a well-ventilated area.
- reapply:
  - after 80 minutes of swimming or sweating
  - immediately after towel drying.
  - at least every 2 hours

- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

## **Inactive Ingredients**

ALCOHOL DENAT. • DIISOPROPYL SEBACATE • TRIETHYLHEXANOIN • C12-15 ALKYL BENZOATE • DIMETHICONE • CAPRYLYL METHICONE • SILICA DIMETHYL SILYLATE • DIPROPYLENE GLYCOL • PPG-3 DIPIVALATE • VITIS VINIFERA (GRAPE) SEED OIL • PEG/PPG-14/7 DIMETHYL ETHER • LECITHIN • GLYCRRHIZA GLABRA (LICORICE) ROOT EXTRACT • PPG-17 • HYDROXYSTEARIC ACID • POLYHYDROXYSTEARIC ACID • POLYAMIDE-8 • PALMITIC ACID • LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE • BHT • TOCOPHEROL • PENTAERYTHRITYL TETRA-DI-T-BUTYL HYDROXYHYDROCINNAMATE • PHENOXYETHANOL • FRAGRANCE •

## **Other information**

- protect this product in this container from excessive heat and direct sun.

## **Questions or comments?**

Call toll free 1-800-906-7503

## **PRINCIPAL DISPLAY PANEL - 150 mL Cartridge Carton**

SHISEIDO  
GINZA TOKYO

Ultimate Sun Protector  
Spray

SPF  
40

Invisible and breathable  
protection veil.

BROAD SPECTRUM  
SPF 40

WATER RESISTANT  
(80 MINUTES)

SUNSCREEN

150mL 5 FL. OZ.



**SHISEIDO ULTIMATE SUN PROTECTOR**

avobenzone, homosalate, octisalate, and octocrylene spray

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-891
Route of Administration	TOPICAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>AVOBENZONE</b> (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	3.1533 g in 150 mL
<b>HOMOSALATE</b> (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	13.71 g in 150 mL
<b>OCTISALATE</b> (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	6.855 g in 150 mL
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	6.855 g in 150 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>DIISOPROPYL SEBACATE</b> (UNII: J8T3X564IH)	
<b>TRIETHYLHEXANOIN</b> (UNII: 7K3W1BIU6K)	
<b>ALKYL (C12-15) BENZOATE</b> (UNII: A9EJ3J61HQ)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>CAPRYLYL TRISILOXANE</b> (UNII: Q95M2P1KJL)	
<b>SILICA DIMETHYL SILYLATE</b> (UNII: EU2PSP0G0W)	
<b>DIPROPYLENE GLYCOL</b> (UNII: E107L85C40)	
<b>GRAPE SEED OIL</b> (UNII: 930MLC8XGG)	
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>GLYCYRRHIZA GLABRA</b> (UNII: 2788Z9758H)	
<b>PPG-17</b> (UNII: OV0Q322E0U)	
<b>12-HYDROXYSTEARIC ACID</b> (UNII: 933ANU3H2S)	
<b>POLYHYDROXYSTEARIC ACID (2300 MW)</b> (UNII: YXH47AOU0F)	
<b>POLYAMIDE-8 (4500 MW)</b> (UNII: 77723GV81A)	
<b>PALMITIC ACID</b> (UNII: 2V16E095H1)	
<b>LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE</b> (UNII: 25G622K2RA)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)	
<b>PENTAERYTHRITOL TETRAKIS(3-(3,5-DI-TERT-BUTYL-4-HYDROXYPHENYL)PROPIONATE)</b> (UNII: 255PIF62MS)	
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-891-20	150 mL in 1 CARTRIDGE; Type 0: Not a Combination Product	03/01/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	03/01/2024	

## Labeler - SHISEIDO AMERICAS CORPORATION (193691821)

## Establishment

Name	Address	ID/FEI	Business Operations
Shiseido America Inc.		782677132	analysis(58411-891) , manufacture(58411-891)

Revised: 1/2026

SHISEIDO AMERICAS CORPORATION