

WALGREENS HEADACHE RELIEF EXTRA STRENGTH- acetaminophen 250mg, aspirin 250mg and caffeine 65mg tablet, coated
Walgreens

Walgreens Headache Relief Extra Strength GELTAB

DRUG FACTS

DRUG FACTS

Active Ingredient (in each softgel)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever

Pain reliever aid

Uses

Temporarily relieves minor aches and pains due to:

- headache ■ a cold ■ arthritis ■ muscular aches
- toothache ■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing)

■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 8 gels in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat.

Do not use

■ if you have ever had an allergic reaction to

acetaminophen, aspirin or any other pain reliever/fever reducer

■ with any other drug containing acetaminophen (prescription or nonprescription). if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

■ you have liver disease ■ stomach

bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking

Ask a doctor or pharmacist before use if you are taking

■ a prescription drug for diabetes, gout, or arthritis

■ any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding:

■ feel faint ■ vomit blood ■ have bloody or black stools

■ have stomach pain that does not get better

■ ringing in the ears or loss of hearing occurs

■ painful area is red or swollen

■ pain gets worse or lasts for more than 10 days

■ fever gets worse or lasts for more than 3 days

■ any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before

use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a

doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

■ do not use more than directed (see overdose warning)

■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 2 gels every 6 hours; not more than 8 gels in 24 hours

■ children under 12 years: ask a doctor

Other information

■ store at room temperature between 20-25°C (68-77°F)

■ avoid excessive heat, cold and humidity

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, edible black ink, ethyl cellulose, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hypromellose, isopropyl alcohol, lactose monohydrate, light mineral oil, microcrystalline cellulose, pregelatinized

potato starch, purified water, pvpk-30, sorbitol sorbitan solution, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?

1-888-333-9792 Monday-Friday 8am-4pm EST



Walgreens

Compare to the active ingredients in Excedrin® Extra Strength*

Headache Relief

ACETAMINOPHEN 250 mg / PAIN RELIEVER
ASPIRIN (NSAID) 250 mg / PAIN RELIEVER
CAFFEINE 65 mg / PAIN RELIEVER AID

Extra Strength

120

GELTABS

ACTUAL SIZE

Drug Facts

Active ingredients (in each geltab)
Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID) Pain reliever
Caffeine 65 mg Pain reliever aid
*Nonsteroidal anti-inflammatory drug

Uses
Temporarily relieves minor aches and pains due to:
Headache Cold Flu/Influenza Muscular aches
Nervousness and menstrual cramps

Warnings
Keep's symptoms. Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Always alert: Acetaminophen may cause skin reactions. Symptoms may include: skin redness, blisters, or rash. If a skin reaction occurs, stop use and seek medical help right away. Always alert: Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, or asthma (wheezing). Always alert: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 gels in 24 hours, which is the maximum daily amount. Do not take more than 8 gels in 24 hours, which is the maximum daily amount. Do not take more than 8 gels in 24 hours, which is the maximum daily amount. Do not take more than 8 gels in 24 hours, which is the maximum daily amount.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 65 or older, have had stomach ulcers or bleeding problems, take a blood thinner (warfarin) or steroid drug, take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

Child use warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer with any other drug containing acetaminophen (prescription or over-the-counter), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if:
You have liver disease
Stomach bleeding warning applies to you
You have a history of stomach problems such as heartburn
You have high blood pressure, heart disease, liver problems, or kidney disease
You are taking a diuretic
You have asthma
Ask a doctor or pharmacist before use if you are taking:
A prescription drug for diabetes, gout, or arthritis
Any other drug, or are under a doctor's care for any serious condition
Stay on and seek advice if:
An allergic reaction occurs. Seek medical help right away.
You experience any of the following signs of stomach bleeding:
Low fever, vomit blood, or have bloody or black stools
Have stomach pain that does not get better
Ring in the ears or loss of hearing occurs
Get itchy or itchy or swollen
Pain gets worse or lasts for more than 3 days
Liver gets worse or lasts for more than 3 days
Any new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless a health professional directs you to do so because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions
Do not use more than directed (see overdose warning)
Drink a full glass of water with each dose
Adults and children 12 years and over: take 2 gels every 6 hours, not more than 8 gels in 24 hours
Children under 12 years: ask a doctor

Drug Facts (continued)

Other information
Store at room temperature between 20-25°C (68-77°F)
Avoid excessive heat, cold and humidity

Inactive ingredients
corn starch, DSC yellow #10 aluminum lake, edible black ink, erythritol, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerol, hypromellose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose, light mineral oil, microcrystalline cellulose, polyethylene glycol, polyethylene glycol, purified water, pvpk-30, sorbitol, sorbitan, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?
1-888-333-9792 Monday-Friday 8am-4pm EST

HOW DO THEY HOLD COMPARE TO COMPLETELY PAPER, PLASTIC OR BOTTLE? IMPROVED DOWN BY 4 TIMES! MULTIPLE RECYCLES BOTTLE TO RECYCLE

Your pharmacist recommended the Walgreens brand. Why? It's you to compare to national brands.

This product is not manufactured or distributed by WALGREEN CO. S.A., owner of the registered trademark EXCEDRIN®.

DISTRIBUTED BY: WALGREEN CO.
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
ITEM 767159 W0000-0000-0
1 1917 08167 0

W3OR00126-F NDC 093-4633-12

Walgreens
Compare to the active ingredients
in Excedrin® Extra Strength†

Headache Relief
ACETAMINOPHEN 250 mg / PAIN RELIEVER
ASPIRIN (NSAID) 250 mg / PAIN RELIEVER
CAFFEINE 65 mg / PAIN RELIEVER AID

Extra Strength

80
GELTABS

ACTUAL SIZE

Drug Facts

Active ingredients (in each geltab) **Purposes**
 Acetaminophen 250 mg Pain reliever
 Aspirin 250 mg (NSAID) Pain reliever
 Caffeine 65 mg Pain reliever and
 "nonsteroidal anti-inflammatory drug"

Uses
 Temporarily relieves minor aches and pains due to:
 ■ headache ■ a cold ■ arthritis ■ muscular aches
 ■ toothache ■ premenstrual and menstrual cramps

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 8 geltabs in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or other oral drug
 ■ take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product.
 ■ take more or for a longer time than directed

CELEBRATIONS
80
Extra Strength
ACETAMINOPHEN 250 mg / PAIN RELIEVER
ASPIRIN (NSAID) 250 mg / PAIN RELIEVER
CAFFEINE 65 mg / PAIN RELIEVER AID

Headache Relief
Walgreens

Drug Facts (continued)

Caffeine warnings: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat.

Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
 ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use ■ if you have liver disease
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you are taking a diuretic
 ■ you have asthma

Ask a doctor or pharmacist before use if you are taking
 ■ a prescription drug for diabetes, gout, or arthritis
 ■ any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ red, dark, or vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ ringing in the ears or loss of hearing occurs
 ■ painful area in mid or lower back
 ■ pain gets worse or lasts for more than 10 days
 ■ fever gets worse or lasts for more than 3 days
 ■ any new symptoms occur

These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children.
Overdose warning: In cases of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions
 ■ do not use more than directed (see overdose warning)
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 2 geltabs every 4 hours; not more than 8 geltabs in 24 hours
 ■ children under 12 years: ask a doctor

Other information
 ■ store at room temperature between 20-25°C (68-77°F)
 ■ avoid excessive heat, cold and humidity

Inactive ingredients
 corn starch, GPC yellow #10 aluminum lake, edible black ink, stibic calcium, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hypromellose, isopropyl alcohol, lactose monohydrate, light mineral oil, microcrystalline cellulose, pregelatinized potato starch, purified water, poly-20, sorbitol solution, stearic acid, talc, titanium dioxide, vitamin

Questions or comments?
 1-888-838-8782 Monday-Friday 9am-4pm EST

DISTRIBUTED BY: WALGREEN CO.
 OBERFIELD, IL 60455
 100% SATISFACTION GUARANTEED
 walgreens.com
 ©2005 Walgreen Co. MADE IN INDIA
 ITEM 757136 W00000-0000-0



3 11917 08172 4



†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by WALGREEN GELTABS, owner of the registered trademark EXCEDRIN®.

W30R00126-F NDC (683-693)-60

READ AND KEEP OUTER CAUTION ON COMPLETE PRODUCT WARNINGS AND INFORMATION

DAIPIR EVIDENT; DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



WALGREENS HEADACHE RELIEF EXTRA STRENGTH

acetaminophen 250mg, aspirin 250mg and caffeine 65mg tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6693
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	

WATER (UNII: 059QF0KO0R)
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)
STEARIC ACID (UNII: 4ELV7Z65AP)
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)
HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)
POVIDONE K30 (UNII: U725QWY32X)
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)
ISOPROPYL ALCOHOL (UNII: ND2M416302)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
HYPROMELLOSE 2910 (15 MPA.S) (UNII: 36SFW2JZ0W)
TALC (UNII: 7SEV7J4R1U)
STARCH, CORN (UNII: O8232NY3SJ)
GLYCERIN (UNII: PDC6A3C0OX)
FERROSO FERRIC OXIDE (UNII: XM0M87F357)

Product Characteristics

Color	green (Dark Green and White Opaque)	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	43
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6693-20	1 in 1 CARTON	05/01/2026	
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:0363-6693-80	1 in 1 CARTON	05/01/2026	
2		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:0363-6693-12	1 in 1 CARTON	05/01/2026	
3		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/01/2026	

