

**FAST ACTING MUCUS RELIEF MAXIMUM STRENGTH SEVERE COUGH AND CONGESTION- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid**  
**Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

-----

## **Drug Facts**

### **Active ingredients (in each 20 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

### **Purposes**

Cough suppressant

Expectorant

Nasal decongestant

### **Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

### **Warnings**

#### **Do not use**

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### **Ask a doctor before use if you have**

- heart disease
- thyroid disease
- diabetes
- high blood pressure

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

**When using this product,  
do not use more than directed**

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in a 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

**Other information**

- **each 20 mL contains:**sodium 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate

**Inactive ingredients**

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, sodium citrate, xanthan gum

**Principal Display Panel**

Compare to the active ingredients of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough\*

Maximum Strength

**Fast-Acting**

**Mucus Relief**

Severe Cough & Congestion

Multi-Symptom Relief

Dextromethorphan HBr

Guaifenesin

Phenylephrine HCl

Cough suppressant

Expectorant

Nasal decongestant

fl oz (mL)

For ages 12 and over

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE,

GOODLETTSVILLE, TN 37072

**Package Label**

**DG** health

Compare to the active ingredients of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough\*

**Maximum Strength Fast Acting Mucus Relief Severe Congestion & Cough**

Multi-Symptom Relief

Dextromethorphan HBr

Guaifenesin

Phenylephrine HCl

Cough Suppressant

Expectorant

Nasal Decongestant

For ages 12 and over

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

100\* Satisfaction Guaranteed! (888) 309-9030

A0514 PLD-E283D LB008190



9 fl oz  
(266 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

PLD-C283D  
LB003126

**Drug Facts**

**Active ingredients Purposes (in each 20 mL)**

Dextromethorphan HBr 20 mg.....  
.....Cough suppressant  
Guaifenesin 400 mg.....Expectorant  
Phenylephrine HCl 10 mg.....  
.....Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
- nasal congestion due to a cold

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

**Drug Facts (continued)**

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.**

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

**Other information**

- each 20 mL contains: sodium 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium

**Drug Facts (continued)**

benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**DOLLAR GENERAL HEALTH Maximum Strength Mucus Relief**

# FAST ACTING MUCUS RELIEF MAXIMUM STRENGTH SEVERE COUGH AND CONGESTION

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-037
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D45NN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-037-09	266 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/31/2014	02/01/2027

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/31/2014	02/01/2027

**Labeler** - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 11/2025

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)