

**XRAY PAIN RELIEF- menthol, (plus)-, camphor, (-)-, and methyl salicylate cream**  
**Genomma Lab USA**

-----  
**XRay® Pain Relief Cream**

***Drug Facts***

<b><i>Active ingredients</i></b>	<b><i>Purpose</i></b>
Camphor 4%	Topical analgesic
Menthol 10%	Topical analgesic
Methyl salicylate 30%	Topical analgesic

**Uses**

- temporarily relieves the minor aches and pains of muscles and joints associated with:
  - sprains
  - simple backache
  - arthritis
  - strains
  - bruises

**Warnings**

**For external use only**

**Allergy alert**

If prone to allergic reaction from aspirin or salicylates, consult a doctor before use.

**Do not use**

- on wounds or damaged skin
- with a heating pad
- on a child under 12 years of age with arthritis-like conditions

**Ask a doctor before use if you have** redness over the affected area.

**When using this product**

- do not use otherwise than as directed
- avoid contact with eyes, mucous membranes or rashes
- do not bandage tightly

**Stop use and ask a doctor if**

- redness is present or irritation or rash develops
- condition worsens or symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- use only as directed
- adults and children 12 years of age and older: apply to affected area not more than 3 to 4 times daily
- children under 12 years of age: ask a doctor

### **Other information**

store at room temperature 20-25°C (68-77°F)

### **Inactive ingredients**

ethylparaben, glyceryl monostearate, polysorbate 80, purified water, sorbitan monooleate, stearic acid

### **Questions or comments?**

Call toll free 1 877 994 3666

Monday to Friday 8:00 am – 6:00 pm CST

### **Distributed by:**

**Genomma Lab USA, Inc.,  
Houston, TX 77098**

### **PRINCIPAL DISPLAY PANEL - 56.7 G Tube Carton**

X RAY®

DOL

ULTRA STRENGTH

Pain Relief Cream

FOR DAILY TREATMENT OF  
ARTHRITIS AND JOINT PAIN\*

DEEP PENETRATING

FAST ACTING  
RELIEF

Topical Analgesic

\*See Drug Facts for more information

Camphor 4%

Menthol 10%

Methyl salicylate 30%

NET WT.  
2 OZ (56.7 G)

GLUE AREA / No Ink / No Varnish

# X RAY<sup>®</sup> DOL

MÁXIMA POTENCIA

## Crema para Alivio del Dolor

PARA EL TRATAMIENTO DIARIO DEL DOLOR ARTICULAR Y DE ARTRITIS\*



ACCIÓN PROFUNDA

RÁPIDO ALIVIO

Analgésico Tópico

\*Lea la Información de Medicamento

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPEN OR FOIL UNDER CAP IS BROKEN OR MISSING.

Drug Facts

Active Ingredients	Purpose
Camphor 4%.....	Topical analgesic
Menthol 10%.....	Topical analgesic
Methyl salicylate 30%.....	Topical analgesic

**Uses** ■ temporarily relieves the minor aches and pains of muscles and joints associated with: ■ sprains ■ simple backache ■ arthritis ■ strains ■ bruises

**Warnings**  
For external use only  
Allergy alert: If prone to allergic reaction from aspirin or salicylates, consult a doctor before use.  
**Do not use** ■ on wounds or damaged skin ■ with a heating pad ■ on a child under 12 years of age with arthritis-like conditions  
**Ask a doctor before use if you have redness over the affected area.**

**When using this product** ■ do not use otherwise than as directed ■ avoid contact with eyes, mucous membranes or rashes: ■ do not bandage tightly

**Stop use and ask a doctor if** ■ redness is present or irritation or rash develops ■ condition worsens or symptoms persist for more than 7 days ■ symptoms clear up and occur again within a few days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**  
■ use only as directed ■ adults and children 12 years of age and older: apply to affected area not more than 3 to 4 times daily  
■ children under 12 years of age: ask a doctor

**Other information**  
store at room temperature 20-25°C (68-77°F)

**Inactive ingredients**  
ethylparaben, glyceryl monostearate, polysorbate 80, purified water, sorbitan monooleate, stearic acid

**Questions or comments?**  
Call toll free 1 877 994 3666  
Monday to Friday 8:00 am – 6:00 pm CST

Made in / Hecho en China  
Distributed by / Distribuido por:  
Genomma Lab USA, Inc.,  
Houston, TX 77098



CONSERVE LA CAJA EXTERNA PARA ADVERTENCIAS E INFORMACION COMPLETAS DEL PRODUCTO.

EVIDENCIA DE VIOLACION: NO UTILIZAR SI LA CAJA ESTA ABIERTA O LA LAMINA BAJO LA TAPA ESTA ROTA O FALTA.

Información del Medicamento

Ingredientes activos	Propósitos
Alcanfor 4 %.....	Analgésico tópico
Mentol 10%.....	Analgésico tópico
Salicilato de metilo 30%.....	Analgésico tópico

**Usos** ■ alivia temporalmente los dolores y molestias menores de músculos y articulaciones asociados con: ■ esguinces ■ dolor de espalda ■ artritis ■ distensiones ■ contusiones

**Advertencias**  
Solo para uso externo  
**Alerta de alergia:** si es propenso a una reacción alérgica a la aspirina o los salicilatos, consulte a un médico antes de usar.  
**No lo use** ■ en heridas o piel dañada ■ con una almohadilla térmica ■ en niños menores de 12 años con condiciones similares a la artritis  
**Consulte a un médico antes de usar si tiene enrojecimiento en el área afectada.**

**Cuando use este producto** ■ no lo use de otra manera que no sea como se indica. ■ evite el contacto con los ojos, las membranas mucosas o las erupciones ■ no use vendajes apretados

**Deje de usarlo y consulte a un médico si** ■ hay enrojecimiento, irritación o sarpullido ■ la condición empeora o los síntomas persisten durante más de 7 días ■ los síntomas desaparecen y vuelven a aparecer en unos pocos días

**Si está embarazada o amamantando,** consulte a un profesional de la salud antes de usar.

**Mantener fuera del alcance de los niños.**  
Si se ingiere, obtenga ayuda médica o comuníquese con un Centro de control de envenenamiento (1-800-222-1222) de inmediato.

**Instrucciones** ■ use solo como se indica ■ adultos y niños mayores de 12 años: aplique en el área afectada no más de 3 a 4 veces al día ■ niños menores de 12 años: consulte a un médico

**Otra información** almacene a temperatura ambiente 20-25° C (68-77°F)

**Ingredientes inactivos**  
etilparabeno, monoestearato de glicerilo, polisorbato 80, agua purificada, monooleato de sorbitán, ácido esteárico.

**¿Preguntas o comentarios?**  
Lláme lada sin costo 1 877 994 3666.  
Lunes a viernes 8:00 am – 6:00 pm CST

IM TP-Z0P124 R2  
IM MD-Z0P112 R2

2000011966

GLUE AREA / No Ink / No Varnish

# X RAY<sup>®</sup> DOL

ULTRA STRENGTH

## Pain Relief Cream

FOR DAILY TREATMENT OF ARTHRITIS AND JOINT PAIN\*



DEEP PENETRATING

FAST ACTING RELIEF

Topical Analgesic

\*See Drug Facts for more information

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPEN OR FOIL UNDER CAP IS BROKEN OR MISSING.

Drug Facts

Active Ingredients	Purpose
Camphor 4%.....	Topical analgesic
Menthol 10%.....	Topical analgesic
Methyl salicylate 30%.....	Topical analgesic

**Uses** ■ temporarily relieves the minor aches and pains of muscles and joints associated with: ■ sprains ■ simple backache ■ arthritis ■ strains ■ bruises

**Warnings**  
For external use only  
Allergy alert: If prone to allergic reaction from aspirin or salicylates, consult a doctor before use.  
**Do not use** ■ on wounds or damaged skin ■ with a heating pad ■ on a child under 12 years of age with arthritis-like conditions  
**Ask a doctor before use if you have redness over the affected area.**

**When using this product** ■ do not use otherwise than as directed ■ avoid contact with eyes, mucous membranes or rashes: ■ do not bandage tightly

**Stop use and ask a doctor if** ■ redness is present or irritation or rash develops ■ condition worsens or symptoms persist for more than 7 days ■ symptoms clear up and occur again within a few days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**  
■ use only as directed ■ adults and children 12 years of age and older: apply to affected area not more than 3 to 4 times daily  
■ children under 12 years of age: ask a doctor

**Other information**  
store at room temperature 20-25°C (68-77°F)

**Inactive ingredients**  
ethylparaben, glyceryl monostearate, polysorbate 80, purified water, sorbitan monooleate, stearic acid

**Questions or comments?**  
Call toll free 1 877 994 3666  
Monday to Friday 8:00 am – 6:00 pm CST

Made in / Hecho en China  
Distributed by / Distribuido por:  
Genomma Lab USA, Inc.,  
Houston, TX 77098



CONSERVE LA CAJA EXTERNA PARA ADVERTENCIAS E INFORMACION COMPLETAS DEL PRODUCTO.

EVIDENCIA DE VIOLACION: NO UTILIZAR SI LA CAJA ESTA ABIERTA O LA LAMINA BAJO LA TAPA ESTA ROTA O FALTA.

Información del Medicamento

Ingredientes activos	Propósitos
Alcanfor 4 %.....	Analgésico tópico
Mentol 10%.....	Analgésico tópico
Salicilato de metilo 30%.....	Analgésico tópico

**Usos** ■ alivia temporalmente los dolores y molestias menores de músculos y articulaciones asociados con: ■ esguinces ■ dolor de espalda ■ artritis ■ distensiones ■ contusiones

**Advertencias**  
Solo para uso externo  
**Alerta de alergia:** si es propenso a una reacción alérgica a la aspirina o los salicilatos, consulte a un médico antes de usar.  
**No lo use** ■ en heridas o piel dañada ■ con una almohadilla térmica ■ en niños menores de 12 años con condiciones similares a la artritis  
**Consulte a un médico antes de usar si tiene enrojecimiento en el área afectada.**

**Cuando use este producto** ■ no lo use de otra manera que no sea como se indica. ■ evite el contacto con los ojos, las membranas mucosas o las erupciones ■ no use vendajes apretados

**Deje de usarlo y consulte a un médico si** ■ hay enrojecimiento, irritación o sarpullido ■ la condición empeora o los síntomas persisten durante más de 7 días ■ los síntomas desaparecen y vuelven a aparecer en unos pocos días

**Si está embarazada o amamantando,** consulte a un profesional de la salud antes de usar.

**Mantener fuera del alcance de los niños.**  
Si se ingiere, obtenga ayuda médica o comuníquese con un Centro de control de envenenamiento (1-800-222-1222) de inmediato.

**Instrucciones** ■ use solo como se indica ■ adultos y niños mayores de 12 años: aplique en el área afectada no más de 3 a 4 veces al día ■ niños menores de 12 años: consulte a un médico

**Otra información** almacene a temperatura ambiente 20-25° C (68-77°F)

**Ingredientes inactivos**  
etilparabeno, monoestearato de glicerilo, polisorbato 80, agua purificada, monooleato de sorbitán, ácido esteárico.

**¿Preguntas o comentarios?**  
Lláme lada sin costo 1 877 994 3666.  
Lunes a viernes 8:00 am – 6:00 pm CST

IM TP-Z0P124 R2  
IM MD-Z0P112 R2

2000011966

GLUE AREA / No Ink / No Varnish

# X RAY<sup>®</sup> DOL

ULTRA STRENGTH

## Pain Relief Cream

GLUE AREA / No Ink / No Varnish

Alcanfor 4% | Mentol 10% | Salicilato de metilo 30%

NET WT. 2 OZ (56.7 G)

GLUE AREA / No Ink / No Varnish

Camphor 4% | Menthol 10% | Methyl salicylate 30%

NET WT. 2 OZ (56.7 G)

LOT NO. \_\_\_\_\_  
EXP. \_\_\_\_\_  
Blister Packs must clear the LOT and EXP.



6 50240 06454 8

# XRAY PAIN RELIEF

menthol, (plus)-, camphor, (-)-, and methyl salicylate cream

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50066-456
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MENTHOL, (+)-</b> (UNII: C6B1OE8P3W) (MENTHOL, (+)- - UNII:C6B1OE8P3W)	MENTHOL, (+)-	100 mg in 1 g
<b>CAMPHOR, (-)-</b> (UNII: 213N3S8275) (CAMPHOR, (-)- - UNII:213N3S8275)	CAMPHOR, (-)-	40 mg in 1 g
<b>METHYL SALICYLATE</b> (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:O414PZ4LPZ)	METHYL SALICYLATE	300 mg in 1 g

## Inactive Ingredients

Ingredient Name	Strength
<b>glyceryl monostearate</b> (UNII: 230OU9XXE4)	
<b>Water</b> (UNII: 059QF0KO0R)	
<b>Polysorbate 80</b> (UNII: 6OZP39ZG8H)	
<b>sorbitan monooleate</b> (UNII: 06XEA2VD56)	
<b>stearic acid</b> (UNII: 4ELV7Z65AP)	
<b>ethylparaben</b> (UNII: 14255EXE39)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50066-456-20	1 in 1 CARTON	12/26/2023	
1		56.7 g in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M017	12/26/2023	

**Labeler** - Genomma Lab USA (832323534)

Revised: 12/2023

Genomma Lab USA