

QUALITY CHOICE LIDOCAINE PAIN RELIEF PATCH- lidocaine patch
Quality Choice (CHAIN DRUG MARKETING ASSOCIATION)

Quality Choice Lidocaine Pain Relief Patch

Active Ingredients

Lidocaine 4% Topical Anesthetic

Purpose

Topical anesthetic

Indications and Usage

For temporary relief of pain.

Warnings

For external use only.

Do not Use

- more than one patch on your body at a time
- on cut, irritated or swollen skin
- on puncture wounds
- for more than one week without consulting a doctor
- if you are allergic to any active or inactive ingredients
- if pouch is damaged or opened.

When using this product

- use only as directed
- read and follow all directions and warnings on this carton
- do not allow contact with the eyes
- do not use at the same time as other topical analgesics
- do not bandage tightly or apply local heat (such as heating pads) to the area of use
- do not microwave
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse

effects if a child or pet chews or ingests this patch.

Stop use and consult a doctor

- condition worsens
- redness is present
- irritation develops
- symptoms persist for more than 7 days or clear up and occur again within a few days
- you experience signs of skin injury, such as pain, swelling or blistering where the product was applied.

ask a health professional before use.

Keep out of reach of children and pets.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Dosage and Administration

Adults and children 12 years of age and over:

- Clean and dry affected area, free of lotions, ointments and creams
- Carefully remove backing from patch starting at a corner
- Apply sticky side of patch to affected area
- Do not use more than one patch in a 12 hour period
- Maximum 2 patches per day
- Discard patch after single use

Children under 12 years of age:

Consult a physician

Other Safety Information

Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Inactive Ingredients

crospovidone, dihydroxyaluminum aminoacetate, edetate disodium, glycerin, kaolin, methylparaben, polyacrylic acid, polysorbate 80, propylene glycol, propylparaben, sodium polyacrylate, tartaric acid, titanium dioxide, water

Questions

800-935-2362



Maximum OTC Strength Patch
Lidocaine Pain Relief
4% Lidocaine, Topical Anesthetic



Maximum OTC Strength Patch
Lidocaine Pain Relief
4% Lidocaine, Topical Anesthetic

NDC 83324-257-05

*Compare to the Active Ingredient in Aspercreme® Lidocaine Patch

Drug Facts

Active Ingredient Lidocaine 4.0%	Purpose Topical anesthetic
Use For temporary relief of pain.	
Warnings For external use only.	
Do not use ■ More than one patch on your body at a time ■ On cut, irritated or swollen skin ■ On puncture wounds ■ Over raw or broken or irritated areas ■ For more than one week without consulting a doctor ■ If you are allergic to any active or inactive ingredients ■ If pouch is damaged or opened.	
When using this product ■ Use only as directed ■ Read and follow all directions and warnings on this carton ■ Do not allow contact with the eyes ■ Do not use at the same time as other topical anesthetics ■ Do not package tightly or apply local heat (such as heating pads) to the area or use ■ Do not microwave ■ Dispose of used patch in container that always keeps product away from children and pets. Used patches still contain the drug product and can produce serious adverse effects if a child or pet chews or ingests this patch.	
Stop use and consult a doctor if ■ Condition worsens ■ Redness is present ■ Irritation develops ■ Symptoms persist for more than 7 days or clear up and occur again within a few days ■ You experience signs of skin injury, such as pain, swelling or blistering where the product was applied.	
If pregnant or breastfeeding, ask a health professional before use.	
Keep out of reach of children and pets. If swallowed, get medical help or contact Poison Control Center (1-800-222-1222) right away.	
Directions ■ Adults and children 12 years of age and over ■ Clean and dry affected area, use ointments, ointments and creams. Carefully remove banding from patch starting at a corner. Apply sticky side of patch to affected area. Do not use more than one patch in a 12-hour period. Maximum 2 patches per day. Discard patch after single use ■ Children under 12 years of age: consult a physician.	
Other information Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.	
Inactive ingredients croscellose, dibutyltinyltin diisobutyrate, edetate disodium, glycerin, kaolin, methylparaben, polyacrylic acid, polyisobutylene, polyethylene glycol, propylparaben, sodium polyacrylate, tartaric acid, titanium dioxide, water	

*This product is not manufactured or distributed by Chastain, Inc., owner of the registered trademark Aspercreme®.

Maximum OTC Strength Patch
Lidocaine Pain Relief

Maximum OTC Strength Patch
Lidocaine Pain Relief

5 Patches 3.93 in. x 5.51 in. (10 cm x 14 cm)



6 35515 99627 4

ITEM NO. 89-911



Distributed by: COMA, Inc.
New, HI 48275
www.qualitychoice.com
Guarantee: 800-935-2262

Made in China
32-63047 05/25
VC310047

LOT
EXP

QUALITY CHOICE LIDOCAINE PAIN RELIEF PATCH

lidocaine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83324-257
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
PHENOXYETHANOL (UNII: H1E492ZZ3T)	
DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6)	
GLYCERIN (UNII: PDC6A3C0OX)	
KAOLIN (UNII: 24H4NWX5CO)	
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)	

POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
CROSPVIDONE (UNII: 2S7830E561)	
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05115JN12J)	
TARTARIC ACID (UNII: W4888I119H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
METHYLPARABEN (UNII: A2I8C7HI9T)	

Product Characteristics

Color		Score	
Shape	RECTANGLE	Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83324-257-05	30 in 1 CARTON	05/01/2026	
1		9 g in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	05/01/2026	

Labeler - Quality Choice (CHAIN DRUG MARKETING ASSOCIATION) (011920774)

Establishment

Name	Address	ID/FEI	Business Operations
Shanghai Chuangshi Medical Technology (Group) Co., Ltd		546872672	manufacture(83324-257)

Revised: 5/2026

Quality Choice (CHAIN DRUG MARKETING ASSOCIATION)