

ASPIRIN- aspirin tablet
United Exchange Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

United Exchange 44-249

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- headache
- toothache
- muscle pain
- pain and fever of colds
- menstrual pain
- minor pain of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take more of for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient

corn starch

Questions or comments?

1-800-426-9391

Principal display panel

ASPIRIN

(NSAID)

Pain Reliever • Fever Reducer

For the temporary relief from the pain of:

- headache • muscle pain • toothache
- menstrual pain • pain and fever of colds
- minor pain of arthritis

100 TABLETS

Actual Size

325 mg each

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

50844 REV0318A24912

Distributed By: UNITED EXCHANGE CORP.

Cypress, CA 90630 USA

1-888-645-8204

<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients 	<p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ you have asthma ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for</p> <ul style="list-style-type: none"> ■ gout ■ diabetes ■ arthritis 	<p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur <p>These could be signs of a serious condition.</p>	<p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>	<p>Directions</p> <ul style="list-style-type: none"> ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package 	<p>Inactive ingredient corn starch</p> <p>Questions or comments? 1-800-426-9391</p>
--	--	--	---	--	---

44-249

<div>ASPIRIN</div> <div>aspirin tablet</div>			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65923-249
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg
Inactive Ingredients			
Ingredient Name			Strength
STARCH, CORN (UNII: O8232NY3SJ)			

Product Characteristics				
Color	WHITE	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;249;ASPIRIN	
Contains				

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65923-249-12	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	08/04/1993	

Labeler - United Exchange Corporation (840130579)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(65923-249)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	MANUFACTURE(65923-249)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(65923-249)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(65923-249)