

**COLD AND FLU MULTI-SYMPATOM DAYTIME/NIGHTTIME SEVERE- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, guaifenesin and phenylephrine hcl
COSTCO WHOLESALE CORPORATION**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Kirkland 44-503A473C-R

***Active ingredients (in each caplet)
(Daytime Cold & Flu Severe)***

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

***Active ingredients (in each caplet)
(Nighttime Cold & Flu Severe)***

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Antihistamine
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - headache
 - sore throat
 - minor aches and pains
 - nasal congestion
 - sinus congestion and pressure (*Nighttime only*)
 - sneezing and runny nose (*Nighttime only*)
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive (*Daytime only*)
- helps clear nasal passages (*Nighttime only*)

- relieves cough to help you sleep (**Nighttime only**)
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- heart disease
- glaucoma (**Nighttime only**)
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis (**Nighttime only**)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Nighttime only**)

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children (**Nighttime only**)
- marked drowsiness may occur (**Nighttime only**)
- avoid alcoholic beverages (**Nighttime only**)
- use caution when driving a motor vehicle or operating machinery (**Nighttime only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nighttime only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Read each section carefully. Do not take DAYTIME and NIGHTTIME products at the same time.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole - do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- see end flap for expiration date and lot number
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients (Daytime only)

corn starch, crospovidone, D&C yellow #10 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Inactive ingredients (Nighttime only)

corn starch, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391



Kirkland 44-503A473C

COLD AND FLU MULTI-SYMPTOM DAYTIME/NIGHTTIME SEVERE

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, guaifenesin and phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:6398 1-702
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:63981-702-95	1 in 1 CARTON	04/09/2020
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1		8
Part 2		4

Part 1 of 2

COLD AND FLU MULTI-SYMP TOM DAYTIME SEVERE

acetaminophen, dextromethorphan hbr and phenylephrine hcl tablet, film coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 2S7830E561)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;503
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	04/09/2020	

Part 2 of 2

COLD AND FLU MULTI-SYMP TOM NIGHTTIME SEVERE

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr and phenylephrine hcl tablet, film coated

Product Information

Route of Administration	ORAL
-------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FD&C BLUE NO. 2--ALUMINUM LAKE (UNII: 4AQJ3LG584)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	BLUE	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	44;473
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	04/09/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	04/09/2020	

Labeler - COSTCO WHOLESALE CORPORATION (103391843)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(63981-702)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(63981-702)

Revised: 5/2020

COSTCO WHOLESALE CORPORATION