


MENTAL SKIN CARE LIP SHIELD IMPACT- zinc oxide lotion
Mental Skin

Mental Skin Care Lip Shield Impact

Mental Skin Care Lip Shield Impact



DRUG FACTS (continued)

To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including:

- limit your time in the sun, especially from 10 a.m. — 2 p.m.
- wear long-sleeved shirts, pants, hats, and sunglasses

Other Information - Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients: Hydrogenated Poly(C6-14 Olefin), Squalane, Butyltolyl Salicylate, Cetearyl Dimethicone, Isododecane, Ethylene/Propylene/Styrene Copolymer, Caprylyl Methicone, Synthetic Fluorophlogopite, Iron Oxides, Lecithin, Dimethiconol/Propylsilsesquiox

ane/Silicate Crosspolymer, Titanium Dioxide, Phenylpropanediol, Propylene Glycol, Tin Oxide, Butylene/Ethylene/Styrene Copolymer, Pentaerythrityl Tetra-*n*-butyl Hydroxyhydrocinnamate, Dimethicone, Tocopherol.

*less or equal to 1%

You are stronger than you think...

Mental Skincare, LLC
 Irvine, CA 92620
 Question or
 Comments?
 US 888-883-1891
 MentalSkincare.com
 MADE IN USA

DRUG FACTS


Active Ingredients **Purpose**
 Zinc Oxide 10% Sunscreen

Uses - Helps prevent sunburn. If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer & early skin aging caused by the sun.

Warnings - For external use only. • Do not use on damaged or broken skin. • When using this product keep out of eyes. Rinse with water to remove. • Stop use and ask a doctor if skin rash occurs. • Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions - apply liberally 15 minutes before sun exposure. • Reapply at least every 2 hours. • use a water-resistant sunscreen if swimming or sweating. • Children under 6 months of age. Ask a doctor. • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging.

Mental Skin Care Lip Shield Impact



Mental
Broad Spectrum SPF 35

LIP TREATMENT SUNSCREEN
BROAD SPECTRUM SPF 35 UVA/UVB
Net 0.11 fl. oz. / 3.5 mL

IMPACT

DRUG FACTS (continued)
To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including:
• limit your time in the sun, especially from 10 a.m. – 2 p.m. • wear long-sleeved shirts, pants, hats, and sunglasses

Other Information • Protect the product in this container from excessive heat and direct sun.
Inactive Ingredients: Hydrogenated Poly(C6-14 Olefin), Squalane, Butyloctyl Salicylate, Octyl Dimethicone, Isododecane, Ethylene/Propylene/Styrene Copolymer, Caprylyl Methicone, Synthetic Fluorophlogopite, Iron Oxides, Lecithin, Dimethiconol/Propylsilsesquioxane/Silicate Crosspolymer, Titanium Dioxide, Phenylpropanol, Propanediol, Caprylyl Glycol, Tin Oxide, Butylene/Ethylene/Styrene Copolymer, Panthenol, Tetra-*n*-butyl hydroxyethylpyridone-3-carboxylate, Dimethicone, Tocopherol.

*less or equal to 1%

You are stronger than you think...

Mental Skincare, LLC
Irvine, CA 92620
Question or Comments?
US 848-883-1891
mentalskincare.com
MADE IN USA

DRUG FACTS
Active Ingredients
Zinc Oxide 10%
Purpose
Sunscreen
Uses • Helps prevent sunburn • If used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer & early skin aging caused by the sun.


Warnings • For external use only. • Do not use on damaged or broken skin • When using this product keep out of eyes. Rinse with water to remove • Stop use and ask a doctor if skin rash occurs • Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions • apply liberally 15 minutes before sun exposure • Reapply at least every 2 hours • use a water-resistant sunscreen if swimming or sweating • Children under 6 months of age: Ask a doctor. • Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging.

Mental Skin Care Lip Shield Impact




<p>DRUG FACTS (continued)</p> <p>To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including:</p> <ul style="list-style-type: none"> • limit your time in the sun, especially from 10 a.m. – 2 p.m. • wear long-sleeved shirts, pants, hats, and sunglasses 	<p>Other Information • Protect the product in this container from excessive heat and direct sun.</p> <p>Inactive Ingredients: Hydrogenated Poly(C9-14 Olefin), Squalane, Butylloctyl Salicylate, Cetearyl Dimethicone, Isododecane, Ethylene/Propylene/Styrene Copolymer, Caprylyl Methicone, Synthetic Fluorophlogopite, Iron Oxides, Lecithin, Dimethiconol/Propylsilsesquiox</p>	<p>ane/Silicate Crosspolymer, Titanium Dioxide, Phenylpropanediol, Pentaerythrityl Glycol, Tin Oxide, Butylene/Ethylene/Styrene Copolymer, Pentaerythrityl Tetra-di-<i>n</i>-butyl Hydroxyhydrocinnamate, Dimethicone, Tocopherol.</p> <p>*less or equal to 1%</p>
---	---	--



<p>DRUG FACTS</p> <p>Active Ingredients Zinc Oxide 10%</p> <p>Purpose Sunscreen</p> <p>Uses • Helps prevent sunburn • If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer & early skin aging caused by the sun.</p>	<p>Warnings • For external use only. • Do not use on damaged or broken skin. • When using this product keep out of eyes. Rinse with water to remove. • Stop use and ask a doctor if skin rash occurs. • Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.</p>	<p>Directions • apply liberally 15 minutes before sun exposure • Reapply at least every 2 hours • use a water-resistant sunscreen if swimming or sweating • Children under 6 months of age: Ask a doctor. • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging.</p>
---	--	--

Mental Skin Care Lip Shield Impact



DRUG FACTS (continued)

To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including: • limit your time in the sun, especially from 10 a.m. – 2 p.m. • wear long-sleeved shirts, pants, hats, and sunglasses

Other Information • Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients: Hydrogenated Poly(C6-14 Olefin), Squalane, Butyloctyl Salicylate, Octyl Dimethicone, Isododecane, Ethylene/Propylene/Styrene Copolymer, Caprylyl Methicone, Synthetic Fluorophlogopite, Iron Oxides, Lecithin, Dimethiconol/Propylsiloxane

ane/Silicate, Crosspolymer, Titanium Dioxide, Phenylpropanol, Propanediol, Caprylyl Glycol, Tin Oxide, Butylene/Ethylene/Styrene Copolymer, Panthenol, Tetra-*n*-butyl hydroxyhydroxamate, Dimethicone, Tocopherol.

*less or equal to 1%

You are stronger than you think...

Mental Skincare, LLC
 Irvine, CA 92620
 Question or Comments?
 US 888-883-1891
 MentalSkincare.com
 MADE IN USA

DRUG FACTS


Active Ingredients	Purpose
Zinc Oxide 10%	Sunscreen

Uses • Helps prevent sunburn • If used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer & early skin aging caused by the sun.

Warnings • For external use only. • Do not use on damaged or broken skin • When using this product keep out of eyes. Rinse with water to remove • Stop use and ask a doctor if skin rash occurs • Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions • apply liberally 15 minutes before sun exposure • Reapply at least every 2 hours • use a water-resistant sunscreen if swimming or sweating • Children under 6 months of age. Ask a doctor. • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging.

Mental Skin Care Lip Shield Impact



DRUG FACTS (continued)

To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including: • limit your time in the sun, especially from 10 a.m. – 2 p.m. • wear long-sleeved shirts, pants, hats, and sunglasses

Other Information • Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients: Hydrogenated Poly(C6-14 Olefin), Squalane, Butylcetyl Salicylate, Cetearyl Dimethicone, Isododecane, Ethylene/Propylene/Styrene Copolymer, Caprylyl Methicone, Synthetic Fluorophlogopite, Iron Oxides, Lecithin, Dimethiconol/Propylsesquiox

ane/Silicate, Crosspolymer, Titanium Dioxide, Phenylpropanol, Propylalcohol, Caprylyl Glycol, Tin Oxide, Butylene/Ethylene/Styrene Copolymer, Pentaerythrityl Tetra-di-t-butyl Hydroxyhydrocinnamate, Dimethicone, Tocopherol.

*less or equal to 1%

You are stronger than you think...

Mental Skincare, LLC
Irvine, CA 92620
Question or
Comments?
US 888-883-1891
MentalSkincare.com
MADE IN USA

DRUG FACTS


Active Ingredients Purpose
Zinc Oxide 10% Sunscreen

Uses • Helps prevent sunburn • If used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer & early skin aging caused by the sun.

Warnings • For external use only. • Do not use on damaged or broken skin. • When using this product keep out of eyes. Rinse with water to remove. • Stop use and ask a doctor if skin rash occurs. • Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions • apply liberally 15 minutes before sun exposure • Reapply at least every 2 hours • use a water-resistant sunscreen if swimming or sweating • Children under 6 months of age. Ask a doctor. • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging.

Mental Skin Care Lip Shield Impact



DRUG FACTS (continued)

To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including:

- limit your time in the sun, especially from 10 a.m. — 2 p.m.
- wear long-sleeved shirts, pants, hats, and sunglasses

Other Information - Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients: Hydrogenated Poly(C6-14 Olefin), Squalane, Butyltolyl Salicylate, Cetearyl Dimethicone, Isododecane, Ethylene/Propylene/Styrene Copolymer, Caprylyl Methicone, Synthetic Fluorophlogopite, Iron Oxides, Lecithin, Dimethiconol/Propylsilsesquiox

ane/Silicate Crosspolymer, Titanium Dioxide, Phenylpropanol, Propenediol, Caprylyl Glycol, Tin Oxide, Butylene/Ethylene/Styrene Copolymer, Pentaerythrityl Tetra-*n*-butyl Hydroxyhydrocinnamate, Dimethicone, Tocopherol.
*less or equal to 1%

IMPACT

You are stronger than you think...

Mental Skincare, LLC
Irvine, CA 92620
Question or Comments?
US 888-693-1891
MentalSkincare.com
MADE IN USA

DRUG FACTS


Active Ingredients Purpose
Zinc Oxide 10% Sunscreen

Uses - Helps prevent sunburn - If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer & early skin aging caused by the sun

Warnings - For external use only. - Do not use on damaged or broken skin - When using this product keep out of eyes. Rinse with water to remove. - Stop use and ask a doctor if skin rash occurs - Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions - apply liberally 15 minutes before sun exposure - Reapply at least every 2 hours - use a water-resistant sunscreen if swimming or sweating - Children under 6 months of age: Ask a doctor - Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging.

Mental Skin Care Lip Shield Impact



DRUG FACTS (continued)
To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including:
• limit your time in the sun, especially from 10 a.m. – 2 p.m. • wear long-sleeved shirts, pants, hats, and sunglasses


Other Information • Protect the product in this container from excessive heat and direct sun.
Inactive Ingredients: Hydrogenated Poly(C6-14 Olefin), Squalane, Butyloctyl Salicylate, Octyl Dimethicone, Isododecane, Ethylene/Propylene/Styrene Copolymer, Caprylyl Methicone, Synthetic Fluorophlogopite, Iron Oxides, Lecithin, Dimethiconol/Propylsiloxane/ane/Silicate Crosspolymer, Titanium Dioxide, Phenylpropanol, Propanediol, Caprylyl Glycol, Tin Oxide, Butylene/Ethylene/Styrene Copolymer, Pentaerythrityl Tetra-*n*-butyl hydroxyhydroxamate, Dimethicone, Tocopherol.
*less or equal to 1%

You are stronger than you think...

Mental Skincare, LLC
Irvine, CA 92620
Question or Comments?
US 888-883-1881
mentalskincare.com
MADE IN USA

DRUG FACTS Active Ingredients Zinc Oxide 10% Purpose Sunscreen	Warnings • For external use only. • Do not use on damaged or broken skin • When using this product keep out of eyes. Rinse with water to remove • Stop use and ask a doctor if skin rash occurs • Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.	Directions • apply liberally 15 minutes before sun exposure • Reapply at least every 2 hours • use a water-resistant sunscreen if swimming or sweating • Children under 6 months of age. Ask a doctor. • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging.
---	--	--

Mental Skin Care Lip Shield Impact



MENTAL SKIN CARE LIP SHIELD IMPACT

zinc oxide lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82696-0348
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	95 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SQUALANE (UNII: GW89575KF9)	
ISODODECANE (UNII: A8289P68Y2)	
BUTYLOCTYL SALICYLATE (UNII: 2EH13UN8D3)	
PROPANEDIOL (UNII: 5965N8W85T)	
PHENYLPROPANOL (UNII: 0F897O3O4M)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
DIMETHICONOL/PROPYLSILSESQUIOXANE/SILICATE CROSSPOLYMER (450000000 MW) (UNII: 9KB5R958PB)	
TOCOPHEROL (UNII: R0ZB2556P8)	
HYDROGENATED POLY(C6-14 OLEFIN; 4 CST) (UNII: 7C6HV4SSLA)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
CAPRYLYL GLYCOL (UNII: 00YIU5438U)	
TIN OXIDE (UNII: KM7N50LOS6)	
PENTAERYTHRITYL TETRA-DI-T-BUTYL HYDROXYHYDROCINNAMATE (UNII: 255PIF62MS)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
CAPRYLYL METHICONE (UNII: Q95M2P1KJL)	
MAGNESIUM POTASSIUM ALUMINOSILICATE FLUORIDE (UNII: YK3DC63Y5M)	
DIMETHICONE (UNII: 92RU3N3Y1O)	

Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:82696			

1	NDC:82096-0348-2	1 in 1 CARTON	03/13/2024	
1	NDC:82696-0348-1	3.5 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020		03/13/2024	

Labeler - Mental Skin (115554947)

Revised: 5/2026

Mental Skin