

MEDIQUE MEDI SELTZER- aspirin, citric acid, sodium bicarbonate tablet, effervescent
Unifirst First Aid Corporation

Medique Medi Seltzer

Drug Facts

Active ingredient (in each tablet)

Aspirin (NSAID*) 325 mg

Citric Acid 1000 mg

Sodium bicarbonate (heat-treated) 1916 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever /fever reducer

Antacid

Antacid

Uses

For the temporary relief of

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin,

ibuprofen, naproxen, or others)

- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/ fever reducer
- if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you are on a sodium restricted diet
- you reached age 60 or older

Ask a doctor or pharmacist before use if you are

- taking any other drugs (prescription or non-prescription). Antacids may interact with certain drugs.
- under a doctor's care for any serious condition

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - you feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin (***NSAIDS**) **at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage
- to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

Adults and children 12 to 60 years of age: 2 tablets every 4 hours. Do not exceed 8 tablets in 24 hours or do not use the maximum dosage for more than 10 days.

Children under 12 years of age: Do not give to children under 12 years of age.

60 years and older: Do not exceed 4 tablets in 24 hours or do not use the maximum dosage for more than 10 days.

Other information

- each tablet contains: Sodium 524 mg
- store at room temperature 59°-86° F (15°-30° C)
- read all product information before using
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

none

Questions or comments? 1-800-634-7680

Medique Medi-Seltzer Label

Medique®

Medi-Seltzer

Effervescent Antacid/Pain Reliever

Pull to Open

This Package is for Households without Young Children.

Aches, Fever ● Aspirin (NSAID) 325 mg

Antacid ● Citric Acid 1000 mg Sodium Bicarbonate 1916 mg

72 Tablets

(36 x 2)

Tamper Evident Unit Dose Packets

Medique® Medi-Seltzer
Effervescent Antacid/Pain Reliever

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

Collect Medibucks from certain large size Medique Brand products and redeem them for great gifts visiting www.mediqueproducts.com

15 MEDIBUCKS

Medique® Medi-Seltzer
Effervescent Antacid/Pain Reliever

Pull to Open

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

72 Tablets (36 x 2)

Tamper Evident Unit Dose Packets

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin (NSAID) 325 mg _____ Pain reliever/fever reducer
Citric Acid 1000 mg _____ Antacid
Sodium bicarbonate (heat-treated) 1916 mg _____ Antacid
Nonsteroidal anti-inflammatory drug

Uses
For the temporary relief of:
● heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
● upset stomach with headache from overindulgence in food or drink
● headache, body aches, and pain alone

Warnings
Riley's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Riley's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:
● hives ● facial swelling
● asthma (wheezing) ● shock
If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Stomach Bleeding Warning: This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you:
● are age 60 or older
● have had stomach ulcers or bleeding problems
● take a blood thinning (anticoagulant) or steroid drug
● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
● have 3 or more alcohol drinks every day while using this product
● take more or for a longer time than directed

Do not use
● if you have ever had an allergic reaction to any other pain reliever/fever reducer
● if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if
● stomach bleeding warning applies to you
● you have a history of stomach problems such as heartburn
● you have high blood pressure, heart disease, liver disease, or kidney disease
● you are taking a diuretic
● you are on a sodium restricted diet
● you reached age 60 or older
(continued on the next panel)

Directions
2 tablets every 4 hours. Do not exceed 8 tablets in 24 hours or do not use the maximum dosage for more than 10 days.

Children under 12 years of age: Do not give to children under 12 years of age.

60 years and older: Do not exceed 4 tablets in 24 hours or do not use the maximum dosage for more than 10 days.

Other information
● each tablet contains: Sodium 524mg
● store at room temperature 59°-86° F (15°-30° C)
● read all product information before using
● avoid excessive heat and humidity
● tamper evident sealed packets
● do not use any opened or torn packets

Inactive ingredients None

Questions or comments: 1-800-634-7680

Retain carton for complete product information

Reorder #13524
Manufactured for Medique Products
Fort Myers, FL 33967
1-800-634-7680
www.mediqueproducts.com

3 47682 13524 9

MEDIQUE MEDI SELTZER

aspirin, citric acid, sodium bicarbonate tablet, effervescent

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:47682-135 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---------------------------------------------------------------------------------------------|-----------------------|----------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 325 mg |
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL) | ANHYDROUS CITRIC ACID | 1000 mg |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20) | SODIUM BICARBONATE | 1916 mg |

Product Characteristics

| | | | |
|-----------------|---------------|---------------------|----------|
| Color | white (white) | Score | no score |
| Shape | ROUND (ROUND) | Size | 25mm |
| Flavor | | Imprint Code | L421 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--------------------------------------------------|----------------------|--------------------|
| 1 | NDC:47682-135-20 | 18 in 1 BOX | 12/30/2008 | |
| 1 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 2 | NDC:47682-135-24 | 36 in 1 BOX | 12/30/2008 | |
| 2 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 3 | NDC:47682-135-60 | 300 in 1 BOX | 12/30/2008 | |
| 3 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| OTC Monograph Drug | M013 | 12/30/2008 | |

Labeler - Unifirst First Aid Corporation (832947092)

