

ISOPROPYL ALCOHOL- isopropyl alcohol liquid
Medline Industries, LP

800 Isopropyl Alcohol

Active ingredient

Isopropyl Alcohol 70%

Purpose

First Aid Antiseptic

Uses

For first aid to help prevent the risk of infection in minor:

- cuts
- scrapes
- burns

Warnings

For external use only.

Flammable. Keep away from fire or flame.

Do not use

with electrocautery procedures. Use only in a well ventilated area; fumes may be toxic

When using this product do not

- get into eyes
- apply over large areas of the body. In case of deep or puncture wounds, animal bites or serious burns, consult a doctor.

Stop use and ask a doctor if

condition persists or gets worse or lasts for more than 72 hours. Do not use longer than 1 week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Clean the affected area. Apply a small amount of the product on the area 1 to 3 times daily.
- May be covered with a sterile bandage.

- If bandaged, let dry first.

Other Information

- Does not contain, nor is intended as a substitute for grain or ethyl alcohol.
- Will produce serious gastric disturbances if taken internally.
- Protect from freezing and avoid excessive heat.

Inactive ingredients

Water

Package/Label Principal Display Panel

MEDLINE

NDC: 53329-800-06

70%

ISOPROPYL ALCOHOL

First aid antiseptic

WARNING FLAMMABLE Keep away from heat, spark, electrical fire or flame.

Product sealed for your protection.
Use only in a well-ventilated area; fumes can be harmful.

16 FL OZ
(1 PT) 473 mL

L0015301FA

Drug Facts	
Active ingredient	Purpose
Isopropyl alcohol 70%.....	First aid antiseptic
Uses first aid to help prevent the risk of infection in: ■ minor cuts ■ scrapes ■ burns	
Warnings For external use only	
Flammable ■ keep away from fire or flame, heat, spark, electrical	
Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns	
When using this product ■ do not get into eyes ■ do not inhale ■ do not apply over large areas of the body ■ do not use longer than 1 week	
Stop use and ask a doctor if condition persists or gets worse	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Caution – Fumes can be acutely irritating to skin, eyes and the respiratory system. Do not apply to irritated skin or if excessive irritation develops. Avoid getting into the eyes or on mucous membranes. Avoid inhaling this product.	
Directions ■ clean the affected area ■ apply a small amount of this product on the area 1 to 3 times daily ■ may be covered with a sterile bandage ■ if bandaged, let dry first	
Other information ■ does not contain, nor is intended as a substitute for grain or ethyl alcohol ■ will produce serious gastric disturbances if taken internally	
Inactive ingredient water	

REF MDS098003Z

www.medline.com Made in US with US and foreign components for Medline Industries, LP, Three Lakes Drive, Northfield, IL 60093 USA. 1-800-MEDLINE V1 RJ24VJO L0015301BA

0 80196 74864 1

ISOPROPYL ALCOHOL

isopropyl alcohol liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53329-800
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	70 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53329-800-06	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/16/2009	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	02/16/2009	

Labeler - Medline Industries, LP (025460908)

Registrant - Medline Industries, LP (025460908)

Revised: 12/2024

Medline Industries, LP