# GOOD SENSE COMPLETE LICE TREATMENT KIT- piperonyl butoxide, pyrethrum extract L. Perrigo Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### **Perrigo Complete Lice Treatment Kit Drug Facts**

#### **Active ingredients**

Piperonyl butoxide 4%

Pyrethrum extract (equivalent to 0.33% pyrethrins)

#### Purpose

Lice treatment

#### Uses

treats head, pubic (crab), and body lice

#### **Warnings**

#### For external use only

#### Do not use

- near eyes
- inside nose, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

#### Ask a doctor before use if you are

allergic to ragweed. May cause breathing difficulty or an asthmatic attack.

#### When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets in eyes, flush with water right away
- scalp itching or redness may occur

#### Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

#### Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

 Important: Read warnings before use Adults and children 2 years and over:

#### Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

#### **Treat**

- shake well before using
- apply thoroughly to **DRY HAIR** or other affected area. For head lice, first apply behind ears and to back of neck.
- allow product to remain for 10 minutes, but no longer
- use warm water to form a lather, shampoo, then thoroughly rinse
- for head lice, towel dry hair and comb out tangles

#### Remove lice and their eggs (nits)

- use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove).
- hair should remain slightly damp while removing nits
- if hair dries during combing, dampen slightly with water
- for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours.
- lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.
- pin back each strand of hair after combing
- clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
- check daily for any lice/nits that you missed
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice
- if infestation continues, see a doctor for other treatments

#### **Children under 2 years:** ask a doctor

#### Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat
- keep carton for important product information
- see Consumer Information Insert for additional information

#### **Inactive ingredients**

benzyl alcohol, C13-C14 isoparaffin, fragrance, polyoxyethylene (1,1,3,3-tetramethyl butyl) phenyl ether, purified water, xanthan gum

#### **Questions & comments?**

1-800-719-9260

#### **Consumer Information**

3-STEP PROCESS FOR LICE TREATMENT

See carton for complete product information.

#### Step 1 ● Treat

## **Apply Lice Killing Shampoo**

See Usage Guidelines to estimate how much product you will need.

#### NOTE:

- Shake well before using.
- Use towels to protect child's eyes from treatment and prevent clothes from getting wet.
- Apply Lice Killing Shampoo thoroughly to **DRY HAIR** or other affected area.
- Wetting the hair dilutes the treatment making it less effective.
- Also, lice can hold their breath when immersed in water making it difficult for the active ingredient to penetrate the lice.



• For head lice, thoroughly massage shampoo into scalp, behind the ears and to the back of the neck. Lice can crawl up and down the shaft of the hair very quickly so it is important to apply the treatment from the roots to the ends of the hair.



• Allow product to remain on the hair (or other affected area) for 10 minutes, but no longer.



- Add warm water and massage to form a lather.
- Rinse thoroughly.



- For head lice, towel dry hair and comb out tangles.
- Proceed to Step 2.
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice.

	Hair	Short		Medium		Long	
Guidelines	Length	(ear length	or shorter)	(shoulder le	ength)	(past should	ler length)
	Approximate	First	1  oz - 2  oz	First	2 oz - 3 oz	First	3  oz - 4  oz
	Amount for 1	Application	n:	Application	<b>.</b>	Application	:
	Adult/Child	Second	1  oz - 2  oz	Second	2  oz - 3  oz	Second	3  oz - 4  oz
		Application	n:	Application	• •	Application	:
		Total:	2 oz – 4 oz	Total:	4 oz – 6 oz	Total:	6 oz – 8


#### Step 2 • Remove

# Comb out lice and eggs with lice comb or other fine-tooth comb and Lice & Egg Comb-Out Spray


This product should be used **AFTER** Lice Killing Shampoo (Step 1). Combing is an important step to remove eggs.

#### NOTE:

- A fine-toothed comb or the special lice/nit removing comb (included) must be used to remove lice, eggs, and nits from hair.
- Before starting have on hand:
  - extra towels
  - bobby pins or hair clips
  - facial tissues or paper towels
  - regular comb for parting and detangling hair
  - throw-away gloves
- Find a distraction for your child (e.g. books, TV, game).
- Comb out damp hair with a regular wide-tooth comb to take out snarls and tangles.



Apply a small amount of Lice & Egg Comb-Out Spray and use a wide-tooth comb to remove any tangles.



Lift a 1-2" section of hair. Start as close to scalp as possible, slowly comb section all the way to ends of hair with lice comb, using additional Lice & Egg Comb-Out Spray to remove tangles as needed. Use a tissue or paper towel to wipe any lice or eggs off comb.



Pin back this section of hair. Continue to comb out one 1-2" section at a time until entire head has been combed and pinned.



Remove pins and rinse thoroughly with warm water. You may use your regular shampoo now if desired. Combs should be disinfected for reuse by soaking in hot water (above 130°F) for 10 minutes.

#### Step 3 • Control

#### Treat the home with Home Lice, Bedbug & Dust Mite Spray

Use Home Lice, Bedbug & Dust Mite Spray to kill lice and their eggs on mattresses, furniture, car interiors and other non-washable items.

#### Not for use on humans.

#### **Additional Tips for Success**

- Be sure to complete all three steps in the Lice Treatment process. This includes a second shampooing between 7-10 days after the initial treatment, but no sooner. If any eggs were not initially combed out, lice may hatch and can be killed with the second treatment. Newly hatched lice are not mature enough to lay new eggs.
- Carefully inspect infested family members daily for at least 2 weeks after the second shampooing. If you see eggs or nits, consult your doctor.
- Teach children not to share personal items, such as hats, combs and brushes and to avoid

prolonged contact with a person who has lice.

#### Did You Know?

#### **HEAD LICE**

- Lay small white eggs (nits) on hair shaft close to scalp.
- Nits are most easily found on back of neck or behind ears.
- Disinfect hats, hair ribbons, scarves, coats, towels, and bed linens by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes.
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be drycleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again.
- Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose.
- Soak all combs and brushes in hot water (above 54°C (130°F)) for at least 10 minutes.
- Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.

#### **PUBIC (CRAB) LICE**

- May be transmitted by sexual contact. Sexual partners should be treated simultaneously to avoid reinfestation.
- Lice are very small and look like brown or grey dots on skin.
- Usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface.
- May be present on the short hairs of groin, thighs, trunk, and underarms, and occasionally on the beard and mustache.
- Disinfect underwear by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes.

#### **BODY LICE**

- Body lice and their eggs (nits) are generally found in the seams of clothing particularly in waistline and armpit area.
- Body lice feed on skin then return to clothing to lay their eggs.
- Disinfect clothing by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes.
- Do not seal clothing in a plastic bag because nits can remain dormant for up to 30 days.

#### Save this insert for future reference

Distributed By

Perrigo®

Allegan, MI 49010

:6Y600 00 J1

#### Keep carton for important product information.

Lice Killing Shampoo

**Drug Facts** 

#### **Active ingredients**

Piperonyl butoxide 4%

Pyrethrum extract (equivalent to 0.33% pyrethrins)

#### **Purpose**

Lice treatment

#### Uses

treats head, pubic (crab), and body lice

#### Warnings

#### For external use only

#### Do not use

- near eyes
- inside nose, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

**Ask a doctor before use if you are** allergic to ragweed. May cause breathing difficulty or an asthmatic attack.

### When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets in eyes, flush with water right away
- scalp itching or redness may occur

#### Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

Important: Read warnings before use

#### Adults and children 2 years and over:

#### Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

#### **Treat**

- shake well before using
- apply thoroughly to **DRY HAIR** or other affected area. For head lice, first apply behind ears and to back of neck.

- allow product to remain for 10 minutes, but no longer
- use warm water to form a lather, shampoo, then thoroughly rinse
- for head lice, towel dry hair and comb out tangles

#### Remove lice and their eggs (nits)

- use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw- away glove).
- hair should remain slightly damp while removing nits
- if hair dries during combing, dampen slightly with water
- for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours.
- lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.
- pin back each strand of hair after combing
- clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
- check daily for any lice/nits that you missed
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice
- if infestation continues, see a doctor for other treatments

#### **Children under 2 years:** ask a doctor

#### Other information

• store at 20-25°C (68-77°F). Avoid excessive heat.

**Inactive ingredients** benzyl alcohol, C13-C14 isoparaffin, fragrance, polyoxyethylene (1,1,3,3-tetramethyl butyl) phenyl ether, purified water, xanthan gum

Questions? 800-719-9260

#### Lice & Egg Comb-Out Spray

#### **Product Information**

**Use** detangles hair in order to make lice and egg comb-out easier

#### **Warnings**

#### For external use only. Keep out of reach of children. Avoid contact with eyes.

#### Directions for use

- use AFTER shampooing with a lice killing shampoo
- towel dry hair. Apply a small amount of this product and use a wide-toothed comb to remove any tangles.
- comb out the dead lice, eggs and nits, one small section at a time, with the enclosed lice comb. **This step is very important.**
- use additional product to remove tangles, if needed, during the comb-out process
- after you have combed entire head, rinse hair thoroughly with warm water. Disinfect combs with hot water (130°F).

**Ingredients** purified water, polysorbate 20, glycerin, propylene glycol, phenoxyethanol, amodimethicone, fragrance, DL-panthenol, trideceth-12, cetrimonium chloride,

methylchloroisothiazolinone, methylisothiazolinone

#### Home Lice, Bedbug & Dust Mite Spray

Not for use on humans.

#### **ACTIVE INGREDIENT:**

100.00%

†Cis/trans ratio: Max. 55% (+/-) cis and Min. 45% (+/-) trans.

††Contains petroleum distillate.

#### **FIRST AID**

#### If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

#### If swallowed:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

#### If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-222-1222.

#### NOTE TO PHYSICIAN

Contains petroleum distillate – vomiting may cause aspiration pneumonia.

#### PRECAUTIONARY STATEMENTS

#### **Hazards To Humans & Domestic Animals**

**CAUTION:** Harmful if absorbed through skin. Avoid inhalation of spray mist. Causes moderate eye

irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco or the toilet. Remove and wash contaminated clothing before reuse.

#### PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54°C) may cause bursting.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **Use Restrictions:**

- Do not allow adults, children, or pets to enter until vapors, mists and aerosols have dispersed, and the treated area has been thoroughly ventilated.
- Do not allow adults, children, or pets to enter the treated area until sprays have dried.
- In the home, cover all food processing surfaces and utensils during treatment and thoroughly wash before use. Cover or remove exposed food.
- Remove pets, birds and cover any water inhabited by fish (such as aquariums and ornamental fish ponds) before application. Turn off aquarium filtering systems before spraying and wait approximately 15 minutes after application before turning the filter back on.
- **This product is not for use on humans.** If lice infestations should occur on humans, use a product labeled for use on humans.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
- Not for broadcast use on indoor residential surfaces.

#### SHAKE WELL BEFORE USING

Remove protective cap. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 12 to 15 inches for baseboards, moldings and when applying to floors as a spot spray and 8 to 10 inches for clothing, furniture and bedding. Test product on an inconspicuous area before applying to check for possible staining or discoloration. Inspect again after drying. Spray to the point of dampness but not run-off. Over application may cause damage. Will not stain surfaces or fabrics where water alone causes no damage.

To kill *Lice* and *Louse eggs*: Clean home. Use this product for non-washable items only. Washables should be washed in hot water and dried in high heat. Spray only those articles and parts of bedding, including mattresses and furniture that cannot be either laundered or dry-cleaned. Do not use on sheets or pillowcases. Allow all sprayed articles to dry thoroughly before use. If lice infestation occurs on humans, use a product labeled for use on humans.

**Bedbugs**: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Spray to the point of dampness but not run-off. After mattress is dry, cover with mattress cover and sheet. Do not use mattress without cover. Retreat if infestation occurs, but not more than once every two weeks.

To control *House Dust Mites (Dermatophagoides spp.)*: This product controls house dust mites that may cause asthma, hay fever, rhinitis, watery eyes and sneezing. One treatment lasts for up to 8 weeks. **On Uphols tered Furniture** – Test fabric in an inconspicuous area to insure dyes will not bleed. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray all fabric surfaces until damp. Spray to the point of dampness but not run-off. Do not use on delicate fabrics. Allow furniture to dry thoroughly before using. **On Carpets** – Apply as a spot spray to areas where dust mites are likely (including along baseboards and moldings). Hold nozzle

approximately 12 to 18 inches from the carpet surface. Spray until damp. After application, brush the surface of the carpet with a broom or brush to insure deep penetration. Allow carpet to dry thoroughly before reentering the treated area. Drying time can be accelerated by using a fan. **On Mattresses** — Spray the top, bottom and sides of mattresses until damp, paying particular attention to tufts and seams. Allow to dry thoroughly before using. After mattress is dry, cover with mattress cover and sheet. Do not use mattress without cover. Vacuum entire area once dry. Since vacuuming stirs dust up in the air those with allergies should have another person vacuum or should wear a dust mask and use vacuum with a high efficiency particulate (HEPA) filter.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry area away from children and pets and away from heat, sparks and open flame. Do not transport or store below 32°F (0°C). Protect from freezing. Do not freeze.

**Pesticide Disposal and Container Handling:** Do Not Puncture or Incinerate.

**If empty:** Place in trash or offer for recycling if available.

**If partly filled:** Call your local solid waste agency or 1-800-CLEANUP (253-2687) for disposal instructions. Never place unused product down any indoor or outdoor drain.

**To the extent consistent with applicable law,** purchaser assumes all responsibility for safety and use not in accordance with directions.

See carton for EPA numbers.

#### Package/Label Principal Display Panel

Maximum Strength Shampoo

Complete Lice Treatment Kit

3 Step Lice Control System

STEP 1 Lice Killing Shampoo

Proven effective treatment

Leaves no chemical residue behind

STEP 2 Lice & Egg Comb-Out Spray

Makes combing faster and easier

Combing is an essential step to remove eggs

The included comb is 100% effective in the removal of eggs and lice from hair♦

♦As demonstrated in laboratory testing

STEP 3 Control In-Home Lice, Bedbug & Dust Mite Spray

Kills lice and their eggs on mattresses, furniture, car interiors and other non-washable items

KEEP OUT OF REACH OF CHILDREN


#### **CAUTION:**

See back and side panel for additional information

Compare to active ingredients of RID® Lice Killing Shampoo

100% MONEY BACK GUARANTEE

NET 4 FL OZ (118 mL) SHAMPOO - NET WT 5 OZ (142 g) SPRAY



**Maximum Strength Shampoo** 

# Complete Lice Treatment Kit

3 Step Lice Control System

STEP 1 Lice Killing Shampoo

- Proven effective treatment
- •Leaves no chemical residue behind

STEP 2 Lice & Egg Comb-Out Spray

- Makes combing faster and easier
- Combing is an essential step to remove eggs

The included comb is 100% effective in the removal of eggs and lice from hair +

STEP 3 Control In-Home Lice, Bedbug & Dust Mite Spray

 Kills lice and their eggs on mattresses, furniture, car interiors and other non-washable items

KEEP OUT OF REACH OF CHILDREN

CAUTION:

See back and side panel for additional information

Compareto active ingredients of RID® Lice Killing Shampoo



\*As demonstrated in

NET 4 FL OZ (118 mL) SHAMPOO • NET WT 5 OZ (142 g) SPRAY NET 2 FL OZ (59 mL) COMB-OUT SPRAY • ONE NIT COMB

Complete lice control for your family and home

#### STEP 1 TREAT

- Apply Lice Killing Shampoo according to label directions.
   Report this step 7 to 10 days letter to help prevent.
- Repeat this step 7 to 10 days later to help prevent reinfestation.

#### STEP 2 REMOVE


- After Step 1, comb out the eggs and nits in the hair with a fine-toothed comb or with the comb included.
- Apply Lice & Egg Comb-Out Spray on damp hair to make removal faster and easier.

#### STEP 3 CONTROL

- Use Home Lice, Bedbug & Dust Mite Spray to kill lice and their eggs on mattresses, furniture, car interiors and other non-washable items.
- Wash bed linens, clothing and other items in hot water and dry in high heat.

This kit contains 3 products for complete lice control for your family and home. The 4 oz bottle of shampoo and 2 oz bottle of combing spray should be enough product to treat one person. However, amount of productneeded will vary by hair length. See below for guidelines.

Lice Killing S	hampoo		
Hair Length	Approximate Amount For 1 Adult/Child:		
Short (ear length	First Application: Second Application:	1 oz - 2 oz 1 oz - 2 oz	
or shorter)	Total:	2 oz - 4 oz	
Medium (shoulder length)	First Application: Second Application:	2 oz - 3 oz 2 oz - 3 oz	
20000	Total:	4 oz - 6 oz	
Long (past shoulder	First Application: Second Application:	3 oz - 4 oz 3 oz - 4 oz	
length)	Total:	6 oz - 8 oz	



## 3 Step Lice Control System

# Completel

## COODSENSE<sup>®</sup>

#### Lice Killing Shampoo

Drug Facts	
Active ingredients	Purpose
Piperonyl butoxide 4%Lice Pyrethrum extract	treatment
	treatm ent

#### Uses treats he ad, pubic (crab), and body lice

## Warnings

Do not use in near eyes in inside nose, mouth, or vagina mon lice in eyebrows or eyelashes. See a doctor if lice are present in the se areas.

Ask a doctor before use if you are allergic to regweed. May cause breathing difficulty or an authorized attack.

When using this product in keep eyes bightly doesdand protect eyes with a washcloth or lower

n scalp riching or redness may occur nif product gets in eyes, flush with water right away

Stop use and sales doctor if meye imbation occurs white thing difficulty occurs makinor acalpi mbation continues or infection occurs Keep out of reach of children. Hawallowed, get medical ta Poison Control Centerright away help or contacts: Por (1-800-222-1222).

Directions
Important: Read warnings before use
Adults and children 2 years and over:

laspect • check each household member with a magnifying glass in

brightlight for licelnits (eggs) wlook for in y nits near scalp, beginning at back of neck and

behind ears necemine small sections of hair statime numble dandruff which moves when bouched, nits stick to

the hair mili either lice or nits are found, treat with this product

Ingite well before using market within process.

Ingite well before using market reflected area. For feed loe, first apply behind ass and to back of reds. allow product to remain for 10 minutes, but no longer more warm water to form a larker, shampoo, from suce warm water to form a larker, shampoo, from

floroughly rinse in for headlice, towel dry hair and comb out tangles

wite hacilities, towed dy hair and comb outstangles. Remove lice and their eggs (pith) muses a fine-both or special lookint comb. Remove any remaining mits by hard busing a frice-wavey glove). What is should remain slightly drap white removing mits with the dies during controling, dumper slightly with water in the had look, part their into sections. Do one section at a time starting on top of head. Longer heir any take 1 to 2 hours. will at 1-bi2-in-third de strand of hair. Place comb as close to salp as possible and comb with a firm, even motion savey from a colp.

away mon acap. mpinbaskeach strand of heir after combing mdian comb often. Wee nits away with tissue and discard in a plestic bag. Seal bag and discard to prevent lice from components.

matter combing, thoroughly recheck for lice/hits. Repeat

combing fine-sessay.

In check daily for any lice/hits that you missed

a second trustment must be done in 7 to 10 days to kill any

Drug Facts (continued)

milimestation continues, see a diorior for other treatments Children under 2 years: ask a doctor

Other information

store at 20-25°C (68-77°F)

nikeep carton for important product information nisee Consumer Information Insert for additional information

Inactive ingredients beraylalcohol, C13-C14 isopartiin, fagrance, polycoyethylene (1, 1, 3, 3-letrametry) butyliphenyl ether, purihed water, vanthan gum

Questions or comments ?1 -800-719-9260

#### Lice & Egg Comb-Out Spray

Product Information

Use detangles hair in order to make lice and egg comb-out

Warnings For external use only. Keep out of reach of children. Avoid contact with eyes.

Directions for us

complete directions
use ATER shampcoing with all icelelling shampco
I towel dry hair. Apply a small amount of this product and use a
wide-bothed combinerance any langles.

comb out the dead fice, eggs and nits, one small section at a time, with the enclosed lice comb. (See insert for additional

combing instructions). This step is very important. ups additional product to remove tangles, if needed, during

the comb-out process

after you have combed entire head, rinse hair thoroughly w
warm water. Disinfect combs with hot water (130 °F).

Ingredients purified water, polysorbate 20, glycerin, propylene glycol, phenoxyethanol, amedine thiopre, traipance propylene glycol, phenosyethanol, amodimetricone DL-panthenol, trideceth-12, entrimonium chloride, methyl chloroi softiazoli rone, methyl isothiszoli rone

Questions or comments ?1 -800-719-9260

#### Home Lice, Bedbug & Dust Mite Spray Not for use on humans.

### Permetrin OTHER INCREDENTS: ...0.50% ...99.50% 100.00% †Cis/transinstic: Max. 55% (+/-) dis and Min. 45% (+/-) trans. ††Contains petroleum distillate.

Take of contaminated dofting. Pinse disnimmed stoly with plenty of water for 15-20 minutes.

Call a poison control senter or doctor for

trustment advice. Holdeye open andrinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first

5 minutes, then confinue mising eye.

Call a poison control center or declor for

Immediately call a poison control center or doctor.

Do not induce vornifing unless tolet to do so by a poison control center or doctor.

Do not give any liquid to the person. Do not give anything by mouth to an uncons

Move person to freshair.
If person is not breathing, sall 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

Have the product container or label with you when calling a poision control center or doctor, or going for treatment. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may sall 1-800-222-1222.

Contains petroleum distillate - vomiting may cause aspiration

FIRM: Harmful if absorbed through skin. Avoid inhalation of spray mist. Causes moderate eye initation, Avaid contact with skin, eyes or clothing. Wash thoroughly with seap and water after handing and before eating, drinking, chewing gum, or using tobacco or the toilet. Pernove and wash conta dofting before reuse.

Contents under pressure. Do not use or store near heat or open fame. Denot puncture or incinerate container. Exposure temperatures above 130°F (64°C) may cause bursting.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- one restructions:

  Constallow soldts, children, or prict trenter until vapora, mists and servosiolistive dispersed, and the treated direction been thoroughly verificated.

  Constallow soldts, children, or prict trenter the treated areas until graps have direct.

  In the home, cover all lood processing surfaces, and utencils during treatment and thoroughly weatherfore use. Cover or remove exposed food.

  Represented to the children of the control of the co

- Remove pets, birds and eover any water inhabited by fish (such as aquariums and ornamental fish pondis) befor application. Turn of aquarium filtering systems before spraying and wait approximately 15 minutes after application before turning the filter back on.
- This product is not for use on humans. If fice infestations should occur on humans, use a product lebeled for use onhumans.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, atorm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. • Not for broadcast use on indoor ruside nital surfaces

Remove protective cap, Bald container up gight with needs sway from you. Dupness whe and apray from a distance of 12 to 15 inches for Doueboards, makings and when applying to floors are spectappy and 8 to 10 inches for cathing, farmings and bedding, later product on an incompisuous area before applying to check for possible staming or disostonation, haper Lagan after drying. Spray to the point of data posses but not uncell. Over application may cause damage. Will not stain surfaces or fabrics where water alone causes no demonstrate.

Clean home. Use this product for mon-washable ichtif Levend Lauwenger: Gean home, bei für productfür non-washelbeimen ofly Washelbeimholdes wesseld in hotweis and dried in high beit. Spray only hose arficks and ports of bedding, induding mattresses and furnium that carnot be either launden der dry-skannel. De not use on she bei prijewasses. Albe will sprayed artisks be dry throughly before use. If lies infestation ceeus on humans, use a product labeled für use on humans.

Bedfuge: Spray mathesises, particularly around fuffs and seems. Take beds apart and agray into all pints. Their baseboards, moldings and flows. Spray to the paint of dampinass but notion-off. After notifiees to dy, cover with maintess over and sheet. Do not use nathesis without cover. Retmatrif infestation occurs, but not note than once every howevers.

To kill Riosee Dust Mikes (Demostophagodieu app.): This product controls house dust mins that may cause authma, hey fever, rhinitis, watery eyes and needing. The restort ent last form to be week. On Uphobitised Furniture—Test faint cings incorporation are see to incure dyes will not be all hald contains using that when way from you. Uppers when and parry from a distance of 8 to 10 inches. Spray all listine authors until damp. Spray for the point of dampness but not turn off. Do not use and elected faintes. Allow marker to but from coglitic testing allowed the seed of the point of dampness but not turn of Do not use and elected faintes. Allow marker to but from coglitic testing, the Campels—Apply as a supplying reliable for the sample of the cappet within a post purpose. Spray until damp, After application, but allowed the cappet with a broom or brush in such companies of the cappet that a broom or brush to have one prevention at the cappet to the cappet that become or brush in such as the cappet to the cappet that a broom or brush to have a dust the sample of the cappet that a cappet that the cappet t

Bo noticentaminate water, load or each by storage or disposal.

Perficiels Storage: Store in a cool, dry area away from children and pets and away from heat, pare is and op inflame. Do not than apport or store below 22°F, PCQ, Portest: from becamp, to not than apport or store below 22°F, PCQ, Portest: from becamp, to not the reach perfect to the perfect of the performance of the performance

NUTICE: for the extent consistent with applicable law, purchaser assume all responsibility for safety and use not in assordance with directions.

EPA Reg. No. 71-1 EPA Est. No. 89209-MI-001

MPORTANT: Read detailed directions inside before using

Distributed By L. Perrigo Co. 515 Eastern Ave, Allegan, MI 49010

MONEY BACK GUARANTEE



### GOOD SENSE COMPLETE LICE TREATMENT KIT

piperonyl butoxide, pyrethrum extract kit

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0113-0108

#### **Packaging**

# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:0113-0108-62	1 in 1 CARTON; Type 0: Not a Combination Product	11/20/2018	

#### **Quantity of Parts**

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	118 mL
Part 2	1 BOTTLE, PUMP	59 mL

#### Part 1 of 2

#### LICE KILLING

piperonyl butoxide, pyrethrum extract shampoo

#### **Product Information**

Route of Administration TOPICAL

#### **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
(	PIPERONYL BUTOXIDE	4 g in 100 mL
	PYRETHRUM EXTRACT	.33 g in 100 mL

#### Inactive Ingredients

macave ingredients			
Ingredient Name	Strength		
BENZYL ALCOHOL (UNII: LKG8494WBH)			
C13-14 ISOPARAFFIN (UNII: E4F12ROE70)			
WATER (UNII: 059QF0KO0R)			
XANTHAN GUM (UNII: TTV12P4NEE)			

#### **Product Characteristics**

Color YELLOW (cloudy to transparent, cream to straw) Score

Shape	Size	
Flavor	Imprint Code	
Contains		

l	Packaging			
ı	# Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
ı	1	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part358G			

## Part 2 of 2

## LICE AND EGG COMB OUT

other hair preparations

### **Product Information**

Route of Administration TOPICAL

Other Ingredients				
Ingredient Kind	Ingredient Name	Quantity		
INGR	WATER (UNII: 059QF0KO0R)			
INGR	POLYSORBATE 20 (UNII: 7T1F30V5YH)			
INGR	GLYCERIN (UNII: PDC6A3C0OX)			
INGR	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
INGR	PHENOXYETHANOL (UNII: HIE492ZZ3T)			
INGR	PANTHENOL (UNII: WV9CM0O67Z)			
INGR	TRIDECETH-12 (UNII: YFY3KG5Y7O)			
INGR	CETRIMO NIUM CHLO RIDE (UNII: UC9 PE95IBP)			
INGR	METHYLCHLOROISOTHIAZOLINONE (UNII: DEL7T5QRPN)			
INGR	METHYLISOTHIAZOLINONE (UNII: 229 D0 E1QFA)			

l	Packaging				
	#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
	1		59 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product		

	Marketing Information				
ı	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	

Cosmetic					
Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part358G	11/20/2018			

## Labeler - L. Perrigo Company (006013346)

Revised: 2/2019 L. Perrigo Company