

IODINE PRO CONCENTRATE- iodine solution, concentrate
Aspen Veterinary

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

Attention

For External Use Only

Wash hands thoroughly after handling.

Get medical advice/attention if you feel unwell.

FOR INDUSTRIAL USE ONLY

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: Contact local poison control.

DIRECTIONS:

IODINE PRO CONCENTRATE MUST BE DILUTED BEFORE USE

DO NOT MIX WITH ANYTHING BUT COOL, POTABLE WATER

FOR USE ONLY IN A DISPENSING SYSTEM RECOMMENDED BY ECOLAB

Dilution Parameter(s):

Percent Iodine	Iodine Pro Concentrate By Part	Water By Part	Final RTU Product
1.00%	1	3	4
0.75%	1	4.30	5.30
0.50%	1	7	8
0.25%	1	15	16

IMPORTANT: Do not mix Iodine Pro Concentrate or diluted Iodine Pro Concentrate with any other products unless specified in the product use instructions. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for Iodine Pro Concentrate. If product in dip cup becomes visibly dirty, discard contents and replenish with diluted, ready-to-use product. Do not reuse or return any unused product to the original container.

Before Milking: Wash teats thoroughly just prior to next milking with appropriate udder wash solution or pre-milking teat dip to avoid contamination of milk. Teats should then be dried with single-service towels. Use proper procedures for udder washing or pre-dipping.

Application of Diluted Iodine Pro Concentrate

Directions for Teat Dipping

Pre-Milk Dipping: Fill teat dip cup with diluted Iodine Pro Concentrate. Before each cow

is milked, dip each teat full-length into the teat dip cup containing diluted Iodine Pro Concentrate. Wipe teats dry after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Immediately after each milking, dip each teat full-length into the teat dip cup containing diluted Iodine Pro Concentrate. Allow to air dry. Do not wipe. Always use fresh, diluted Iodine Pro Concentrate. Do not turn cows out in freezing weather until teat dip is completely dry.

Directions for Teat Spraying

Pre-Milk Spraying: Before each cow is milked, spray entire teat with diluted Iodine Pro Concentrate. Wipe teats after spraying using single-service towels to avoid contamination of milk.

Post-Milk Spraying: Immediately after each milking, spray entire teat with diluted Iodine Pro Concentrate. Allow to air dry. Do not wipe. Always use fresh, diluted Iodine Pro Concentrate. Do not turn cows out in freezing weather until diluted Iodine Pro Concentrate is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.

PRECAUTION: Iodine Pro Concentrate is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

NOTE: Use blended Iodine Pro Concentrate within six (6) months of blended date.

Principal display panel and representative container

ASPEN IODINE PRO CONCENTRATE

ACTIVE INGREDIENT:

Titrateable iodine.....4.0%

INERT INGREDIENTS:..... 96.0%

(contains glycerin, propylene glycol)

TOTAL:.....100.0%

Iodine Teat Dip Concentrate

An aid in reducing the spread of organisms which may cause mastitis

208 L (55 US GAL)

6302354

Manufactured for:

Aspen Veterinary Resources®, Ltd. 3155 Heartland Drive, Liberty, MO 64068 1-800-203-5620

778199/5300/1021



IODINE PRO CONCENTRATE



ACTIVE INGREDIENT:	
Titratable Iodine	4.0%
INERT INGREDIENTS:	96.0%
(contains glycerin, propylene glycol)	
TOTAL:	100.0%

Iodine Test Dip Concentrate
 An aid in reducing the spread of organisms which may cause mastitis
FOR INDUSTRIAL USE ONLY
DIRECTIONS:
IODINE PRO CONCENTRATE MUST BE DILUTED BEFORE USE
DO NOT MIX WITH ANYTHING BUT COOL, POTABLE WATER
FOR USE ONLY IN A DISPENSING SYSTEM RECOMMENDED BY ECOLAB
Dilution Parameters:

Percent Iodine	Iodine Pro Concentrate By Part	Water By Part	Final RTU Product
1.00%	1	3	4
0.75%	1	4.30	5.30
0.50%	1	7	8
0.25%	1	15	16

IMPORTANT: Do not mix Iodine Pro Concentrate or diluted Iodine Pro Concentrate with any other products unless specified in the product use instructions. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for Iodine Pro Concentrate. If product in dip cup becomes visibly dirty, discard contents and replenish with diluted, ready-to-use product. Do not reuse or return any unused product to the original container.

Before Milking: Wash teats thoroughly just prior to next milking with appropriate udder wash solution or pre-milking teat dip to avoid contamination of milk. Teats should then be dried with single-service towels. Use proper procedures for udder washing or pre-dipping.
Application of Diluted Iodine Pro Concentrate
Directions for Teat Dipping
Pre-Milk Dipping: Fill test dip cup with diluted Iodine Pro Concentrate. Before each cow is milked, dip each teat full-length into the test dip cup containing diluted Iodine Pro Concentrate. Wipe teats dry after dipping using single-service towels to avoid contamination of milk.
Post-Milk Dipping: Immediately after each milking, dip each teat full-length into the test dip cup containing diluted Iodine Pro Concentrate. Allow to air dry. Do not wipe. Always use fresh, diluted Iodine Pro Concentrate. Do not turn cows out in freezing weather until teat dip is completely dry.

Directions for Teat Spraying
Pre-Milk Spraying: Before each cow is milked, spray entire teat with diluted Iodine Pro Concentrate. Wipe teats after spraying using single-service towels to avoid contamination of milk.
Post-Milk Spraying: Immediately after each milking, spray entire teat with diluted Iodine Pro Concentrate. Allow to air dry. Do not wipe. Always use fresh, diluted Iodine Pro Concentrate. Do not turn cows out in freezing weather until diluted Iodine Pro Concentrate is completely dry.
Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.
PRECAUTION: Iodine Pro Concentrate is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.
NOTE: Use blended Iodine Pro Concentrate within six (6) months of blended date.

KEEP OUT OF REACH OF CHILDREN
NOT FOR HUMAN USE
Attention
 For External Use Only
 Wash hands thoroughly after handling.
 Get medical advice/attention if you feel unwell.
READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT
EMERGENCY HEALTH INFORMATION: Contact local poison control.

Yodo concentrado para el tratamiento de pezones
 Ayuda a evitar la proliferación de organismos que pueden causar mastitis
PARA USO INDUSTRIAL SOLAMENTE
INSTRUCCIONES:
IODINE PRO CONCENTRATE DILUIR ANTES DE USAR
NO MEZCLAR CON NADA QUE NO SEA AGUA POTABLE FRÍA
USAR ÚNICAMENTE EN UN SISTEMA DE DOSIFICACIÓN RECOMENDADO POR ECOLAB
Parámetro(s) de dilución:

Porcentaje de yodo	Iodine Pro Concentrate Por Parte	Agua Por Parte	Producto RTU final
1.00%	1	3	4
0.75%	1	4.30	5.30
0.50%	1	7	8
0.25%	1	15	16

IMPORTANTE: No mezcle Iodine Pro Concentrate o Iodine Pro Concentrate diluido con otros productos a menos que se indique en las instrucciones de uso del producto. Si el producto fuese transferido desde este envase a cualquier otro, asegúrese de que ese otro envase haya sido íntegramente prelavado y que contenga el etiquetado del envase de Iodine Pro Concentrate. Si el producto en el recipiente para realizar el baño está visiblemente sucio, deseché el contenido y rellene el recipiente con producto diluido y listo para usar. No vuelva a utilizar el mismo producto, ni vuelva a colocar el producto sin usar dentro del envase original.

Antes de ordeñar: Lave los pezones completamente justo antes de ordeñar con una solución de lavado apropiada para ubres o baño antiséptico para pezones previo al ordeño, o rocíe el pezón para evitar la contaminación de la leche. Posteriormente, los pezones se deben secar con toallas de servicio único o descartables. Utilice el procedimiento adecuado para el lavado o el prebañado antiséptico de las ubres.
Aplicación de dilución Iodine Pro Concentrate
Instrucciones para realizar el sellado del pezón
Sellado del pezón previo al ordeño: Llene el recipiente para el baño de pezones con Iodine Pro Concentrate diluido. Antes de ordeñar cada vaca, bañe toda la superficie del pezón en el recipiente que contiene Iodine Pro Concentrate diluido. Seque los pezones después de realizar el sellado, utilizando toallas desechables para evitar la contaminación de la leche.
Sellado del pezón posterior al ordeño: Inmediatamente después de ordeñar cada vaca, bañe toda la superficie del pezón en el recipiente que contiene Iodine Pro Concentrate diluido. Deje secar al aire. No utilice paños. Siempre utilice Iodine Pro Concentrate fresco y diluido. Hasta que Iodine Pro Concentrate no se haya secado completamente, no saque las vacas al exterior en climas fríos.

Instrucciones para el Rociado de los Pezones
Rociado previo al ordeño: Antes de ordeñar cada vaca, rocíe toda la superficie del pezón con Iodine Pro Concentrate diluido. Seque los pezones después de rociarlos utilizando toallas de un solo servicio para evitar la contaminación con la leche.
Rociado posterior al ordeño: Inmediatamente después de ordeñar cada vaca, rocíe toda la superficie del pezón con Iodine Pro Concentrate diluido. Deje secar al aire. No utilice paños. Siempre utilice Iodine Pro Concentrate fresco y diluido. No deje las vacas en un clima helado antes de que Iodine Pro Concentrate diluido se haya secado por completo.
Uso Expandido: Al refrescar a las vacas, comience con un baño antiséptico de sus pezones dos veces por día por aproximadamente 10 días antes de parir.
PRECAUCIÓN: Iodine Pro Concentrate no está diseñado para curar o ayudar con la cicatrización de los pezones agrietados o irritados. Como cualquier otro germicida, puede provocar irritación o sensibilización en animales sensibles. En caso de irritación o agrietamiento de los pezones, haga examinar dichas afecciones y, de ser necesario, hágaslas tratar con un veterinario.
AVISO: Utilice Iodine Pro Concentrate mezclado dentro de los seis (6) meses a partir de la fecha de mezclado.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
NO ES PARA USA HUMANO
Atención
 Sólo de uso externo
 Lavarse las manos cuidadosamente después de la manipulación.
 Consultar a un médico si la persona se encuentra mal.
LEA LA FICHA DE DATOS DE SEGURIDAD ANTES DE USAR EL PRODUCTO
INFORMACIÓN DE EMERGENCIA SOBRE SALUD: contacto local de veneno.

208 L (55 US GAL)
6302354



Manufactured for:
 Aspen Veterinary Resources*, Ltd.
 3155 Heartland Drive, Liberty, MO 64068
 1-800-203-5620
 778199/5300/1021

IODINE PRO CONCENTRATE

iodine solution, concentrate

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:46066-427
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	40 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46066-427-17	56800 mL in 1 DRUM		
2	NDC:46066-427-21	208000 mL in 1 DRUM		
3	NDC:46066-427-44	1134000 mL in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		11/23/2021	

Labeler - Aspen Veterinary (627265361)

Revised: 3/2025

Aspen Veterinary