

## **NICOTINE POLACRILEX MINT- nicotine polacrilex gum, chewing Walgreens**

-----  
**Nicotine Polacrilex Gum Mint 2 mg and 4 mg - Walgreens**

### ***Drug Facts***

#### ***Active ingredient (in each chewing piece) - 2 mg***

Nicotine polacrilex 2 mg (nicotine)

#### ***Active ingredient (in each chewing piece) - 4 mg***

Nicotine polacrilex 4 mg (nicotine)

### ***Purpose***

Stop smoking aid

### ***Use***

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

### ***Warnings***

**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

### **Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

### **Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may

need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.**

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions - 2 mg**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

<b>Weeks 1 to 6</b>	<b>Weeks 7 to 9</b>	<b>Weeks 10 to 12</b>
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider

**Directions - 4 mg**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

<b>Weeks 1 to 6</b>	<b>Weeks 7 to 9</b>	<b>Weeks 10 to 12</b>
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

***Other information - 2 mg***

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

***Other information - 4 mg***

- **each piece contains:** calcium 120 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

***Inactive ingredients - 2 mg***

carnauba wax, flavors, gum base, magnesium oxide, sodium bicarbonate, sodium carbonate, talc and xylitol.

***Inactive ingredients - 4 mg***

carnauba wax, flavors, gum base, magnesium oxide, sodium bicarbonate, sodium carbonate, talc and xylitol.

## Questions or comments?

Call 1-**877-753-3935** Monday-Friday 9AM-5PM EST

## Principal display panel 2 mg

Compare to the active ingredient in Nicorette® Gum††

UNCOATED

Nicotine Gum

NICOTINE POLACRILEX GUM 2 mg (NICOTINE)

STOP SMOKING AID

Sugar Free

2 mg

Mint Flavor

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP,

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Gum, 4 mg.

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks
3. Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. Use with a support program as directed in the enclosed User's Guide.
  - NOT FOR SALE TO THOSE UNDER 18 YEARS OF AGE
  - PROOF OF AGE REQUIRED
  - NOT FOR SALE IN VENDING MACHINES OR FROM ANY SOURCE WHERE PROOF OF AGE CANNOT BE VERIFIED

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

††This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., owner of the registered trademark Nicorette®.

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

**Package label 2 mg**

**TO INCREASE YOUR SUCCESS  
IN QUITTING:**

1. You must be motivated to quit.
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. Use with a support program as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing starting at corner with loose edge.



Push gum through foil.

For tools, resources and motivation, please visit  
**Walgreens.com/QuitSmoking**

NDC 0365-1405-08

**Walgreens**



Compare to the active ingredient in Nicorette® Gum™

UNCOATED  
**Nicotine Gum**

NICOTINE POLACRILEX GUM 2 mg (NICOTINE)  
STOP SMOKING AID

Sugar Free

**170** PIECES **2** mg

2 mg EACH



ACTUAL SIZE



Mint flavor

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.  
If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Gum, 4 mg.

THEFT SURVEILLANCE TAG AREA




how2recycle.info

QUESTIONS OR COMMENTS? Call 1-877-759-3828. Member Since 1960. Walgreens.com

**Directions**  
If you are under 18 years of age, ask a doctor before use.  
Before using this product, read the enclosed Patient Safety Information.  
Directions and use instructions are printed on the inner tray.  
If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum.  
If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum.  
Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to cause drowsiness and dizziness. When used, pieces of gum may stick to clothing and hair. Wash hands with soap and water after use.  
Contact a Patient Care Center (1-800-222-1222) right away if you have symptoms of an allergic reaction (such as difficulty breathing or rash), dizziness, nausea, and rapid heartbeat.

Walgreens.com ©2021 Walgreen Co.  
**100% SATISFACTION GUARANTEED**

ITEM 605048 W10024-1022-L  
PLD-0618A FC008323  
W30R00122-F REV-0922  
PRODUCT OF IRELAND



## WALGREENS Uncoated Nicotine Gum Mint Flavor

### Principal display panel 4 mg

Compare to the active ingredient in Nicorette® Gum††

UNCOATED

Nicotine Gum

NICOTINE POLACRILEX GUM 4 mg (NICOTINE)

STOP SMOKING AID

Sugar Free

4 mg

Mint Flavor

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex Gum, 2 mg.

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks
3. Use Long Enough - Use Nicotine Polacrilex Gum for the first full 12 weeks.
4. Use with a support program as directed in the enclosed User's Guide.
  - NOT FOR SALE TO THOSE UNDER 18 YEARS OF AGE
  - PROOF OF AGE REQUIRED
  - NOT FOR SALE IN VENDING MACHINES OR FROM ANY SOURCE WHERE PROOF OF AGE CANNOT BE VERIFIED

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN

††This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., owner of the registered trademark Nicorette®.

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOTRD., DEERFIELD, IL 60015

**Package label 4 mg**



**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing starting at corner with loose edge.



Push gum through foil.

For tools, resources and motivation, please visit [Walgreens.com/QuitSmoking](http://Walgreens.com/QuitSmoking)

NDC 0363-1401-06

**Walgreens**



Compare to the active ingredient in Nicorette® Gum\*

UNCOATED

# Nicotine Gum

NICOTINE POLACRILEX GUM 4 mg (NICOTINE)  
STOP SMOKING AID

Sugar Free

**170** **4 mg**  
PIECES

4 mg EACH



ACTUAL SIZE



Mint flavor

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.  
If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex Gum, 2 mg.

THEFT SURVEILLANCE TAG AREA



**Active Ingredients** Contains weak, flavor, gum base, polyvinyl alcohol, sodium bicarbonate, sodium carbonate, zinc stearate.

**Questions or comments?** Call 1-877-755-8285 Monday-Friday 9AM-9PM EST.

**Directions:** If you are under 18 years of age, ask a doctor before use. Do not use Nicotine Polacrilex Gum for longer than 12 weeks. Do not use Nicotine Polacrilex Gum if you are pregnant or breastfeeding. Do not use Nicotine Polacrilex Gum if you are allergic to any of the ingredients. Do not use Nicotine Polacrilex Gum if you are taking any other medications. Do not use Nicotine Polacrilex Gum if you are taking any other tobacco products. Do not use Nicotine Polacrilex Gum if you are taking any other nicotine products. Do not use Nicotine Polacrilex Gum if you are taking any other smoking cessation products. Do not use Nicotine Polacrilex Gum if you are taking any other smoking cessation products. Do not use Nicotine Polacrilex Gum if you are taking any other smoking cessation products.


DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015

**100% SATISFACTION**  
Guaranteed

Walgreens.com ©2021 Walgreen Co.

ITEM 609044 W10024-1022-L  
PLD-D624A FC008324  
W30H0122-F REV-0922

PRODUCT OF IRELAND



## WALGREENS Uncoated Nicotine Gum Mint Flavor

### NICOTINE POLACRILEX MINT

nicotine polacrilex gum, chewing

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-1421
<b>Route of Administration</b>	BUCCAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

#### Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
MAGNESIUM OXIDE (UNII: 3A3U0G171G)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
TALC (UNII: 7SEV7J4R1U)	
XYLITOL (UNII: VCQ006KQ1E)	

#### Product Characteristics

<b>Color</b>	white (off-white to tan)	<b>Score</b>	no score
<b>Shape</b>	SQUARE	<b>Size</b>	14mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1421-98	170 in 1 CARTON	01/01/2014	

1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076568	01/01/2014	

## NICOTINE POLACRILEX MINT

nicotine polacrilex gum, chewing

<b>Product Information</b>				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-1420	
Route of Administration	BUCCAL			
<b>Active Ingredient/Active Moiety</b>				
Ingredient Name	Basis of Strength	Strength		
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg		
<b>Inactive Ingredients</b>				
Ingredient Name	Strength			
CARNAUBA WAX (UNII: R12CBM0EIZ)				
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)				
SODIUM BICARBONATE (UNII: 8MDF5V39QO)				
SODIUM CARBONATE (UNII: 45P3261C7T)				
TALC (UNII: 7SEV7J4R1U)				
XYLITOL (UNII: VCQ006KQ1E)				
<b>Product Characteristics</b>				
Color	white	Score	no score	
Shape	SQUARE	Size	14mm	
Flavor	MINT	Imprint Code		
Contains				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1420-98	170 in 1 CARTON	01/01/2014	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076569	01/01/2014	

**Labeler** - Walgreens (008965063)

Revised: 10/2022

Walgreens